



Effect of Boron and Bio-fertilizers on Growth and Yield of Sesame (*Sesamum indicum* L.)

**Kanakapudi Bhavana^{a≡*}, Vikram Singh^{b∞}, Shruti Grace George^{a#}
and Shrish Kumar Singh^{ct}**

^a Department of Agronomy, NAI, SHUATS, Prayagraj-211007, (U.P.), India.

^b Directorate of Seed and Farm, NAI, SHUATS, Prayagraj-211007, (U.P.), India.

^c Department of Agronomy, TDPG College, Jaunpur, (U.P.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i1631025

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86441>

Received 14 February 2022

Accepted 22 April 2022

Published 27 April 2022

Original Research Article

ABSTRACT

A field experiment was conducted during *kharif* 2021 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The experimental soil used in this experiment was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.36 %), available N (171.48 kg/ha), available P (15.2 kg/ha) and available K (232.5 kg/ha). The experiment was laid out in Randomized Block Design with ten treatments each replicated three times on the basis of one year experimentation. The treatments which were Control (RDF), 2.5 kg B/ha + PSB, 2.5 kg B/ha + *Azotobacter*, 2.5 kg B/ha + PSB + *Azotobacter*, 5.0 kg B/ha + PSB, 5.0 kg B/ha + *Azotobacter*, 5.0 kg B/ha + PSB + *Azotobacter*, 7.5 kg B/ha + PSB, 7.5 kg B/ha + *Azotobacter* and 7.5 kg B/ha + PSB + *Azotobacter* are used. The results showed that application of 7.5 kg B/ha + PSB + *Azotobacter* showed significantly higher plant height (85.36 cm), No. of Branches/plant (6.57), Plant dry weight (19.53 g/plant), Crop growth rate (20.89 g/m²/day), capsules/plant (52.57), Seeds/capsule (62.60), Test weight (3.38 g), Seed yield (1.1635 t/ha), Stover yield (3.3819 t/ha), Biological yield (4.5556 t/ha), gross returns (Rs.79723.65/ha), net return (Rs.52849.25/ha) and

[≡]M.Sc. Scholar;

[∞]Associate Professor;

[#]Ph.D. Scholar;

[†]Associate Professor & Head;

*Corresponding author: E-mail: bhavanakanakapudi98@gmail.com;

benefit cost ratio (1.96) as compared to other treatments. Therefore, treatment with application of 7.5 B/ha along with PSB 10 g/kg seeds and Azotobacter 10 g/kg seeds was more productive and can be recommended to farmers after further trials.

Keywords: Azotobacter; boron; economics; growth; PSB; sesame; yield.

1. INTRODUCTION

“Among the oilseed crops, sesame (*Sesamum indicum* L.) is one of the world most important and oldest known oil seed crops (Abou Gharbia et al., 1997). Sesame is a member of the padaliaceae plant family. Its cultivation has started since 1500 BC in the Middle East, Asia & Africa” [1]. “India is the major producer of sesame (*Sesamum indicum* L.) and ranks first in both area (1.78 M ha) and production (0.81 Mt) with average productivity of 455 kg ha⁻¹. Odisha produces 0.09 million tonnes sesame seeds annually with average productivity of 403 kg ha⁻¹” [2]. Higher nutritional, medicine and cooking quality has recognized it as ‘the queen of oilseeds’. There is a decline in productivity in sesame due to its cultivation in marginal and sub-marginal lands and moreover poor crop management practice.

The sesame seed has been considered as ‘Queen of Oilseed’ for its high oilseed content and quality and traditionally categorized as a health food in China, Japan and Asian countries. It is considered to have both nutritional and medicinal value. “Moreover, seed is a rich source of edible oil (48-55%) and protein (20-28%) with anti-oxidants lignans such as sesamol and sesamin which prevents rancidity and gives sesame oil a shelf life. Sesame oil is called ‘poor man’s ghee’. The lignin content has useful physiological effect in human and animal health” [3]. The seeds are very rich in iron, magnesium, manganese, copper and calcium and contain vitamin E, A, B and B1. The seed contain phytosterols associate with reducing the level of blood cholesterol and also phytoestrogens with anti- oxidants an anticancer property.

“Among the micronutrient deficiency, boron deficiency is the second most dominant problem globally. Among the micronutrients, boron deficiency is one of the most widespread micronutrient deficiency in India. Plants require boron for a number of growth processes like development of meristematic tissue, proper pollination and seed set, translocation of sugars, starches. It has been reported that boron is

required for pollen germination and pollen tube growth” [4]. “Boron deficiency can affect pollen viability and abortion of stamen and pistils which contribute to low seed set reported by” [5].

“Biofertilizers like Azotobacter and phosphate solubilizing bacteria play a vital role in the nutrition of plants. Indian soils are mostly deficient in microorganisms and nutrients. So application of biofertilizers and inorganic fertilizers are necessary to increase the crop productivity” [6]. “Application of Azotobacter and Azospirillum incubated with FYM with 50 per cent moisture in soil resulting 13 and 14 times increase in population respectively, as compare to initial microbial population and increase in yield by 46 per cent in okra” [7]. “Growth and yield of crop is enhanced by the presence of sufficient quantities of available form of nutrients in the soil for plant uptake. So application of organics like FYM, biofertilizers and chemical fertilizers in integrated manner is beneficial to achieve higher seed yield and yield attributes” (Gablw et al., 2008).

2. MATERIALS AND METHODS

The present examination was carried out during Kharif 2021 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, UP, which is located at 25.28°N latitude, 81.54°E longitude and 98 m altitude above the mean sea level. The experiment was laid out in Randomized Block Design which consist of nine treatments with Control (RDF), 2.5 kg B/ha + PSB, 2.5 kg B/ha + Azotobacter (*Azotobacter beijerinckii*), 2.5 kg B/ha + Phosphate solubilizing bacteria (*Pseudomonas putida*) + Azotobacter, 5.0 kg B/ha + PSB, + Azotobacter, 5.0 kg B/ha + PSB + Azotobacter, 7.5 kg B/ha + PSB, 7.5 kg B/ha + Azotobacter and 7.5 kg B/ha + PSB + Azotobacter are used. The experimental site was uniform in topography and sandy loam in texture, nearly neutral in soil reaction (p^H -7.1), low in Organic carbon (0.38%), medium available N (225 kg ha⁻¹), higher available P (19.50 kg ha⁻¹) and medium available K (213.7 kg ha⁻¹). In the period from germination to harvest several plant growth parameters were recorded at frequent intervals along with it after

harvest several yield parameters were recorded those parameters are growth parameters, plant height, no. of branches/plant and plant dry weight are recorded. The yield parameters like No. of capsules/plant, No. of seeds/capsules, Test weight (g), seed yield, stover yield and harvest index were recorded and statistically analyzed using analysis of variance (ANOVA) as applicable to Randomized Block Design (Gomez and Gomez 1984).

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

3.1.1 Plant height

At harvest, significantly highest plant height (85.36 cm) was recorded in the treatment 7.5 kg B/ha + PSB + *Azotobacter* over all the other treatments. However, the treatments with application of 5.0 kg B/ha + PSB + *Azotobacter* (85.13 cm) and 7.5 kg B/ha + *Azotobacter* (85.27 cm) which were found to be at par with treatment 7.5 kg B/ha + PSB + *Azotobacter* as compared to all the treatments. The increase in plant height might be due to the involvement of Boron in different physiological processes like enzymes activation, electron transport, chlorophyll formation, stomatal regulation, etc. which gradually increased plant height. The results were found in accordance with Choudhary et al. [8]. The increase in plant height due to the application of PSB might be due to the presence of plant growth hormones i.e. IAA, cytokinin, gibberellins, and phosphorus availability to the plant similar results were found by Dutta and Singh [9].

3.1.2 Number of branches/plant

At harvest, maximum Number of Branches/plant (6.57) was observed in the treatment with 7.5 kg B/ha + PSB + *Azotobacter*. However, treatment with 5.0 kg B/ha + *Azotobacter* (6.52) and 7.5 kg B/ha + PSB (6.41) which was found to be statistically at par with 7.5 kg B/ha + PSB + *Azotobacter*.

The higher levels of boron attributed to best results because boron might attributed to the favorable influence of them on plant metabolism and biological process activity and their stimulating effect on photosynthetic pigments and enzyme activity which in turn encouraged vegetative growth. The results were found similar with Ravichandra et al. [10].

3.1.3 Plant dry weight (g/plant)

At harvest, treatment with 7.5 kg B/ha + PSB + *Azotobacter* was recorded with significantly maximum dry weight (19.53 g/plant) over all the treatments. However, the treatments 7.5 kg B/ha + PSB (19.48 g/plant) and 5.0 kg B/ha + PSB + *Azotobacter* (19.42 g/plant) were found to be statistically at par with 7.5 kg B/ha + PSB + *Azotobacter*.

The application of boron generally influences cell division and nitrogen absorption from soil might enhanced plant growth reflects in terms of plant dry weight. "The findings were in harmony with Mamatha et al. [11]. Inoculation of PSB solubilization of inorganic insoluble phosphates by microorganisms to the production of organic acids, chelating oxoacids from sugars, and exchange reactions in growth environment, the results were found to similar with" Kumar et al. [12].

3.2 Yield Parameters

3.2.1 Number of capsules/plant

Significantly Maximum Number of capsules/plant (52.57) was recorded with the treatment of application of 7.5 kg B/ha + PSB + *Azotobacter* over all the treatments. However, the treatments 5.0 kg B/ha + PSB + *Azotobacter* (51.96) and 7.5 kg B/ha + PSB (52.25) which were found to be statistically at par with 7.5 kg B/ha + PSB + *Azotobacter*.

Significant increase in number of capsules/plant is due to increase in the availability of Nitrogen and Phosphorous through bio fertilizer inoculation by which more spikelets are produced due to increased rates of spikelets primordial production, similar results were found Hadiyal et al. [13].

3.2.2 Number of seeds/capsules

Significantly Maximum Number of Seeds / capsule (62.60) was recorded with the treatment of application of 7.5 kg B/ha + PSB + *Azotobacter* over all the treatments. However, the treatments 5.0 kg B/ha + PSB + *Azotobacter* (61.87) and 7.5 kg B/ha + PSB (62.22) which were found to be statistically at par with 7.5 kg B/ha + PSB + *Azotobacter*.

"The application of boron to sesame generally improves capsule growth by synthesizing

tryptophan and auxin. The enhancement effect on capsules/plant, seeds per capsule attributed to the favorable influence of boron application to crops on nutrient metabolism, biological activity and growth parameters which in turn influenced higher enzyme activity which in turn encouraged more capsules/plant and seeds/capsule. Similar findings were observed by” Yadav et al. [14].

3.2.3 Test weight (g)

Significantly highest Test weight (3.38 g) was recorded with the treatment of application of 7.5 kg B/ha + PSB + *Azotobacter* over all the treatments. However, the treatments 5.0 kg B/ha + PSB + *Azotobacter* (3.29 g) and 7.5 kg B/ha + PSB (3.35 g) which were found to be statistically at par with 7.5 kg B/ha + PSB + *Azotobacter*.

3.2.4 Seed yield (t/ha)

Significantly high seed yield (1.1635 t/ha) was recorded with the treatment (7.5 kg B/ha + PSB + *Azotobacter*) over all the treatments. However, the treatments with (1.1141 t/ha) in 5.0 kg B/ha + PSB + *Azotobacter* and with (1.1447 t/ha) in 7.5 kg B/ha + PSB which were found to be statistically at par with 7.5 kg B/ha + PSB + *Azotobacter*.

“Boron plays the vital role in increasing because it takes place in many physiological process of plant such as chlorophyll formation, stomatal

regulation, starch utilization which enhanced seed yield”, Mallik and Raj (2015).

3.2.5 Stover yield (t/ha)

Significantly high stover yield (3.3819 t/ha) was recorded with the treatment (7.5 kg B/ha + PSB + *Azotobacter*) over all the treatments. However, the treatments with (3.2999 t/ha) in 5.0 kg B/ha + PSB + *Azotobacter* and 7.5 kg B/ha + PSB (3.3443 kg/ha) which were found to be statistically at par with 7.5 kg B/ha + PSB + *Azotobacter*.

“Increase in yield and stover yield through bio-fertilizer might be attributed to supply of more plant hormones (auxin, cytokinin, gibberellin etc.) by the microorganisms inoculated or by the root resulting from reaction to microbial population and more availability of phosphorous and nitrogen. Similar results were obtained by” Kalita et al. (2019).

3.3 Harvest Index (%)

Significantly high harvest index (26.1%) was recorded with the treatment (2.5 kg B/ha + PSB + *Azotobacter*) over all the treatments. However, the treatments with (25.7%) in 5.0 kg B/ha + PSB + *Azotobacter* and 7.5 kg B/ha + *Azotobacter* (26.0%) which were found to be statistically at par with 7.5 kg B/ha + PSB + *Azotobacter*.

Table 1. Effect of boron and bio-fertilizers on growth of cowpea

Treatments	At harvest			30-45 DAS	
	Plant height (cm)	Number of branches / plant	Dry weight (g)	CGR (g/m ² /day)	RGR (g/g/day)
Control (RDF)	83.78	5.65	18.26	20.49	0.061
2.5 kg B/ha + PSB	83.83	5.31	18.26	19.82	0.057
2.5 kg B/ha + <i>Azotobacter</i>	84.05	6.15	18.31	19.60	0.057
2.5 kg B/ha + PSB + <i>Azotobacter</i>	84.49	5.45	18.35	20.02	0.056
5.0 kg B/ha + PSB	84.63	5.43	18.37	20.56	0.058
5.0 kg B/ha + <i>Azotobacter</i>	84.43	6.52	18.62	19.78	0.056
5.0 kg B/ha + PSB + <i>Azotobacter</i>	85.13	5.27	19.42	20.00	0.054
7.5 kg B/ha + PSB	85.27	6.41	19.48	19.82	0.052
7.5 kg B/ha + <i>Azotobacter</i>	84.89	6.29	19.32	20.24	0.057
7.5 kg B/ha + PSB + <i>Azotobacter</i>	85.36	6.57	19.53	20.89	0.052
F test	S	S	S	S	NS
Sem±	0.13	0.16	0.06	0.21	0.00
CD (P=0.05)	0.39	0.56	0.17	0.62	-

Table 2. Effect of boron and bio - fertilizers on yield of sesame

Treatments	At Harvest						
	Number of capsules / plant	Number of seeds / capsules	Test weight (g)	Seed yield (t/ha)	Stover yield (t/ha)	Biological Yield (t/ha)	Harvest index (%)
Control (RDF)	47.5	54.6	2.89	0.75	2.23	2.99	24.7
2.5 kg B/ha + PSB	48.2	56.3	3.05	0.84	2.58	3.42	24.5
2.5 kg B/ha + <i>Azotobacter</i>	47.8	55.0	3.01	0.78	2.34	3.14	24.9
2.5 kg B/ha + PSB + <i>Azotobacter</i>	48.9	57.7	3.08	0.99	2.81	3.79	26.1
5.0 kg B/ha + PSB	50.3	58.2	3.19	1.05	3.03	4.06	25.7
5.0 kg B/ha + <i>Azotobacter</i>	48.5	57.1	3.01	0.91	2.73	3.64	24.8
5.0kg B/ha + PSB + <i>Azotobacter</i>	52.0	61.9	3.28	1.12	3.29	4.46	25.0
7.5 kg B/ha + PSB	52.3	62.2	3.35	1.15	3.35	4.52	25.3
7.5 kg B/ha + <i>Azotobacter</i>	51.2	60.5	3.23	1.09	3.11	4.18	26.0
7.5 kg B/ha + PSB + <i>Azotobacter</i>	52.6	62.6	3.38	1.17	3.39	4.56	25.5
F test	S	S	S	S	S	S	S
Sem±	0.22	0.32	0.04	0.02	0.16	0.16	0.12
CD (P=0.05)	0.66	0.94	0.13	0.06	0.49	0.49	0.33

4. CONCLUSION

It is concluded that application of treatment 7.5 kg B/ha+ PSB10 g/kg seeds + *Azotobacter* 10 g/kg seeds (T10) had performed better in growth and yield parameters. As it was more productive it can be recommended to farmers after further trails.

ACKNOWLEDGEMENTS

The work was supported by my Advisor Dr. Vikram Singh and all faculty members, Department of Agronomy, Naini Agricultural institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, (U.P.) India for providing field and assistance in conducting the research trail.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ali GM, Yasumoto S, Seki-Katsuta M. Assessment of genetic diversity in Sesame (*Sesamum indicum* L.) detected by Amplified Fragment Length Polymorphism Markers. Electronic Journal of Biotechnology. 2007;10:12-23.
2. Anonymous. Odisha Agriculture Statistics 2012-13. Directorate of Agriculture and Food production, Bhubaneswar, 2014, Odisha.
3. Mane SG. Studies on growth, yield and quality of sesame (*Sesamum indicum*) as influenced by chemical fertilizers and liquid bio fertilizers. Thesis, M.Sc. (Agri.), VNMKV, Parbhani; 2019.
4. Dugger WM. Boron in plant metabolism. Encyclopedia of Plant Nutrition. 1983;626-650.
5. Chitralekha C, Nirmala N. Developmental aberrations in seeds of boron deficient

- sunflower and recovery. Journal of Plant Nutrition. 2000;6:835-841.
6. Patel HA, Raj AD. Nutrient content as well as uptake of summer sesame as affected by nitrogen, phosphorus and biofertilisers under south Gujarat conditions. International Journal of Chemical Studies. 2017;5(6):01-04.
 7. Pattanayak SK, Mohanty RK, Sethy AK. Response of Okra to Azotobacter and Azospirillum inoculation grown in Acid soil amended with lime and FYM. II North Eastern Regional Conference on Biofertilisers, AAU, Jorhat, Assam; 2001.
 8. Choudhary Md. MU, Ullah Md. H, Rahman Md. A, Shahidullslam. Effect of boron and nitrogen fertilization on cowpea growth, nodulation and grain yield in Rangamati, Bangladesh. Legume Research. 2000; 23(1):9-14.
 9. Dutta S, Singh MS. Effect of Azotobacter on yield and oil content of rapeseed mustard varieties under Manipur condition. Indian Journal of Hill Farming. 2002;15(2): 44-46.
 10. Ravichandra K, Nagajyothi B, Jaipal S, Joy D, Krupakar A. Growth of groundnut (*Arachis hypogea* L.) and its yield as influenced by foliar spray of Boron along with rhizobium inoculation. Indian Journal of Dryland Agriculture and Development. 2015;30(1):60-63.
 11. Mamatha K, Vidya S, Laxmi N. Influence of different levels of boron along with application of FYM on nutrient uptake and yield of Sesame (*Sesamum indicum* L.). International Journal of Farm Sciences. 2016;6(2):37-45.
 12. Kumar V, Singh G, Shrivastava AK, Singh VK, Singh RK, Kumar A. Effect of integrated nutrient management on growth and yield attributes of Indian mustard. Indian Journal of Ecology. 2016;43(1): 440-443.
 13. Hadiyal JG, Kachhadiya SP, Ichchuda PK, Kalsaria RN. Response of Indian mustard (*Brassica juncea* L.) to different levels of organic manures and bio-fertilizers. Journal of Pharmacognosy and Phytochemistry; 2017.
 14. Yadav SN, Singh SK, Omkar K. Effect of Boron on Yield Attributes, Seed Yield and Oil Content of Mustard (*Brassica juncea* L.) on an Inceptisol. Journal of the Indian Society of Soil Science. 2016;64(3):291-296.

© 2022 Bhavana et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/86441>