



Benefit of Early and Goal Directed Physiotherapy Intervention on Gross Motor Function in a Child with Japanese Encephalitis: A Rare Case Report

**Riya Kale ^{a≡}, Rakesh Krishna Kovela ^{a*ϕ}, Mohammed Irshad Qureshi ^{a#},
Ragini Dadgal ^{a†}, Rebecca Timothy ^{b†} and Snehal Samal ^{c†}**

^a Department of Neurophysiotherapy, Ravi Nair Physiotherapy College, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), Wardha, Maharashtra - 442001, India.

^b Department of Electrotherapy, Ravi Nair Physiotherapy College, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), Wardha, Maharashtra - 442001, India.

^c Department of Kinesiology and Kinesiotherapy, Ravi Nair Physiotherapy College, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), Wardha, Maharashtra - 442001, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2022/v34i2A35358

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/76659>

Case Study

Received 15 October 2021
Accepted 22 November 2021
Published 16 January 2022

ABSTRACT

Due to its widening geographical range, Japanese encephalitis (JE) has lately been named a notifiable disease in India. The disease notification makes it easier to put preventive measures in place and manage cases. JE is a vector-borne disease that is preventable with vaccination. The virus that causes it is the Japanese encephalitis virus (JEV), which belongs to the Flaviviridae family. The present case report is of an infant of 18 months old presented with problems of decreased tone, loss of developmental achieved milestones and convulsions with severe fever. On examination there was hypotonia, lack on neck control, difficulty in rolling, low pitch cry and smile. We managed the child with standard physiotherapy protocol. We framed goals focusing both on

[≡]BPT Student

^ϕAssociate Professor

[#]Professor

[†]Assistant Professor

*Corresponding author: E-mail: rakeshrishna.pt@gmail.com;

preventing complications and promoting recovery. We gave treatment for a period of 4 weeks. We measured Modified Ashworth Scale (MAS) and Gross motor function measure (GMFM-88) Pre and post treatment. We would like conclude our case report by mentioning the importance early and goal specific supportive and restorative care to a child will enable him or her to take more vigorous physiotherapy care in future for restoring sensorimotor and behavioral function.

Keywords: Japanese encephalitis; physiotherapy; rehabilitation; goal oriented therapy; early physiotherapy.

1. INTRODUCTION

Japanese encephalitis is a viral infection. The virus that causes Japanese encephalitis, Japanese Encephalitis virus is spread through mosquito bites. It causes significant brain inflammation. With an estimated 50 000 cases and 15 000 deaths per year, Japanese encephalitis is one of the most common types of viral encephalitis globally [1]. The disease is prevalent in larger parts of Asia. A third of patients die, and half of those who survive have significant neuropsychiatric complications [2].

Encephalitis epidemics have been documented in Japan since the early 1870s. Initially termed as type B encephalitis. Later, the virus was classified as a member of the Flavivirus genus (family Flaviviridae), which was named after the prototype yellowfever virus (Latin: yellow=flavi) [1].

JEV has a 5–15 day incubation period. The clinical disease ranges from a vague febrile sickness to a severe condition characterised by meningoencephalitis, aseptic meningitis, or sudden flaccid paralysis similar to polio [3].

The virus now has a well-established zoonotic transmission cycle involving mosquitoes, pigs, and/or aquatic fowl; humans become infected only by chance when bitten by an infected mosquito and serve as a dead-end host [4].

Japanese encephalitis usually manifests in initial stages as non-specific feverish sickness, which may include coryza, diarrhoea, and rigours. This is followed by a headache, vomiting, and a loss of consciousness, which is frequently preceded by a convulsion [1].

The haemagglutination inhibition test IgM and IgG capture enzyme linked immunosorbant assays (ELISAs) The IgM ELISA has recently been changed to a simple nitrocellulose membrane-based format, with a color change discernible to the naked eye [5].

Limiting outside exposure at dusk and dawn, dressing in clothing that exposes the least amount of skin possible, Japanese encephalitis vaccine and using insect repellents are possible measures to prevent infection [1].

Japanese encephalitis is treated with supportive care, which includes treating convulsions and high intracranial pressure when they occur. Corticosteroids have been used for many years, but an RCT of dexamethasone found no effect [6].

Physiotherapy till date is supportive. Airway, Breathing, Circulation has to be maintained. Positioning plays a vital role to prevent pressure sores [7].

1.1 Aim

To investigate effect of early and goal directed intervention in managing child with Japanese encephalitis.

2. PATIENT CHARACTERISTICS

The present case report is of an infant of 18 months old presented with problems of decreased tone, loss of developmental achieved milestones and convulsions with severe fever. Child was irritated while assessment. Later cooperated with humming music and rhymes. On examination there was hypotonia, lack on neck control, difficulty in rolling, low pitch cry and smile (Fig. 1).

2.1 Patient History

Child was apparently normal till August 2021. Suddenly developed acute pyrexia with diarrhea. Parents immediately rushed the child to nearby hospital where antipyretic medications were given. As the fever was not subsiding, he was referred to our hospital where he was kept in ICU. He was given airway support. During the period of stay in intensive care unit child

developed few episodes of convulsions for which medications were given and they came under control in one week with supportive therapy. After

a stay of 10 days in ICU child was shifted to general ward. Physiotherapy was started from ICU itself.



Fig. 1. Depicting child with Japanese Encephalitis

Table 1. Depicting the Physiotherapy protocol for 4 Weeks

Problem identified	Cause of the problem	Goal	Treatment strategy	Equipment used
Hypotonia	Inflammation of nervous system and decreased input from brain	To develop tone in postural and phasic muscles	Facilitators techniques for Postural and phasic muscles	Bolsters, physioballs, Lap therapy
Lack of head and neck control	Inflammation of nervous system and decreased input from brain	To enhance head and neck control	Lap therapy, vestibular stimulation	Physioballs, wedges and bolsters
Lack of efficient rolling		Promote segmental rolling	Segmental rolling facilitation on physioball	Physioballs, wedges and bolsters
Decreased pitch in cry	Inflammation of nervous system and decreased input from brain	Improve aeration of lungs	Chest Physiotherapy and positioning strategies	Direct handling
pressure sores		Inability to move because of low tone	Managing already developed pressure sores and preventing further sores	Frequent changing of positioning

Table 2. Depicting the Pre and post scores on day 1 and at the end of 4 Weeks

Outcome Measure	Pre-Test scores	Post test scores
MAS(TA, biceps, hip adductors) Rt and Lt sides	0	1
GMFM 88(Lying and rolling component)	09	34

2.2 Physiotherapy Management

Till date there is no specific physiotherapy protocol for Japanese encephalitis. Children with JE used to be managed in same line as encephalitis but the results were not supporting the recovery. We felt standard physiotherapy protocol is important in managing acute cases of JE. We framed goals focusing both on preventing complications and promoting recovery. Goals and goal specific management is given in (table 1). We gave treatment 45 min a day, 5 days a week for a period of 4 weeks. We measured Modified Ashworth Scale (MAS) and Gross motor function measure (GMFM-88) (8)Pre and post (Table 2).

3. RESULTS

Physiotherapy treatment was lasted for 4 weeks. Pre and post scores suggest there is improvement in child gross motor status and also tone. Chief muscles of upper and lower limb were taken into consideration. There was improvement in tone in all the three muscle groups bilaterally. Lying and rolling component was taken from GMFM 88 by keeping the condition of the child in mind.

4. DISCUSSION

Results of the present study depicts that goal oriented treatment definitely plays a major role in improving the child’s functional status. These results are in line previous study by kovala et al in their study on cerebral palsy child mentioned that goal specific management plays a vital role in improving functional status of children [9]. Results of this study are also in line with the previous study by Christy et al on encephalitis children [10]. Identifying the cause, planning goal specific strategies and regular physiotherapy starting from icu to opd helped the child in gaining some tone. Physiotherapy has to be continued for few more months to get some more improvement in the child. Mother has been also trained with specific home program using kovala etal study on mother as a rehabilitative aid [11].

5. CONCLUSION

We would like conclude our case report by mentioning the importance early and goal specific supportive and restorative care to a child will enable him or her to take more vigorous physiotherapy care in future for restoring sensorimotor and behavioral function. As there is scarcity of physiotherapy care in JE, more high-quality studies are required to support the benefit and importance of physiotherapy in early stages of disease and also on cognitive function.

CONSENT

As per international standard, parental written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Solomon T, Dung NM, Kneen R, Gainsborough M, Vaughn DW, Khanh VT. Japanese encephalitis. *J Neurol Neurosurg Psychiatry*. 2000 ;68(4):405–15.
- Mackenzie JS, Gubler DJ, Petersen LR. Emerging flaviviruses: the spread and resurgence of Japanese encephalitis, West Nile and dengue viruses. *Nat Med*. 2004;10(12):S98–109.
- Solomon T, Vaughn DW. Pathogenesis and Clinical Features of Japanese Encephalitis and West Nile Virus Infections. In: Mackenzie JS, Barrett ADT, Deubel V, editors. *Japanese Encephalitis and West Nile Viruses* [Internet]. Berlin, Heidelberg: Springer; 2002 [cited 2021 Oct

- 18]. p. 171–94. (Current Topics in Microbiology and Immunology). Available: https://doi.org/10.1007/978-3-642-59403-8_9
4. Endy TP, Nisalak A. Japanese Encephalitis Virus: Ecology and Epidemiology. In: Mackenzie JS, Barrett ADT, Deubel V, editors. Japanese Encephalitis and West Nile Viruses [Internet]. Berlin, Heidelberg: Springer; 2002 [cited 2021 Oct 18]. p. 11–48. (Current Topics in Microbiology and Immunology). Available: https://doi.org/10.1007/978-3-642-59403-8_2
 5. Solomon T, Thao LTT, Dung NM, Kneen R, Hung NT, Nisalak A, et al. Rapid Diagnosis of Japanese Encephalitis by Using an Immunoglobulin M Dot Enzyme Immunoassay. J Clin Microbiol. 1998 Jul 1;36(7):2030–4.
 6. Hoke CH, Vaughan DW, Nisalak A et al. Effect of High-Dose Dexamethasone on the Outcome of Acute Encephalitis Due to Japanese Encephalitis Virus | The Journal of Infectious Diseases | Oxford Academic [Internet]. [cited 2021 Oct 18]. Available: <https://academic.oup.com/jid/article-abstract/165/4/631/816103>
 7. Kumar R. Prevention, diagnosis, and management of Japanese encephalitis in children. Pediatr Health Med Ther. 2014;5:99–110.
 8. Michaelis U. Gross Motor Function Measure (GMFM-66 & GMFM 88) User's Manual 2nd Edition Clinics in Developmental Medicine Edited by Dianne J Russell, Peter L Rosenbaum, Marilyn Wright, Lisa M Avery London, UK: Mac Keith Press, 2013 £70.00 (Spiral Binding), pp 290 ISBN: 978-1-908316-88-2. Dev Med Child Neurol. 2015;57(12): 1188–1188.
 9. Kovala R, Sinha M. Goal directed Physiotherapy treatment Program for improving lower extremity function in a child with spastic paraplegic cerebral palsy. Indian J Public Health Res Dev. 2018;9.
 10. Christie S, Chan V, Mollayeva T, Colantonio A. Rehabilitation interventions in children and adults with infectious encephalitis: a systematic review protocol. BMJ Open. 2016;6(3): e010754.
 11. Jayashree K, Krishna KR, Thakur A, Priya PR. Effectiveness of Mother as a Rehabilitative Aid (MARA) Program in the Recovery of Children with Cerebral Palsy- An Assessor blinded Randomized Controlled Trial. Indian J Public Health Res Dev. 2020;11(6):419–24.

© 2022 Kale et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/76659>