



## **Prevalence of Helicobacter Pylori Gastritis among Patients with Symptomatic Cholelithiasis**

**Amir Iqbal Memon <sup>a#</sup>, Samina Naz <sup>a≡</sup>, Riaz Ahmed Memon <sup>a≡</sup>,  
Aisha Masroor Bhatti <sup>a\*⊖</sup>, Nayab <sup>a⊖</sup> and Mansoor Ali <sup>b†</sup>**

<sup>a</sup> Department of Surgery, Liaquat Univeristy of Medical & Health Sciences, Jamshoro, Pakistan.

<sup>b</sup> Department of Plastic Surgery, Liaquat Univeristy of Medical & Health Sciences, Jamshoro, Pakistan.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2022/v34i33A36120

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86421>

**Received 18 February 2022**

**Accepted 20 April 2022**

**Published 20 April 2022**

**Original Research Article**

## **ABSTRACT**

**Background:** H. Pylori gastritis is considered to be the causative factor for gallstone disease. Though this infection is reported to be the leading factor contributing to the pathogenesis of gastro-duodenal ulcers, and gastric cancer. Its infestation of the gallbladder and causing chronic cholecystitis and cholelithiasis are still debatable.

**Methodology:** This study was descriptive cross-sectional in nature, performed on 110 patients (chosen via non-probability, consecutive sampling), presented at Surgical OPD with symptomatic cholelithiasis. All the participants underwent for diagnostic endoscopy and a biopsy was taken for confirmation of h.Pylori gastritis. Those who had positive H. Pylori test, were treated by eradication therapy first and then submitted for the laparoscopic cholecystectomy and those who were negative submitted for the laparoscopic cholecystectomy. The study lasted 1 year from February 2021 to February 2022.

**Results:** Among the subjects, 72.7% (n=80) were female, 32.5% were under 40 years old while male gender composed of 17.3% of the sample population. 3/4<sup>th</sup> of the population hailed from the

<sup>#</sup> Professor & HOD;

<sup>≡</sup> Assistant Professor;

<sup>⊖</sup> Post Graduate Trainee;

<sup>†</sup> Senior Medical Officer;

\*Corresponding author: E-mail: aisha\_bhatti\_999@hotmail.com;

rural setting. 70.1% (n=78) of the patients had positive H. Pylori test on histopathology. Upon the findings of diagnostic endoscopy, severe and moderate gastritis was found to be in 25.6% and 74.4% of the participants with positive helicobacter gastritis, respectively. A significant association was found out between familial history of gall stones and helicobacter pylori gastritis (p=0.02).

**Conclusion:** A high prevalence of H. Pylori infection is among patients with symptomatic cholelithiasis, which also crossponds to the endoscopic findings. The association between the familial history of gall stones and helicobacter gastritis merits to be explore further to establish the casuality of the relationship.

**Keywords:** *H. pylori gastritis; symptomatic cholelithiasis; chronic cholecystitis; postcholecystectomy syndrome.*

## 1. INTRODUCTION

Helicobacter pylori is a gram -ve micro-organism [1] which affects about 50% of the world's population [2]. Almost 4.4 billion individuals suffered from H. pylori infection in 2015 [3]. H. Pylori affects the epithelial tissue of the stomach and may lead to gastric and duodenal ulcers [4] gastric carcinoma [5,6] gastric malignant lymphoma of the mucosa-associated lymphoid tissue (MALT) [7]. As the incidence of cholelithiasis is increasing among the adult population approximately 76.66 % of patients with symptomatic cholelithiasis were infected with the concomitant H. pylori gastritis as well [8]. Nowadays chronic cholecystitis and cholelithiasis establish serious health problems with an excessive medical burden with a prevalence of 11% to 36% [6,9]. H. Pylori could reach the gallbladder directly from the stomach or through the portal blood circulation. Hence, there might be a possibility of another underlying H. pylori infestation of gall bladder and pathophysiology behind the cholecystitis and gastritis [4,9]. As H. pylori along with the gut microbiome reduce bile acid metabolism and leads to gallstone formation [7,10].

Patients with cholelithiasis present most commonly with abdomen pain either in the right hypochondrium or at the epigastrium followed by heartburn, dyspepsia, and bloating [11,12]. The endoscopy findings of most patients with cholelithiasis were in favor of moderate to severe gastritis with a significant 3/4th ratio of cholelithiasis patients with other gastroduodenal problems. Cholecystectomy in patients with cholelithiasis along with concomitant H. pylori infection having nonspecific gastrointestinal symptoms is unjustifiable as the symptoms persist even after surgery and it will be miss-labeled as a case of the post-cholecystectomy syndrome [11]. The H. Pylori eradication therapy

benefits in decreasing the intensity of symptoms, especially the heartburn, and improves the quality of life for patients with the concomitant diseases even before submitting a cholelithiasis patient for laparoscopic cholecystectomy, as the surgery alone can't relieve the symptoms in this scenario [13]. This study was conducted with an aim to determine the prevalence of H. Pylori gastritis among patients presenting with symptomatic cholelithiasis.

### 1.1 Objective

To determine the prevalence of H. Pylori gastritis among patients presenting with symptomatic cholelithiasis.

## 2. METHODOLOGY

This was a cross-sectional descriptive study which was performed from February 2021 to February 2022, on 110 patients (chosen via non-probability, consecutive sampling), who presented at Surgical OPD with symptomatic cholelithiasis along with ultrasound abdomen pointing towards the stone (single/multiple) at the gallbladder without any sign of acute cholecystitis or choledocholithiasis. Informed written consent was taken from patients. Patients were asked to undergo diagnostic endoscopy. The biopsy was also carried out to evaluate the presence of H. Pylori gastritis. Those who came to be H. Pylori positive will be treated for 10-14 days and then submitted for surgery after resolving the gastritis symptoms. The patients with negative H. Pylori will be admitted to the Department of Surgery at Liaquat University of Medical & Health Sciences (LUMHS) Jamshoro and submitted for the surgery. The exclusion criteria were previous the use of H. pylori eradication therapy. Data were documented using a structured questionnaire, including inquiries related to sociodemographic details, and disease specifics, and observed for

the incidence of H. pylori gastritis among patients with symptomatic cholelithiasis. All the maneuvers (history taking, physical examination, sampling, and data collection) were done by the principal researcher while the data was collected on a pre-designed proforma. The data was analyzed using SPSS version 22.0 Chi-square test was applied to find the association familial history of gall stones & helicobacter gastritis.

### 3. RESULTS

Among the subjects, 72.7% (n=80) were female, 32.5% were under 40 years old while male gender composed of 17.3% of the sample population. The mean age of participants were 43.57 years ( $\pm$  7.54). 3/4<sup>th</sup> of the population hailed from the rural setting. 70.1% (n=78) of the patients tested positive for H. Pylori infection on histopathology. Gastritis was present in 93% of patients (n=102). Upon the findings of diagnostic endoscopy, severe and moderate gastritis was found to be in 25.6% and 74.4% of the participants with positive helicobacter gastritis, respectively. A significant association was found out between familial history of gall stones and helicobacter pylori gastritis (p=0.02).

### 4. DISCUSSION

The aim of this research was to find out H. pylori gastritis incidence in patients with symptomatic cholelithiasis. 70.1% of the patients with symptomatic cholelithiasis had helicobacter infection during histopathology examination. Nonetheless, the intensity of gastritis was not significantly associated with helicobacter gastritis. Family history of gallstone was also related to H. Pylori gastritis in the present research.

A research reported that people with symptomatic cholelithiasis had 37% more chances of having helicobacter gastritis than non-cholelithiasis patient, which is reported more in this study [14]. A study conducted by Zhang et al. found the 9.47% prevalence of cholelithiasis among helicobacter gastritis and 8.46% in non-helicobacter gastritis. They also reported a 9.02% prevalence gallstones in patients who have taken eradication therapy helicobacter [15]. Another study conducted by Sabbaghian et al. found 33.3% positivity rate of helicobacter pylori infection by polymerase-chainreaction who underwent cholecystectomy which was more in this research [16].

**Table 1. Descriptive statistics**

Variable		n (110)	%
<b>Mean Age</b>		<b>43.57 years (+ 7.54)</b>	
Gender	Male	30	17.3%
	Female	80	72.7%
Residence	Urban	24	11.9%
	Rural	86	78.1%
Number of gall stones	Single	63	57.3%
	Multiple	47	42.7%
H. Pylori status	Positive	78	70.1%
	Negative	32	29.9%
Severity of Gastritis	Mild	17	16.6%
	Moderate	62	60.8%
	Severe	23	22.6%

**Table 2. Association of Family History of Galls Stones & Severity of Gastritis with H. Pylori Infection**

Variable		H. Pylori Positive (78)	H. Pylori Negative (32)	P Value
Family History of Gall Stones	Positive	51 (65.4%)	17 (53.1%)	0.02
	Negative	27 (34.6)	15 (46.9%)	0.3
Severity of Gastritis	Mild	0 (0%)	17 (100%)	0.23
	Moderate	58 (74.4%)	4 (12.5%)	0.6
	Severe	20 (25.6%)	3 (9.38%)	0.07

A research upon 58 patients with cholelithiasis reported a total of 17 patients with helicobacter gastritis [17]. This is against the findings of our study. Monstein et al. found that around half of the patients i.e. 55%, suffering from cholesterol cholelithiasis were tested positive helicobacter pylori infection. This rate was lesser in comparison to our finding [18]. Takahashi et al. found the 3.81% incidence rate of cholelithiasis among patients with negative helicobacter status in comparison to 6.08% incident rate of cholelithiasis among positive cases. While those who have been had on eradication therapy, had 4.37% incident rate [19].

Similarly, a study reported a contributing rate of 63% of helicobacter gastritis, which is higher than our obtained results [20]. Overall, incidence of H. Pylori among symptomatic cholelithiasis patients, is reported to be high. The relation between familial history of cholelithiasis and helicobacter gastritis is of statistical significance.

## 5. CONCLUSION

A high prevalence of H. Pylori infection is among patients with symptomatic cholelithiasis, which also corresponds to the endoscopic findings. The association between the familial history of gall stones and helicobacter gastritis merits to be explored further to establish the causality of the relationship.

## CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Srigiri SK, Kanigiri L, Pavuluri N, Thota R. Role of Detection of Helicobacter Pylori in Cholecystectomy Specimens by Using Giemsa Stain. *Journal of Chalmeda Anand*

2. Rao Institute of Medical Sciences. 2020; 20(2):17.
2. Kotilea K, Bontems P, Touati E. Epidemiology, diagnosis and risk factors of Helicobacter pylori infection. *Helicobacter pylori in Human Diseases*. 2019;17-33.
3. Zhang F, Pu K, Wu Z, Zhang Z, Liu X, Chen Z, et al. Prevalence and associated risk factors of Helicobacter pylori infection in the Wuwei cohort of north-western China. *Tropical Medicine & International Health*. 2021;26(3):290-300.
4. Mahmood A, Khan Z, Razzaq S, Mahmood MA, Ahmed N, Iqbal W. The Likelihood of Helicobacter Pylori presence in pre Cholecystectomy gall bladder with or without Cholecystitis-A case control study. *Pak J Med Health Sci*. 2020;14(2):304-6.
5. Silva CS, da Silva Júnior RT, de Sá Santos LK, Apolonio JS, da Costa BT, Cuzzuol BR, et al. Extragastric Manifestations of Helicobacter Pylori Infection: A; 2021.
6. Santos MLC, de Brito BB, da Silva FAF, Sampaio MM, Marques HS, e Silva NO, et al. Helicobacter pylori infection: Beyond gastric manifestations. *World Journal of Gastroenterology*. 2020;26(28):4076.
7. Wang L, Chen J, Jiang W, Cen L, Pan J, Yu C, et al. The relationship between Helicobacter pylori infection of the gallbladder and chronic cholecystitis and cholelithiasis: a systematic review and meta-analysis. *Canadian Journal of Gastroenterology and Hepatology*. 2021; 2021.
8. Svistunov A, Osadchuk M, Mironova ED, Vasil'Eva IN. Helicobacter pylori as a risk factor for the development of metabolic syndrome and gallstone disease. *Медицинский вестник Северного Кавказа*. 2021;16(2):135-9.
9. Jahantab MB, Safaripour AA, Hassanzadeh S, Yavari Barhaghtalab MJ. Demographic, Chemical, and Helicobacter pylori Positivity Assessment in Different Types of Gallstones and the Bile in a Random Sample of Cholecystectomized Iranian Patients with Cholelithiasis. *Canadian Journal of Gastroenterology and Hepatology*. 2021;2021.
10. Cen L, Pan J, Zhou B, Yu C, Li Y, Chen W, et al. Helicobacter Pylori infection of the gallbladder and the risk of chronic cholecystitis and cholelithiasis: A systematic review and meta-analysis. *Helicobacter*. 2018;23(1):e12457.

11. Kunnuru SKR, Kanmaniyan B, Thiagarajan M, Singh BK, Navrathan N. A Study on Efficacy of UGI Scopy in Cholelithiasis Patients before Laparoscopic Cholecystectomy. Minimally Invasive Surgery. 2021;2021.
12. Hussain T, Kavya S. Assessment of the Relationship between the Presence of Helicobacter pylori Infection and Cholelithiasis. IJSS Journal of Surgery. 2021;7(5):63-6.
13. Eslami B, Iranshahi M, Gachkar L, Hadavand F. Gallstone Frequency in Patients with Helicobacter pylori Gastritis. Archives of Clinical Infectious Diseases. 2021;16(1).
14. Attaallah W, Yener N, Ugurlu MU, Manukyan M, Asmaz E, Aktan AO. Gallstones and Concomitant Gastric Helicobacter pylori Infection. Gastroenterol Res Pract. 2013;2013:643109. DOI: 10.1155/2013/643109.
15. Zhang FM, Yu CH, Chen HT, Shen Z, Hu FL, Yuan XP, et al. Helicobacter pylori infection is associated with gallstones: Epidemiological survey in China. World J Gastroenterol. 2015;21(29):8912–9. DOI: 10.3748/wjg.v21.i29.8912.
16. Sabbaghian MS, Ranaudo J, Zeng L, Alongi AP, Perez-Perez G, Shamamian P. Identification of Helicobacter spp. in bile and gallbladder tissue of patients with symptomatic gallbladder disease. HPB (Oxford). 2010;12(2):129–33. DOI: 10.1111/j.1477-2574.2009.00148.x.
17. Lee JW, Lee DH, Lee JI, Jeong S, Kwon KS, Kim HG, et al. Identification of Helicobacter pylori in Gallstone, Bile, and Other Hepatobiliary Tissues of Patients with Cholecystitis. Gut Liver. 2010;4(1):60–7. DOI: 10.5009/gnl.2010.4.1.60.
18. Monstein HJ, Jonsson Y, Zdolsek J, Svanvik J. Identification of Helicobacter pylori DNA in human cholesterol gallstones. Scand J Gastroenterol. 2002; 37(1):112–9. DOI: 10.1080/003655202753387455.
19. Takahashi Y, Yamamichi N, Shimamoto T, Mochizuki S, Fujishiro M, Takeuchi C, et al. Helicobacter pylori infection is positively associated with gallstones: a large-scale cross-sectional study in Japan. J Gastroenterol. 2014;49(5):882–9. DOI: 10.1007/s00535-013-0832-z.
20. Fikry AA. Helicobacter pylori infection in patients with chronic calculous cholecystitis a cross-sectional study. Journal of Surgery. 2014;2(4). DOI: 10.11648/j.js.20140204.12.

© 2022 Memon et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/86421>