



Distribution Study of the Different Forms of Soil Acidity and Available Nutrients in Upper Brahmaputra Valley Zone (UBVZ) of Assam

Manoharmayum Monica Devi ^{a*}, Devajit Bhattacharyya ^b,
Kulendra Nath Das ^b and Keisham Dony Devi ^c

^a Department of Soil Science and Agriculture Chemistry, Central Agricultural University, Imphal, Manipur, India.

^b Department of Soil Science, Assam Agricultural University, Jorhat, Assam, India.

^c Department of Agronomy, Central Agricultural University, Imphal, Manipur, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i183328

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102965>

Original Research Article

Received: 03/05/2023

Accepted: 12/07/2023

Published: 21/07/2023

ABSTRACT

Soil acidity is a major constraint that affect crop growth and production in larger aspect. The objective of the study is to characterize different forms of soil acidity components in relation to the soil properties. So, 14 georeferenced soil samples were collected from Jorhat and Sibsagar district of Assam. The soils of UBVZ of Assam were highly to moderately acidic (pH range 4.27 to 5.34) with sandy loam to silty clay loam in texture and high content of organic carbon ranged from 0.72 to 2.59%. The exchangeable Ca and Mg value ranged from 1.40-2.64 cmol(p+)/kg and 0.7-2.0 cmol(p+)/kg, respectively with CEC from 7.8 to 14.2 cmol(p+)/kg and BS from 25.03 to 43.76%. Correlation between soil physico-chemical properties and soil acidity components revealed that OC

*Corresponding author: E-mail: manoharmonica17@gmail.com;

had positive correlation with total acidity ($r = 0.638^*$) and exchange acidity ($r = 0.551^*$) while BS% had negative significant correlation with total acidity ($r = -0.540^*$). Exchangeable calcium was significantly correlated with total acidity ($r = -0.751^{**}$), exchange acidity ($r = -0.610^*$) and exchangeable H⁺ ($r = -0.557^*$) while Ex Mg had significant negative correlation with exchangeable H⁺ ($r = -0.596^*$). Deficiency of various nutrients in soil can hinder the yield productive prospects in future. So, we must think of the holistic approach to meet the present need of the nutrients.

Keywords: Soil; nutrients; acidity; correlation and properties.

1. INTRODUCTION

Soil acidity is an important constraint in agriculture production. It is a natural process occurred due to heavy rainfall and leaching of cations like calcium, potassium and magnesium. Acidic parent materials and decomposition of organic matter also leads to formation of acidic condition in soil. Acceleration of acidification is caused by acid forming fertilizers usage, disturbance of soil structure and cultivation of high yielding crops [1]. Due to these factors' acid soils are generally poor in nutrient content. In strongly to moderately acidic soils, plant growth and crop yield are hampered to a large extent [2]. Higher soil acidity leads to decline in productive potential of the soils.

Soil acidity is a major agricultural problem affecting productivity worldwide [3]. The strengths of soil acidity and its influences on fertility and productivity are further embellished by the present need for over exploitation of land for food security [4]. In India, soil acidity affects approximately 1/3rd of arable land [5] and out of 245 million hectare of cultivated land, acidic soils occupy nearly 49 Mha area, which occupy nearly 34% of the total cropped area of the country [5,4]. It was recorded that 26 Mha had pH below 5.5 and 23 Mha had pH between 5.6 and 6.5 out of 49 million hectares of acidic soil in India [6].

North- eastern region of India covered about 21 million hectares of acid soils. The region occupies with nearly 65 per cent of the lands which are in extreme forms of soil acidity with pH of < 5.5 [7,4] constituting a major area under acid soils. Approximately 15 Mha of land area of the north-eastern region is acidic with pH varying from strongly acidic to slightly acidic (pH 4.0-6.3) and with pH below 5.5 occupying 54% of the acid soils in the country [4]. In the context of Assam,

more than 60 per cent of the land was affected by soil acidity problem, thereby affecting crops in many ways and its effects were mostly indirect through its influence on chemical factors such as aluminium and manganese toxicity calcium, phosphorus and magnesium deficiency and microbiological process.

Soil acidity marks a tremendous impact on soil chemical and biological properties besides regulating nutrient accessibility and aluminium toxicity [8]. So, amelioration of soil acidity through liming is the best strategy for improving efficiency and productivity of acid pretentious soil. Soil acidity is usually defined in terms of different forms of soil acidity components viz., total acidity, exchange acidity exchangeable aluminium, etc. The different form of these acidity components and their proportion in soil regulates fixation of nutrients and extent of lime requirement. Precise information of the soil physico-chemical properties and indication of the different forms of acidity components assist in understanding as well as management of acid soils. In light to this information, attempt has been made to study the different soil physico-chemical properties, characterized acidity components and correlates their relationship among them.

2. MATERIALS AND METHODS

2.1 Soil Sample Collection

The studied soil samples were collected from Jorhat and Sibsagar districts of UBZ of Assam. A total of 14 Geo referenced surface soil (0-15 cm) samples were collected. Analysis of the different soil properties for the illustration of the key features of soil were done by the standard procedures.




Fig. 1. Location map of Jorhat and Sibsagar district of Assam

2.2 Processing of Soil Samples

The collected soil samples were grinded using wooden mortar and pestle and mixed uniformly. It was then aired dried in shade and sieved with 2 mm sieve and thereafter packed in zipped polythene bags with proper labelling.

2.3 Analysis of Soil Samples

The particle size distribution of the soil samples was determined by International Pipette Method [9]. Soil moisture at field capacity (0.3 bars) was determined using Pressure plate and pressure membrane apparatus. The collected soil samples were then analysed for pH, EC, CEC and K₂O using standard procedures as described by Jackson [9]. The collected soil samples were analysed for available nitrogen [10], available P₂O₅ [11], neutral ammonium acetate extractable K₂O [9], organic carbon [12], and Lime requirement (LR) by Shoemaker, McLean and Pratt (SMP) Method [13]. The exchangeable calcium and magnesium were estimated by Versene Titration Method [14]. Percentage Base Saturation (PBS) was determined by using the following formula:

$$\text{Percent Base Saturation (PBS)} = (\text{Ca} + \text{Mg} + \text{Na} + \text{K}) / \text{CEC} * 100.$$

Acidity components of the soils were determined as per the procedure outlined by Baruah and Barthakur [154]. Soil acidity components were characterized as follows: total acidity was determined by 1N NaOAc solution. The exchange acidity and exchangeable Al were extracted by 1N KCl solution and determined by titration of 0.1N NaOH using phenolphthalein as indicator and by back titration after acidification with 40g/L NaF with 0.1N HCl solution respectively. Exchangeable H⁺ was determined by subtracting exchangeable Al from exchangeable acidity i.e.

$$\text{Exchangeable H}^+ = \text{Exchangeable acidity} - \text{Exchangeable Al}$$

2.4 Statistical Interpretation

Simple correlation study between different forms of acidity and physico-chemical properties of soil and with soil available nutrients were computed using simple correlation coefficient.

3. RESULTS AND DISCUSSION

3.1 Physico-chemical Properties of the Soils

The data of physico-chemical properties of the soils collected from the Upper Brahmaputra Valley Zone (UBVZ) of Assam is presented in the Table 1. Texture of the soils of UBVZ region of Assam were loamy sand, sandy loam, loam and silty clay loam (Table 1). Sand, silt and clay fraction of soil varies from 14.22 to 75.85, 10.00 to 61.90 and 4.90 to 29.70 per cent, respectively with mean values of sand, silt and clay were 53.75, 25.34 and 16.24 per cent, respectively. The removal of finer particles from the surface soil due to heavy rainfall in rainy season might have contributed to loamy sand and sandy loam texture of the surface soils [16]. Moisture content of the soil at field capacity (0.3 bars) at varied from 16.08 to 32.79 per cent with the mean value of 21.79 per cent (Table 1). The variation in sand, silt and clay fractions of the soils might have resulted in variations in moisture content at field capacity.

The pH of the soil varied from highly acidic to strongly acidic in reaction ranging from 4.27 to 5.34 with a mean value of 4.96. This might be attributed to higher degree of pedogenic development resulting in leaching in basic cations and soluble salts and dominance of exchangeable H⁺ and Al³⁺ as reported by Sharma and Sarangthem [17]. The electrical conductivity (EC) of the soils varied from 0.49 to 2.01 dS/m. The organic carbon content of the soils ranged from 0.72 to 2.59 per cent with mean value of 1.61 per cent of organic carbon which might be due to more vegetation and climatic conditions of the study area (Table 1).

Data of exchangeable bases in the soils of UBVZ of Assam are presented in Table 1. The calcium was the dominant exchangeable basic cation of the soils and it ranged from 1.40 to 2.64 cmol(p+)/kg with the mean value of 2.11 cmol(p+)/kg. The range of exchangeable magnesium, exchangeable sodium and exchangeable potassium content of the soils were from 0.70 to 2.00 cmol(p+)/kg, 0.11 to 0.79 cmol(p+)/kg and 0.05 to 0.11 cmol(p+)/kg, respectively. The mean value of these basic cations viz., Ca, Mg, Na and K were 2.11, 1.39, 0.25 and 0.09 cmol(p+)/kg, respectively. Baruah et al. [18] also made similar observation in the soils of north-eastern region. Base saturation of the soils of UBVZ of Assam varied from 25.03 to

43.76 per cent which were considered low BS% status. Das and Sudhakar [19] reported that low B.S % (< 60 %) of the soil of North-eastern region and attributed it due to leaching of the bases under high rainfall conditions. The CEC of the soil of the UBVZ of Assam ranged from 7.80 to 14.20 cmol(p+)/kg (Table 1). The variation in CEC of different soils might be attributed to variation in clay of organic carbon and free iron and Al oxides content of the soils [5].

The range and mean of Lime Requirement (LR) of the soils were 0.56 to 11.42 t/ha and 7.52 t/ha, respectively (Table 1). Magnitude of lime requirements in soils of UBVZ of Assam ranged from 0.56 to 11.42 t/ha with mean value of 7.42 t/ha (Table 1). Report available in acid soils of Assam revealed that LR of the soils varied with clay content, CEC and pH of the soil [20]. Reports also revealed that in acid soils of Assam LR of the soils not only governed pH alone but by the combined influence of soil pH, exchange acidity and exchangeable Al [21]. Significant positive correlation of LR of the soils of UBVZ of Assam was also observed with clay content ($r= 0.535^*$) in the present investigation (Table 3).

3.2 Available Macronutrients

Available nitrogen (N) status in the soil of UBVZ of Assam ranged from low to medium with its content ranging from 100.80 to 260.22 kg/ha as shown in Table 2. The N-content of the soils depend on factors like soil type, soil pH, climate vegetation, fertilizer management etc. Among these, climate seemed to have most striking effect on organic carbon content in soils of Brahmaputra valley [22] which in turn determine N- content of these soils.

Available phosphorus (P₂O₅) content of the soils of UBVZ of Assam varied from low 12.73 kg/ha to medium 29.80 kg/ha (Table 2). Hence available P₂O₅ status of these soils can be considered as low. In acid soils of Assam fixation of both native and applied P is generally considered as its low availability in these soils [18]. High positive correlation existed between available P (Bray'I) and Al-P and Fe-P indicated that these factors mostly contribute to availability of P in acid soils of Assam [23].

The exchangeable potassium (K₂O) content of the soil of UBVZ of Assam varied from 40.86 kg/ha to 90.01 kg/ha which was low in status (Table 2). The type and nature of soils were most

important factors affecting soil K₂O status through fixation and release of K in the soil [18]. Fixation of K was also found to correlate positively with clay fractions of soil but the correlation was not significant with silt and sand fractions suggesting K fixation in soils was related to the clay fractions of the soils (Table 5).

The secondary nutrients exchangeable calcium (Ca) and exchangeable magnesium (Mg) varied from 1.40 to 2.64 and 0.70 to 2.00 cmol(p+)/kg respectively (Table 1). The calcium was the dominant exchangeable basic cation followed by exchangeable Mg of the soils of the soils of UBVZ of Assam. Similar observations were also reported by Gangopadhyay et al. [24] in tea growing soils of Assam. In acid soils of India Ca saturation of exchange complex generally varied from 20 to 40 per cent. Total CaO and MgO of this acidic soils were low (0.1 to 1.0 % and 0.23 to 1.2 %, respectively). Therefore, management of acid soils involved use of liming material [25].

The correlation found in soils of UBVZ of Assam between forms of soil acidity and exchangeable Ca and Mg content of the soils was mostly negative which also indicates that with increase in soil acidity in any form resulted in decrease of Ca and Mg content of these soils (Table 5). However, correlations between total acidity, exchange acidity and exchangeable H⁺ and exchangeable Ca were statistically significant with r values -0.751^{**} , -0.610^* and -0.557^* , respectively. The exchangeable Mg content of the soils had significant negative correlations with exchangeable H⁺ ($r = -0.596^*$) only (Table 5).

3.3 Forms of Soil Acidity

3.3.1 Total acidity

The findings of acidity components of the soils of UBVZ of Assam are presented in Table 2. The results on forms of acidity revealed that total acidity varied from 1.38 to 3.75 cmol(p+)/kg with the mean value of 2.41 cmol(p+)/kg in soils of UBVZ of Assam. It was the most dominant forms of acidity in soils of UBVZ of Assam (Table 2). The correlation study of the different forms of soil acidity with soil physico-chemical properties are shown in Table 4. In soils of UBVZ of Assam significant positive correlation of total acidity ($r=0.638^*$) was found with OC and its correlation with BS% was significant and negative ($r= -0.540^*$) (Table 4).

Table 1. Physico-chemical characteristics of the soils of UBVZ of Assam

Soil No.	pH	EC	OC (%)	% sand	% silt	% clay	CEC (cmol(p+)/kg)	B.S %	M.C (%)	LR (t/ha)	Ca (cmol(p+)/kg)	Mg (cmol(p+)/kg)
1	5.08	0.72	1.23	75.85	19.30	4.90	9.60	43.76	19.82	0.56	2.20	1.80
2	4.98	0.80	2.16	69.02	22.30	15.30	13.11	35.19	19.53	6.56	2.07	2.00
3	4.79	0.90	0.78	56.16	31.60	17.50	13.80	30.53	18.83	11.42	2.13	1.90
4	4.27	0.67	0.93	65.00	15.00	20.00	9.64	40.34	20.80	7.78	2.16	0.86
5	5.16	0.97	1.41	70.00	10.00	20.00	11.09	33.47	18.56	2.92	1.88	1.62
6	4.75	0.96	1.80	39.33	34.90	17.20	12.50	33.50	26.69	8.99	2.20	1.80
7	4.94	0.49	2.59	14.22	61.90	26.70	11.23	25.03	21.15	11.42	1.40	1.10
8	5.25	0.90	0.72	65.80	18.10	8.38	9.80	43.63	16.81	7.78	2.60	1.40
9	4.63	2.01	2.34	56.29	17.33	5.70	8.90	40.61	32.79	10.21	2.20	1.20
10	4.82	1.27	2.19	30.09	27.00	22.30	7.80	41.10	18.35	8.99	2.30	0.70
11	5.04	1.08	1.98	87.41	8.70	9.30	11.30	29.62	24.88	4.13	1.70	1.09
12	5.26	0.84	1.54	23.85	39.00	29.70	14.20	30.92	26.11	11.42	2.64	1.40
13	5.34	1.11	1.53	29.49	34.37	17.13	12.50	29.51	24.67	7.78	2.10	1.20
14	5.13	0.84	1.32	69.95	15.30	13.20	8.50	43.65	16.08	5.35	2.00	1.40
Mean	4.96	0.97	1.61	53.75	25.34	16.24	10.99	35.77	21.79	7.52	2.11	1.39
Range	4.27-5.34	0.49-2.01	0.72-2.59	14.22-87.41	10.00-61.90	4.90-29.70	7.80-14.20	25.03-43.76	16.08-32.79	0.56-11.42	1.40-2.64	0.70-2.00
SD	0.28	0.34	0.57	21.51	13.68	7.16	1.95	6.06	4.48	3.21	0.31	0.38

Table 2. Different forms of soil acidity components (Total Acidity, Exchange Acidity, Exchangeable Aluminium & Exchangeable Hydrogen) and available nutrients (Nitrogen, phosphorus and potassium) in soil of UBVZ of Assam

Sample no.	Total Acidity	Exchange Acidity	Exchangeable Aluminium	Exchangeable Hydrogen	Nitrogen	Phosphorus	Potassium
1	2.13	1.25	0.88	0.38	102.23	22.30	61.02
2	2.75	2.25	1.50	0.75	170.30	20.80	69.87
3	2.75	1.88	1.38	0.50	162.21	23.40	40.86
4	2.25	2.13	0.63	1.50	100.80	29.80	70.03
5	1.88	1.75	1.13	0.63	150.24	24.30	89.84
6	2.25	1.13	0.75	0.38	152.51	14.83	76.82
7	3.75	2.25	1.00	1.25	260.22	17.86	90.01
8	1.38	0.88	0.78	0.10	220.19	27.42	83.17
9	3.00	2.63	1.25	1.38	210.50	23.40	78.76
10	2.13	2.00	1.25	0.75	223.90	20.55	76.07
11	3.63	2.63	1.15	1.48	240.10	12.73	78.71
12	1.88	1.38	0.75	0.63	211.20	16.35	83.06
13	2.00	1.88	1.13	0.75	225.21	24.55	89.24
14	2.00	2.13	1.13	1.00	219.50	14.42	83.60
Mean	2.41	1.87	1.05	0.82	189.22	20.91	76.50
Range	1.38-3.75	0.88-2.63	0.63-1.50	0.10-1.50	100.80-260.22	12.73-29.80	40.86-90.01
SD	0.66	0.52	0.25	0.43	48.02	4.90	12.72

Table 3. Correlation coefficient among Lime Requirement (LR) and physico-chemical properties of the soils

Parameters	Physico-chemical properties of the soils					
	pH (H2O)	OC	Clay	EC	CEC	BS
Lime Requirement	-0.271	0.211	0.535*	0.150	0.315	-0.378

**Significant at 0.05% level of significance*

Table 4. Correlation coefficient among soil acidity components and soil physical and chemical properties

Parameters	Physico-chemical properties					
	pH (H ₂ O)	O.C	CEC	Clay	BS%	LR
Total Acidity	-0.319	0.638*	0.127	0.0004	-0.540*	0.199
Exchange Acidity	-0.355	0.551*	-0.176	-0.030	-0.224	0.081
Exchangeable Aluminium	0.090	0.374	0.071	-0.150	0.006	-0.160
Exchangeable Hydrogen	-0.488	0.451	-0.258	-0.052	-0.161	0.096

**Significant at 0.05% per cent level.*

*N=14, df=12, *P_{0.05}=0.532*

Table 5. Correlation coefficient among soil acidity components and macronutrients

Parameters	Available Macronutrients				
	N	P	K	Ca	Mg
Total Acidity	0.314	-0.386	-0.106	-0.751**	-0.129
Exchange Acidity	0.310	-0.165	-0.042	-0.610*	-0.388
Exchangeable Aluminium	0.273	-0.119	-0.237	-0.319	0.210
Exchangeable Hydrogen	0.217	-0.131	-0.191	-0.557*	-0.596*

**Significant at 0.05% per cent level*

3.3.2 Exchange acidity

The exchange acidity of the soils had a varied from 0.88 to 2.63 cmol(p+)/kg and its mean value was 1.87 cmol(p+)/kg as shown in Table 2. The OC content of the soils is correlated positively and significantly with exchange acidity ($r=0.551^*$). Lime requirement of the soil showed positive correlation with exchange acidity (0.081) even though not significant. However, there is negative correlation between exchange acidity with pH, CEC, clay content and base saturation per cent of the soils (Table 4).

3.4 Exchangeable Aluminium & Exchangeable Hydrogen

The exchangeable aluminium and the exchangeable hydrogen of the soils varied from 0.63 to 1.50 cmol(p+)/kg and 0.10 to 1.48 cmol(p+)/kg, respectively and the mean exchangeable Al and exchangeable H of the soils of UBVZ of Assam were 1.05 and 0.82 cmol(p+)/kg, respectively (Table 2). Similar observations on forms of acidity in acid soils of Assam were also made by Dhananjaya and Ananthanarayana [26]; Sahoo et al. [27]; Reza et al. [28] and Gangopadhyay et al. [24]. Badole et al. [29] and Sangtam et al. [30] reported significant negative correlations of all forms of acidity with pH (H₂O) in acid soils. However, in the soils of UBVZ of Assam though forms of acidity correlate negatively with pH it was not significant (Table 4) [31].

4. CONCLUSION

The analysis of the soil samples collected from Upper Brahmaputra valley zone of Assam showed a variable trend. An amalgamation of different soil properties renders the soil its optimum soil health status and maintain balance crop nutrition in environment. The different soil acidity components viz. total acidity, exchange acidity, exchangeable aluminium and exchangeable hydrogen showed fickle contribution in acidic soils. The total and exchange acidity showed a significant positive correlation with OC while exhibited significant negative correlation with BS% and exchangeable calcium. Among the different soil acidity components, exchangeable hydrogen showed significant negative correlation with exchangeable magnesium. Deficiency of various nutrients in soil can hinder the yield productive prospects in future. The overall study helps in providing a comprehensive idea about soil types,

nutrient deficiency and plant condition. It also assists the farmers to optimize the quantity of different fertilizers for profitable production.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Fageria NK, Baligar VC. Ameliorating soil acidity of tropical Oxisols by liming for sustainable crop production. *Adv. Agron.* 2008;99:345-431.
2. Bhatt JA, Kundu MC, Hazra GC, Santra GH, Mandal B. Rehabilitating acid soils for increasing crop productivity through low-cost liming materials. *Sci. Tol. Environ.* 2010;408:4346-4353.
3. Borgohain L, Tamuly D, Borah N, Dutta S, Nath DJ, Thakuria RK, et al. Kinetics of Nitrogen Mineralization as Influenced by Liming and Organic Amendments: A Laboratory Study. *Intern. J. Curr. Microbiol. Appl. Sci.* 2019;8(1):853-865.
4. Devi MM, Bhattacharyya D, Das KN, Patgiri DK, Kurmi K, Saikia H, Devi KD. Dynamics of acidity components of the soils as influenced by liming in Upper Brahmaputra Valley Zone of Assam. *Eco. Environ. & Cons.* 2023;1:S131-S136.
5. Maji AK, Reddy GP, Sarkar D. Acid soils of India-Their extent and Spatial Variability. *NBSS and LUP, Nagpur* 2012;1:138-145.
6. Thakuria D, Hazarika S, Rangappa K. Soil Acidity and Management Options. *Ind. J. Fert.*, 2016; 12: 40-56.
7. Sharma UC, Singh RP. Acid soils of India: their distribution, management and future strategies for higher productivity. *Fert. News* 2002;47:45-52.
8. Rahaman MA, Lee SH, Ji HC, Kabir AH, Jones CS, Lee AW. Importance of Mineral Nutrition for Mitigation Aluminum Toxicity in Plants on Acidic Soils: Current Status and Opportunities. *Int. J. Mol. Sci.* 2018;19(10):3073.
9. Jackson ML. *Soil Chemical Analysis.* Prentice Hall of India, New Delhi; 1973.
10. Subbiah BV, Asija GL. A rapid procedure for the determination of available nitrogen in soils. *Curr. Sci.* 1956;25:259-260.
11. Bray RH, Kurtz LT. Determination of total, organic and available forms of phosphorus in soils. *Soil Sci.* 1945;59(1): 39-46.

12. Walkley A, Black IA. An examination of the digestion method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.* 1934;37:29-33.
13. Shoemaker HE, McLean EO, Pratt PF. Buffer methods for determining lime requirement of soils with appreciable amounts of extractable aluminium. *Proc. Soil Sci. Soc. Am.* 1961;25:274-277.
14. Barrow HL, Simpson EC. An EDTA Method for the Direct Routine Determination of Calcium and Magnesium in Soils and Plant Tissue. *Soil Sci. Soc. America J.* 1962;26(5):443-445.
15. Baruah TC, Borthakur HP. *Textbook of Soil Analysis.* Vikas Publishing House Pvt. Ltd., New Delhi; 1997.
16. Bandhyopadhyay KK, Hati K, Singh R. Management options for Improving Soils Physical Environment for Sustainable Agricultural Production: A Brief Review. *J. Agril. Physics.* 2009;1:1-8.
17. Sharma LD, Sarangthem I. Nature of Acidity and Lime Requirement in Acid Soils of Manipur. *Intern. J. Adv. Sci. Res. Engg. Trends.* 2018;2(1):51-57.
18. Baruah, Baruah K, Haque A, Das B, Medhi C, Misra AK. Boron in Soil and Water Samples in some Tea Garden Belts of Golaghat district, Assam. *Adv. Appl. Sci. Res.* 2011;2(4):298-305.
19. Das P, Sudhakar S. Land Suitability Analysis for Orange and Pineapple. A Multi Criteria Decision Making Approach Using Geo Spatial Technology. *J. Geographic Informn. Syst.* 2014;6(1):40-44.
20. Borah DK, Nath AK. Effect of pH and moisture regime on the transformation of native and applied phosphate in alluvial soils. *J. Res. Assam Agric. Univ.* 1980; 1(2):148-152.
21. Halder BR, Mandal LN. Lime requirement of acid soils in relation to pH, exchangeable acidity, extractable acidity and exchangeable aluminium content of the soils. *J. Indian Soc. Soil Sci.* 1985;33(3):528-535.
22. Heikkinen J, Keskinen R, Kostensal J, Nuutinen V. Climate change induces carbon loss of arable minerals soils in boreal conditions. *Gob. Chang. Bio.* 2022;28(12):3960-3973.
23. Dey JK, Bhattacharyya D. Soil factors affecting phosphate fixation in inceptisols of Assam. *Indian J. Agric. Sci.* 1996; 66(4):253.
24. Gangopadhyay SK, Nayak DC, Singh SK. Characteristics of tea growing soils in relation to soil acidity in Upper Brahmaputra Valley of Assam. *J. Indian Soc. Soil Sci.* 2016;64 (4):341-350.
25. Pasricha NS, Sarkar AK. Secondary nutrients. In: *Fundamentals of Soil Science.* Goswami NN, Rattan KK, Dev G, Narayanasawy G, Das DK, Sajal SK, Pal DK, Rao DLN. (eds.). *Indian Soc. Soil Sci., New Delhi,* 2009;1:457.
26. Dhananjaya BC, Ananthanarayana DR. Forms of acidity in some acidic Inceptisols of southern Karnataka. *Agropedology.* 2010;20 (1):49-52.
27. Sahoo AK, Sarkar D, Baruah U, Butte PS. Characterization, classification and evaluation of soils of Longhol hill, Manipur for rational land use planning. *J. Indian Soc. Soil Sci.* 2010;58(1):1-6.
28. Reza SK, Baruah U, Bandopadhyya S, Sarkar D, Dutta DP. Characterization of soil acidity under different land uses in Assam. *Agropedology.* 2012;22(2):123-127.
29. Badole S, Datta A, Basak N, Seth A, Padhan D, Mandal B. Liming influences forms of acidity in soils belonging to different orders under subtropical India. *Commun. Soil Sci. Plant Anal.* 2015; 46(16): 2079-2094.
30. Sangtam S, Gohain T, Kikon N. Assessment of nitrogen doses and planting densities for optimizing growth and yield performance of rainfed maize (*Zea mays* L.). *Indian J. Agric. Res.* 2017;51(5): 473-477.
31. Barua NG, Borah DK, Medhi BK, Bhattacharyya D, Borah N. *Soil Chemistry.* In: *Seven Decades of research on Soils of Assam.* Borah, D. K. and Patgiri, D. K. (eds.). *Niche Area of Excellence, Department of Soil Science, AAU, Jorhat.* 2011:75-126.