



Study of Genetic Drifts in Some Fat Mass and Obesity Associated Gene Variatians in Type 2 Diabetics in Some Ethnic Groups Residing in Delta State, Nigeria

H. Osere ^{a*}, H. Brown ^a and K. N. Elechi-Amadi ^a

^a *Department of Medical Laboratory Science, Rivers State University, Port Harcourt, Rivers State, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRIMPS/2024/v13i1244

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111330>

Original Research Article

**Received: 06/11/2023
Accepted: 10/01/2024
Published: 16/01/2024**

ABSTRACT

Aim: Detailing and distribution of some predominant FTO genes among subjects with diabetes mellitus selected from Igbo, Ika, Ijaw and Isoko tribes of Delta state, Nigeria.

Study Design: Case-control, observational study.

Place and Duration of Study: Federal Medical Center, Asaba, Delta State and Safety Molecular Pathology Laboratory Enugu, Nigeria, between March 2020 and Febuary 2022.

*Corresponding author: E-mail: crystalann4jah@yahoo.com;

Methodology: 100 subjects (20 diabetic and 5 non diabetic from each of the four tribes; age range; 28-74 years) Venous blood was collected for the assay of FBS, Hb1ac and molecular assay of four variants of FTO (Fat Mass and Obesity Associated) gene. Results obtained were statistically analysed.

Result: The genotypes identified in the subjects were AA, GA, GG, AC, CC, TT, and AT. AA (Ijaw 17, Urhobo 18, Ika 18, Igbo 19) and GG (Ijaw 17, Urhobo 18, Ika 19, Igbo 18) had highest level of occurrence in all tribes. Recombinant genotype and recessive allele had higher number of subjects as compared to the originating genotypes. Hardy-Weinberg statistics demonstrated significant genetic deviations in rs2388405 (AA -0.12 $x^2=0.0002$, GA 0.12 $x^2=0.002$) rs201041270 (GG -0.12 $x^2=0.0002$, GA 2.12 $x^2=0.002$), and rs8050136 (CC -0.34 $x^2=0.001$, AC 2.32 , $x^2=0.05$) proving a chance of finding new SNP in the studied population with a probability of 2.25 as seen in the odd ratio of T allele in Igbo tribe.

Conclusion: Statistically significant genetic drifts exist in the studied population.

Keywords: Obesity; diabetes; ethnic groups; gene; genetic drifts; fat mass.

1. INTRODUCTION

Diabetes mellitus, defined as a metabolic condition that is characterized by persistent hyperglycemia caused by insulin insufficiency or insulin resistance [1]. Diabetes mellitus (DM) is a major cause of mortality and disability globally, this raises concerns about the financial burden it places on both people and society as a whole [2]. Endocrinology and Metabolism Society of Nigeria (EMSON) reported in 2022 that over 10 million Nigerians are living with diabetes and warned that if preventive measures are not applied, the numbers are expected to double by 2030 [3]. An obesity sensitivity gene, whose intron 1 region has multiple SNPs that are strongly associated with Body Mass Index, energy metabolism and anthropometric indices has been identified, the gene is known as the fat mass and obesity-associated (FTO) gene [4]. The FTO gene exists in different genetic forms known as variants of FTO polymorphisms. FTO gene rs9939609 SNP is significantly associated with obesity, With the AA genotype carriers having a 2.02 increased risk of developing obesity [5]. Interdependence exist between biochemical parameters of T2D with FTO gene, namely insulin, glucose, HOMA, BMI, waist and hips circumference [6].

Results obtained from genetic risk score (GRS) calculated for two FTO single-nucleotide polymorphisms (SNPs) rs8050136 and rs2388405 demonstrated that the two SNPs were significantly related to increased BMI and obesity [7]. Understanding the genetic make-up of FTO gene, its various variants, distribution and how it affects and influences manifestation of diabetes can assist with early intervention measures, improvement of diabetes clinical diagnostic processes, and grant insight on developing targeted treatment options for patients with

diabetes. Reports of results obtained from various studies for FTO gene variants association with obesity and diabetes varies with population and continents. This research will give insight of FTO gene presence and distribution in various ethnic tribes residing in Delta state region of Nigeria.

2. MATERIALS AND METHODS

2.1 Study Area

This research work was done in Delta State, Nigeria (Fig. 1). Samples were collected from subjects attending clinic at Federal Medical Centre, Asaba and General Hospitals in Agbor & Bomadi Delta State, Nigeria. Participants were people from Ijaw, Urhobo, Ika and Igbo ethnic groups residing in Delta State. Nigeria's Delta State capital, Asaba, is located on a terrace of the lower Niger River, with geographical coordinates of $6^{\circ}11'52.23''N$, $6^{\circ}43'42.48''E$. Asaba serves as a link between western, eastern, and northern Nigeria through the Asaba Niger Bridge, which connects East and West, and the Niger River in the north.

2.2 Research Design

This is a case-control, observational study involving the identification of FTO gene allele variants, distribution of the gene evaluation of HbA1c, Fasting blood sugar (FBS), in Type 2 Diabetic (T2D) and non-diabetic control subjects from the selected ethnic groups in Delta State. The bio-data and medical history of the subjects were obtained through questionnaire.

2.3 Sample Size

Sample size for this study was determined using the Cochran formula:

$$N = \frac{Z^2 pq}{d^2} \quad [9]$$

N = the desired sample size

Z = The Standard Normal deviate usually set at 1.962 corresponding to the 95% Confidence level

p = The SNPs Prevalence rates. (Minor Alleles Frequency of SNPs set at >0.02) [10]

q = 1- p

d = degree of accuracy desired set at 0.05

Minimum Size – 30

By adding 10% of non-respondent = 33.

Therefore, total sample size is 33.

However, this study used 100 subjects, with 20 subjects selected from each of the four ethnic groups in Delta State, and 20 control subjects.

2.4 Sampling Method

A multistage sampling technique was used to choose the subjects. Participants were grouped into sections based on duration of pathogenesis (1-5years, 6-10years, 11-15years, 16-20years).

2.5 Selection Criteria

2.5.1 Inclusion criteria

Individuals who are of the selected tribes in Delta State aged at least 21 years and above,

diagnosed with T2D for at least one year. Controls: Five (5) individuals from each of the selected tribes Ika, Urhobo, Ijaw, Igbo with no history of diabetes, and are non-obese and having a fasting blood glucose of less than 6.5mmol/l.

2.5.2 Exclusion criteria

1. Individuals not from the selected tribes.
2. Critically ill subjects.
3. Pregnant female.

2.6 Sample Collection

From each subject, venous blood was collected using standard veni-puncture technique, 2ml was dispensed into fluoride-oxalate bottle for the assay of FBS and Hb1ac another 3ml was dispensed into EDTA tube for molecular assay of FTO genes.

After allowing the sample in the plain tube to retract, it was centrifuged. The analysis was performed within one week of collection at the Federal Medical Centre, Asaba. The first EDTA tube was transported in cold box to Safety Molecular Pathology Laboratory Services located at 44 Rangers Avenue, Enugu for DNA extraction and genotyping (Sequencing). The second EDTA tube for Glycated hemoglobin was for stored at 2-80C and analysis was done within two days of sample collection.



Fig. 1. Map of Delta state Nigeria, showing Ethnic and local government area [8]

2.6.1 Fasting Blood Sugar (FBS)

FBS was performed using glucose oxidase method.

Principle: After enzymatic oxidation in the presence of glucose oxidase, glucose is measured. Under the catalysis of peroxidases, the generated hydrogen peroxide combines with phenol and 4-aminophenazone to create a red-violet quinoneimine dye as an indicator.

2.6.2 Glycosylated Haemoglobin (HbA1c)

Quantitative determination of glycosylated Haemoglobin in blood was done using the modified Ion Exchange Resin method with kit from INTECO Diagnostics, UK [11].

2.7 Genetic Analysis

2.7.1 Genomic DNA extraction

Genomic DNA extractions of the samples was performed using Geneaid DNA Mini Kit (Blood/Cultured Cell).

2.8 Genotyping of SNPs

Genotyping of SNPs of the FTO gene was performed with the Illumina next-generation sequencing (NGS) using NextSeq 2000 Sequencing System. Concentration and purity of isolated DNA was determined by UV/VIS spectrophotometer NanoDrop ND-1000 (Appendix I).

2.9 Statistical Analysis

Data obtained was categorised based on duration of Type-2 diabetes, ethnicity, age and sex. Utilizing GraphPad Prism, version 8.0.2 (California, USA), Hardy Weinberg equilibrium, Student's statistical t-test at $P = .05$, Mean, Standard deviation, Gene counting, Chi-square and Odds ratio (OR) were used to determine genotypic and allelic frequencies, Variance, SNPs, significant differences in the study population of Type 2 diabetic (T2D) and non-diabetic participants.

3. RESULTS AND DISCUSSION

3.1 Metabolic and Immunologic Parameters

In (Table. 1) below presents the Metabolic and Immunologic parameters of subjects with FTO gene variations with varying duration of type 2 diabetes against the control subject. Fasting blood sugar (FBS) values was significantly higher

in subjects at various duration of diabetes pathogenesis than control subjects, with a gradual progressive rise over years of disease duration. Thus, confirming disease status of patients. hyperglycaemia is an indicator of diabetes mellitus [1]. the impact of continuous increase in FBS can be seen in progressive glycation of glucose to haemoglobin, as is reflected by values obtained for HbA1c which were significant as compared to control. This changes could be due to the dysregulation of glucose [12,13]. Increase in HbA1c as seen in (Table 1) suggest serves as an indicator of blood glucose control, and a marker of diabetes [14].

3.2 Distribution of FTO Genotypes

Genotypes identified in the studied population are AA, GA, GG, AC, CC, TT, AT. Distribution of these FTO genotypes according to tribes in (Table 2) below showed that, genotype AA and GG had highest level of occurrence in all tribes while genotype GA had the lowest level of occurrence in all tribes. FTO gene prevalence varies by ethnicity and race [15], AA genotype was significantly higher FTO in studied population [16].

3.3 Hardy-Weinberg Statistical Test of FTO gene Variants

In (Tables 3 and 4) Hardy-Weinberg statistics demonstrated significant negative and positive genetic deviations in rs2388405 (AA -0.12 $x^2 = 0.0002$, GA 0.12 $x^2 = 0.002$) rs201041270 (GG -0.12 $x^2 = 0.0002$, GA 2.12 $x^2 = 0.002$), and rs8050136 (CC -0.34 $x^2 = 0.001$, AC 2.32 , $x^2 = 0.05$) proving that there is a chance of finding new SNP in the studied population, as well as genetic deviations within the population which could be any or a combination of some of the following distribution factors, mutation, natural selection, non-random mating, genetic drift, or gene flow. similar finding exists among Brazilian population [17]. changes from expected frequency could be due to cross-breeding that may have occurred in previous generations [18], considering the multi-ethnic nature of the geographical population of the study area this might apply to the genetic drift observed for the studied FTO variants.

3.4 Genotype and Allele Distribution of rs2388405 Variant of FTO Gene in Obese and T2D Subjects and Control

(Table 5-8) demonstrates types and numbers of genotypes obtained by gene sequencing and

counting, its recombinant genotype and alleles. In this study, rs2388405 Genotype AA had the highest number of occurrences for T2D in all tribes. Recombinant genotype of rs2388405 AA+GA and its recessive allele A, had higher number of occurrences as compared to the originating genotype. Similarly, recombinant genotype GG+GA with the recessive allele G of rs201041270 had higher number of subjects as compared to the originating genotypes, having the highest number of subjects. In rs9939609 Genotype TT had the highest number of occurrences for T2D in all tribes with varying

number of occurrence, recombinant genotype TT+AT had the highest number of possible occurrence in all tribes. Allele T in rs9939609 had the highest number of subjects in Igbo tribe with an odd ratio value of 2.25, the greatest odds ratio for a T2D risk locus found to be 1.57 suggests that a greater number of as-yet-untested variants may be responsible [19]. Genotype CC of rs8050136 was significantly high in Ijaw tribe with an odd ratio of 0.93 as well as it allele C with odd ratio of 0.94 and x2 of 0.002 in both the genotype and allele. high number of allele occurrence and recombinant genotype

Table 1. Mean±SD of metabolic and immunologic parameters of subjects with FTO gene variations with varying duration of T2D against control subjects

Parameters	Control	1-5 yrs	6 -10 yrs	11 -15 yrs	16 -20	Fvalue	Pvalue	Remark
FBS (mmol/L)	4.92±0.65a	8.11±3.41b	8.27±2.88b	8.32±3.56b	7.35±3.37a	5.168	0.0008	S
HbA1c (%)	6.57±0.99a	8.84±1.86b	8.85±1.39b	9.59±2.08b	9.46±1.95b	9.027	<0.0001	S

Keys: S=Significant, NS=Not Significant, FBS=Fasting Blood Sugar, HbA1c= Glycated Haemoglobin, PostHoc (Dunnets Multiple Comparison test): Values within same row with different superscripts differ significantly compared against the control P=.05

Table 2. Distribution of FTO genotypes ACCORDING to tribes

	AA	GA	GG	AC	CC	TT	AT
Ijaw	17	3	17	5	15	14	3
Urhobo	18	2	18	5	15	14	3
Ika	18	2	19	5	15	16	3
Igbo	19	1	18	2	18	17	2
p-value	0.992	0.682	0.990	0.662	0.934	0.931	0.273
χ2-value	100.00	1.500	0.111	1.588	0.429	0.443	0.965

Table 3. Hardy-weinberg statistical test of FTO gene variants of FTO gene in T2D subjects and control

FTO gene Variant	rs2388405			rs201041270		
Genotype	AA	GA	GG	GG	GA	AA
No, Obs	92	8	0	92	8	0
Freq, Allele	0.97	-	0.03	0.97	-	0.03
Freq, Exp	0.94	0.06	0.0009	0.94	0.06	0.0009
No, Exp	92.12	5.88	0.09	92.12	5.88	0.09
Deviation	-0.12	0.12	-0.09	-0.12	2.12	-0.09
X2	0.0002	0.002	0.99	0.0002	0.002	0.99

Key: A|G=wild (Dominant), G|A=Polymorphic, X2 =chi-square, Exp=Expected, Obs=Observations

Table 4. Hardy-weinberg statistical test of some FTO gene variants in T2D subjects and control

FTO gene Variant	rs9939609			rs8050136		
Genotype	TT	AT	AA	CC	AC	AA
No, Obs	77	14	9	81	18	1
Freq, Allele	0.85	-	0.15	0.91	-	0.09
Freq, exp	0.72	0.26	0.02	0.83	0.16	0.008
No, Exp	70.6	25.48	1.96	81.34	15.68	0.8
Deviation	6.4	-11.48	7.04	-0.34	2.32	0.2
X2	0.5	0.26	0.25	0.001	0.5	0.6

T|C=wild (Dominant), A|A=Polymorphic, X2 =chi-square, Exp=Expected, Obs=Observations

Table 5. Genotype and allele distribution of rs2388405 variant of FTO gene in obese and T2D subjects and control

Tribes	Ijaw					Urobo					Ikah					Igbo				
Genotype	T2D	ND	X2	P	OR	T2D	ND	X2	P	OR	T2D	ND	X2	P	OR	T2D	ND	X2	P	OR
	n=20	n=5				n=20	n=5				n=20	n=5				n=20	n=5			
AA	17(89.5%)	5 (100%)	-	-	-	18(94.7%)	5 (100%)	-	-	-	18 (90%)	5 (100%)	-	-	-	19 (95%)	5 (100%)	-	-	-
GA	3 (10.5%)	0 (0.0%)				2(5.2%)	0 (0.0%)				2 (10%)	0 (0.0%)				1 (5%)	0 (0.0%)			
GG	0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)			
AA	17(89.5%)	5 (100%)	0.57	0.45	0.00	18(94.7%)	5 (100%)	0.27	0.60	0.00	18 (90%)	5 (100%)	0.27	0.60	0.00	19 (95%)	5 (100%)	0.27	0.60	0.00
GA + GG	3 (10.5%)	0 (0.0%)				2 (5.2%)	0 (0.0%)				2 (10%)	0 (0.0%)				1 (5%)	0 (0.0%)			
AA+GA	20(100%)	5 (100%)	-	>0.99	-	20 (100%)	5 (100%)	-	>0.99	-	20 (100%)	5 (100%)	-	>0.99	-	20 (100%)	5 (100%)	-	>0.99	-
GG	0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)			
A	36(94.7%)	10(100%)	0.55	0.45	0.00	37(94.7%)	10(100%)	0.26	0.60	0.00	38 (95%)	10(100%)	0.52	0.47	0.00	39(97.5%)	10(100%)	0.26	0.61	0.00
G	2 (5.2%)	0 (0.0%)				1 (5.2%)	0 (0.0%)				2 (5%)	0 (0.0%)				1 (2.5%)	0 (0.0%)			

A=wild (Dominant), G=Polymorphic, T2D=Type 2 Diabetes, ND = Non- Diabetes (control), X2 =chi-square, p= chi-square -pvalue, OR= Odd Ratio

Table 6. Genotype and allele distribution of rs201041270 Variant of FTO gene in T2D subjects and control

Genotype	Ijaw					Urobo					Ikah					Igbo				
	T2D	ND	X2	P	OR	T2D	ND	X2	P	OR	T2D	ND	X2	P	OR	T2D	ND	X2	P	OR
	n=20	n=5				n=20	n=5				n=20	n=5				n=20	n=5			
GG	17 (89.5%)	5 (100%)	-	-	-	18(94.7%)	5 (100%)	-	-	-	19 (95%)	5 (100%)	-	-	-	18 (90%)	5 (100%)	-	-	-
GA	2 (10.5%)	0 (0.0%)				1 (5.2%)	0 (0.0%)				1 (5%)	0 (0.0%)				2 (10%)	0 (0.0%)			
AA	0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)			
GG	17 (89.5%)	5 (100%)	0.57	0.45	0.00	18 (94.7%)	5 (100%)	0.27	0.60	0.00	19 (95%)	5 (100%)	0.27	0.60	0.00	18 (90%)	5 (100%)	0.110	0.73	0.00
GA + AA	3 (10.5%)	0 (0.0%)				2 (5.2%)	0 (0.0%)				1 (5%)	0 (0.0%)				2 (10%)	0 (0.0%)			
GG+GA	20(100%)	5 (100%)	-	>0.99	-	20(100%)	5 (100%)	-	>0.99	-	20 (100%)	5 (100%)	-	>0.99	-	20(100%)	5 (100%)	-	>0.99	-
AA	0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)			
G	36 (94.7%)	10 (100%)	0.54	0.45	0.00	37 (94.7%)	10 (100%)	0.26	0.60	0.00	39 (97.5%)	10(100%)	0.25	0.61	0.00	38 (95%)	10 (100%)	0.52	0.47	0.00
A	2 (5.2%)	0 (0.0%)				1 (5.2%)	0 (0.0%)				1 (2.5%)	0 (0.0%)				2 (5%)	0 (0.0%)			

G=wild (Dominant), A=Polymorphic, T2D=Type 2 Diabetes, ND = Non-Diabetes (control), X2 =chi-square, p= chi-square -pvalue, OR= Odd Ratio

Table 7. Genotype and allele distribution of rs9939609 variant of FTO gene in T2D subjects and control

Tribes	Ijaw			X2	P	OR	Urobo			X2	P	OR	Ikah			X2	P	OR	Igbo			X2	P	OR		
Genotype	T2D n=20	ND n=5					T2D n=20	ND n=5				T2D n=20	ND n=5				T2D n=20	ND n=5				T2D n=20	ND n=5			
TT	14 (73.7%)	4 (80%)	0.58	0.74	-		14 (73.7%)	4 (80%)	1.09	0.57		16 (80%)	4 (80%)	1.87	0.39	-		17 (85%)	4 (80%)	1.63	0.44	-				
AT	3 (15.8%)	1 (20%)					3 (10.5%)	1 (20%)				3 (15%)	0 (0.0%)				2 (10%)	0 (0.0%)								
AA	3(10.5%)	0 (0.0%)					3 (15.8%)	0 (0.0%)				1 (5%)	1 (20%)				1 (5%)	1 (20%)								
TT	14 (73.7%)	4 (80%)	0.08	0.77	0.70		14 (73.7%)	4 (80%)	0.08	0.77	0.70	16 (80%)	4 (80%)	0.00	>0.99	1.0		17 (85%)	4 (80%)	0.07	0.78	1.41				
AT + AA	6 (26.3%)	1 (20%)					6 (26.3%)	1 (20%)				4 (20%)	1 (20%)					3 (15%)	1 (20%)							
TT+AT	17 (94.7%)	5(100%)	0.57	0.45	0.00		20 (100%)	5(100%)	-	>0.99	-	20 (100%)	5(100%)	-	>0.99	-		20(100%)	5(100%)	-	>0.99	-				
AA	3 (10.5%)	0 (0.0%)					0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)					0 (0.0%)	0 (0.0%)							
T	31 (81.6%)	9 (90%)	0.40	0.52	0.49		30 (78.9%)	9 (90%)	0.63	0.42	0.41	35 (87.5%)	8 (80%)	0.37	0.54	1.75		36 (90%)	8 (80%)	0.75	0.38	2.25				
A	7 (18.4%)	1 (10%)					8 (21.1%)	1 (10%)				5 (12.5%)	2 (20%)					4 (10%)	2 (20%)							

A=wild (Dominant), T=Polymorphic, T2D=Type 2 Diabetes, ND = Non-Diabetes (control), X2 =chi-square, p= chi-square -pvalue, OR= Odd Ratio

Table 8. Genotype and allele distribution of rs8050136 variant of FTO gene in T2D subjects and control

Tribes	Ijaw		X2	P	OR	Urobo		X2	P	OR	Ikah		X2	P	OR	Igbo		X2	P	OR
Genotype	T2D n=20	ND n=5				T2D n=20	ND n=5				T2D n=20	ND n=5				T2D n=20	ND n=5			
CC	15 (78.9%)	4 (80%)	-	-	-	15 (78.9%)	5(100%)	-	-	-	15 (75%)	5 (100%)	-	-	-	18 (90%)	4 (80%)	-	-	-
AC	5 (21.1%)	1 (20%)				5(21.1%)	0 (0.0%)				5 (25%)	0 (0.0%)				2 (10%)	1 (20%)			
AA	0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)			
CC	15 (78.9%)	4 (80%)	0.002	0.95	0.93	15 (78.9%)	5(100%)	1.26	0.26	0.00	15 (75%)	5 (100%)	1.56	0.21	0.00	18 (90%)	4 (80%)	0.37	0.53	2.25
AC + AA	5 (21.1%)	1 (20%)				5(21.1%)	0 (0.0%)				5 (25%)	0 (0.0%)				2 (10%)	1 (20%)			
CC+AC	20 (100%)	5 (100%)	-	>0.99	-	20(100%)	5(100%)	-	>0.99	-	20 (100%)	5 (100%)	-	>0.99	-	20 (100%)	5 (100%)	-	>0.99	-
AA	0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)				0 (0.0%)	0 (0.0%)			
C	34 (89.5%)	9(90%)	0.002	0.96	0.94	34 (89.5%)	10(100%)	1.14	0.28	0.00	35 (87.5%)	10 (100%)	1.38	0.23	0.00	38 (95%)	9 (90%)	0.35	0.55	2.11
A	4 (10.5%)	1 (10%)				4 (10.5%)	0 (0.0%)				5 (12.5%)	0 (0.0%)				2 (5%)	1 (10%)			

C=wild (Dominant), A=Polymorphic, T2D=Type 2 Diabetes, ND = Non-diabetes (control), X2 =chi-square, p= chi-square -pvalue, OR= Odd Ratio

observed in the studied population suggests that SNPs exist in the studied population allele-specific expression is less often observed in regions of low recombination [20], allele increase may be due to SNP that occurs in the genes of members of a population [21,22].

4. CONCLUSIONS

The findings from this investigation showed that genotype AA and GG has the highest level of occurrence in all tribes of the studied population. Analysis and comparison of genotype and allele obtained within the four variants of FTO gene analysed for sample size showed that Genotype AA in rs2388405, Genotype GG in rs201041270, Genotype TT in rs9939609, and Genotype CC in rs8050136 had the highest number of occurrences in all tribes. Statistically significant genetic drifts exist in the studied population, which has caused a deviation from expected frequencies in Hardy-Weinberg statistics. This correlates with higher numbers observed in recombinant genotypes rs2388405, rs201041270, rs9939609 and rs8050136, than the originating genotypes as well as notable odd ratios of 2.25 in Igbo tribe and a significant allele with x^2 of 0.002 and odd ratio of C in rs8050136.

CONSENT

All authors declare that written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images.

ETHICAL APPROVAL

Ethical approval was sought and obtained from the Ethical Committee of Federal Medical Centre, Asaba, Delta State, Nigeria.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization Types of Diabetes, Classification of diabetes: Geneva, Licence:CC BY-NC-SA 3.0 IGO; 2019.
2. Amita B, Sara EP. DNA Methylation and its Role in the Pathogenesis of Diabetes, Pediatric Diabetes. 2017;18(3):167-177.

3. Chukwuma M. World Diabetes Day (WDD): Over 10m Nigerians have diabetes, kills more than COVID-19, cancer. The Guardian; 2022. Available:<https://guardian.ng/news/world-diabetes-day-wdd-over-10m-nigerians-have-diabetes-kills-more-than-covid-19-ca>.
4. Loos RJ, Yeo GS. The bigger picture of FTO—the first GWAS identified obesity gene. *Nature Reviews Endocrinology*. 2014;10(1):51-61.
5. Diniz IG, Noce RRD, Pereira AP, da Silva ANLM, Sacuena ERP, Lemes RB, Cardoso-Costa G. L, Araújo GS, Machado JLP, Figueiredo FAL, Hümemeier T, Guerreiro JF. Common BMI and diabetes-related genetic variants: A pilot study among indigenous people in the Brazilian Amazon. *Genetics and Molecular Biology*. 2022;45(2):1-11.
6. Tamer B, Adlija Č, Tanja D, Maja M, Velija-Asim Z, Besim P,
7. Janja M, Jana N, Vladimír P, Sabina S. Association of FTO Gene Variant (rs8050136) with Type 2 Diabetes and Markers of Obesity, Glycaemic Control and Inflammation, *Journal of Medical Biochemistry*. 2019;38(2):153–163.
8. Surendran S, Jayashri R, Drysdale L, Bodhini D, Lakshmipriya N, Shanthi RCS, Sudha V, Lovegrove JA, Anjana RM, Mohan V, Radha V, Pradeepa R, Vimalaswaran KS. Evidence for the association between FTO gene variants and vitamin B12 concentrations in an Asian Indian population. *Genes & Nutrition*. 2019;14:26.
9. Dele O. How Ijaw can produce Delta gov – Community leader; 2021. Available:<https://punchng.com/how-ijaw-can-produce-delta-gov-community-leader/>.
10. Pourhoseingholi MA, Vahedi M, Rahimzadeh M. Sample size calculation in medical studies. *Gastroenterology and hepatology from bed to bench*. 2013;6(1):14–17.
11. Adeyemo A, Chen G, Zhou J, Shriner D, Doumatey A, Huang H, Rotimi C. FTO genetic variation and association with obesity in West Africans and African Americans. *Diabetes*. 2010;59(6):1549-1554.
12. Prathima M, Reshma S, Sushith S, Shetty P, Dsa J, Kalal B. & Ramarajan M. Estimation of Glycated Haemoglobin by Nephelometry, Ion Exchange Resin and High Performance Liquid Chromatography:

- A Cross-sectional Study, *Journal of Clinical and Diagnostic Research*. 2020; 14(10):44408-139680.
13. Reed J, Bain S, Kanamarlapudi V. A review of current trends with type2 diabetes epidemiology, aetiology, pathogenesis, treatments and future perspectives. *Diabetes, metabolic syndrome and obesity: targets and therapy*. 2021;3567-3602.
 14. Sanches JM, Zhao LN, Salehi A, Wollheim CB, Kaldis P. Pathophysiology of type 2 diabetes and the impact of altered metabolic interorgan crosstalk. *The FEBS Journal*. 2021;290(3):620-648.
 15. JOUR K, Vani AR. Correlation of glycated haemoglobin with fasting and post prandial blood glucose in Type 2 diabetes. *International Journal of Clinical Biochemistry and Research*. 2020; 12274-2394-6377.
 16. Shinozaki K, Okuda M. The effects of fat mass and obesity-associated gene variants on the body mass index among ethnic groups and in children and adults. *Indian Journal of Endocrinology and Metabolism*. 2012;16(3):588-95.
 17. Oyeyemi BF, Ologunde CA, Olaoye AB, Alamukii NA. FTO gene associates and interacts with obesity risk, physical activity, energy intake, and time spent sitting: pilot study in a Nigerian population. *Journal of obesity*; 2017.
 18. Diniz IG, Noce RRD, Pereira AP, da Silva, ANLM, Sacuena ERP, Lemes RB, Cardoso-Costa, GL, Araújo GS, Machado JLP, Figueiredo FAL, Hümemeier T, Guerreiro JF. Common BMI and diabetes-related genetic variants: A pilot study among indigenous people in the Brazilian Amazon. *Genetics and Molecular Biology*. 2022;45(2):1-11.
 19. Abramovs N, Brass A, Tassabehji M. Hardy-Weinberg Equilibrium in the Large Scale Genomic Sequencing Era. *Frontiers in Genetics*. 2020;13;11:210.
 20. Prasad RB, Groop L. Genetics of Type 2 Diabetes—Pitfalls and Possibilities. *Genes*. 2015; 6(1):87–123.
 21. Michelle PH, Isabel A, Hilary E, Mawusse A, Philip A. Human Genetics; Recombination affects allele-specific expression of deleterious variants in human populations. *Science Advances*. 2022;8:19. DOI: 10.1126/sciadv.abl38
 22. Shaikh F, Shah T, Madkhali NAB, Gaber A, Alsanie WF, Ali S, Ansari S, Rafiq M, Sayyed RZ, Rind NA, Rind KH, Shar AH, Asdaq SMB. Frequency distribution and association of Fat-mass and obesity (FTO) gene SNP rs-9939609 variant with Diabetes Mellitus Type-II population of Hyderabad, Sindh, Pakistan. *Saudi Journal of Biological Sciences*. 2021;28(8):4183-4190.

APPENDIX i

MATERIALS

Statfax–2200 ELISA Machine (Reader, Washer & Incubator) by Awareness Technology, USA; MB100-4A Thermo-shaker (USA); Erba Chem 5 V3 by ERBA Diagnostics Mannheim GmbH, Germany; 721-Vis Spectrophotometer (Scottfield, England); Hettich Rotofix 32A Centrifuge by Hettich Germany; ELMI Centrifuge & Vortex, USA; Minispin Centrifuge, Eppendorf (Hamburg), Thermostat; Kiptrack blood glucose machine (Taiwan); Grant T100 water bath (UK); Illumin NextSeq 2000 platform; 10ml Syringe & Needle; Vacutainer-Type EDTA & Plain Tubes, Hand Gloves, Alcohol Swabs, Tourniquet; Sharp Container; Eppendorf Tube; Cryo Tube; Pen; Marker; Semi-Auto pipette (5-20ul, 10-50, 10-100, 20-200, 100-1000uL, 10ul, 20uL, 50uL, 200uL, 1000uL); Weighing scale; Measuring tape; Refrigerator with freezer; Test tubes; Test tube rack; Timer; 1.5-ml micro centrifuge tubes or 15- or 50-ml conical polypropylene centrifuge tubes (e.g., Falcon), DNA source plate: 384-well deep-well PCR plate containing 2.5ng/μl DNA of interest, 100mM dNTPs (Applied Biosystems; store at -20°C), 25mM MgCl₂ (QIAGEN; store at -20°C), Ultrapure PCR-grade H₂O (Invitrogen), 5U/μl HotStarTaq Plus DNA polymerase with 10× PCR buffer (QIAGEN; store at -20°C), Forward and reverse primers: 1μM each in multiplex pool, 384-well PCR reaction plate (Eppendorf twin.tec), 1× SAP buffer (Sequenom), 1.7U/μl shrimp alkaline phosphatase (SAP; Sequenom; store at 0°C), PCR products in 384-well PCR plates, Ultrapure PCR-grade H₂O (Invitrogen), Extend primers: from 5 to 10μM each in multiplex pool, PCR products in 384-well PCR plates, cleaned up by SAP reaction, Ultrapure PCR-grade H₂O (Invitrogen), SpectroCLEAN resin (Sequenom; store at room temperature), Post-primer extension products in 384-well plates kept at 4°C, 100% and 50% ethanol, Plates from primer extension, cleaned up with resin and centrifuged, 3-point calibrant (Sequenom).

Sample Purification Beads (SPB), Resuspension Buffer (RSB), Freshly prepared 80% ethanol (EtOH), 96-well 0.8 ml Polypropylene Deepwell Storage Plate (midi plate) (2), 96-well PCR plate, Microseal 'B' adhesive seal, Microseal 'F' foil seal, 1.7 ml microcentrifuge tubes, Nuclease-free water, Enhanced PCR Mix (EPM), Index adapters (tubes or plate), 20 μl multichannel pipettes, 200 μl multichannel pipettes, Tagment Stop Buffer (TSB), Tagment Wash Buffer (TWB), 96-well plate magnet, Microseal 'B' adhesive seal, Bead-Linked Transposomes (BLT), Tagmentation Buffer 1 (TB1), 96-well PCR plate, 8-tube strip.

© 2024 Osere et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111330>