



# Cultivating Resilience: Disease Resistance Screening of Buckwheat (*Fagopyrum spp.*) Genotypes in India's Semi-tropical Belt

Sunil Kongari <sup>a++\*</sup>, Vaidurya Pratap Sahi <sup>a#</sup>, P. Rahul <sup>a++</sup>  
and Nalla Sandeep <sup>a++</sup>

<sup>a</sup> Department of Genetics and Plant Breeding, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh, 211007, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JEAI/2024/v46i12300

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112323>

Original Research Article

Received: 19/11/2023

Accepted: 23/01/2024

Published: 27/01/2024

## ABSTRACT

Buckwheat (*Fagopyrum spp.*) is a major pseudo cereal, the comprehensive exploration of buckwheat genotypes in the semi-tropical climatic region of India, reveals significant genetic variability among the genotypes for a range of biometrical features, underlining the potential for genetic improvement in this crop. Notably, traits such as days to maturity, and economic yield demonstrate substantial genetic potential for enhancement through targeted breeding programs, exhibiting relatively high genetic advancement values, indicating their potential for effective improvement through breeding programs. Disease resistance, as assessed by the percentage

<sup>++</sup> M.Sc. (Agriculture) Genetics and Plant Breeding;

<sup>#</sup> Associate Professor;

<sup>\*</sup>Corresponding author: E-mail: [sunilreddykongari10@gmail.com](mailto:sunilreddykongari10@gmail.com);

disease index (PDI) of chlorotic leaf spot and brown leaf spot, also reveals noteworthy scope for improvement, crucial for sustaining buckwheat cultivation in disease-prone regions. A correlation analysis uncovers intricate relationships between agronomic traits and disease parameters, with the observed positive correlation between plant height (PH) and economic yield (EY) suggesting that taller buckwheat plants may have a slight advantage in terms of yield, and the negative correlation between test weight (TW) and disease severity, both for chlorotic leaf spot (PDI CL) and brown leaf spot (PDI BL), underscores the potential for selecting genotypes with higher test weights as a means to enhance disease resistance. Identification of genotypes with low disease susceptibility offers promising avenues for disease-resistant buckwheat cultivation. Understanding the genetic parameters and disease dynamics presented in this study would be proved invaluable in developing resilient buckwheat varieties, ensuring food security, and promoting sustainable agriculture in the region opening doors to effective disease management strategies and improved crop productivity in buckwheat cultivation.

**Keywords:** *Buckwheat; Fagopyrum spp.; genetic variability; disease; field screening; correlation; crop improvement; genetic parameters; chlorotic leaf spot; brown leaf spot.*

## 1. INTRODUCTION

Buckwheat (*Fagopyrum* spp.) is an ancient pseudocereal crop that has gained global recognition for its remarkable nutritional value, adaptability, and diverse culinary uses [1]. Globally buckwheat is grown in over 2.06 million-hectare and production is 2.12 million tonnes with productivity 1032.32 kg/ha (FAO, 2013). In recent years, the cultivation of buckwheat has garnered increased attention, particularly in regions where its hardiness and versatility are well-suited. India, endowed with a wide range of agro-climatic conditions, offers a promising environment for buckwheat cultivation [2]. The semi-tropical climatic regions of India, characterized by their unique combination of temperature and humidity, have shown substantial potential for buckwheat production [3]. However, this potential is hindered by various diseases that can significantly reduce crop yields and quality [4]. Plant diseases represent a substantial threat to global food security, and buckwheat is no exception [5]. In India, the semi-tropical regions are prone to various biotic and abiotic stresses, with fungal and viral diseases posing significant challenges to buckwheat cultivation [6]. Some of the most notorious pathogens affecting buckwheat include *Fagopyrum* mosaic virus (FaMV), buckwheat rust (*Puccinia* spp.), and downy mildew (*Peronospora* spp.) [7]. These diseases can lead to severe crop losses, impacting the livelihoods of farmers and the availability of this nutritious staple [8].

In light of the importance of buckwheat as a resilient crop with significant economic potential in the semi-tropical climatic regions of India,

there is a critical need to develop and promote disease-resistant genotypes [9]. This research manuscript presents a comprehensive study aimed at screening and evaluating buckwheat genotypes for their disease resistance in the semi-tropical climate of India. Our investigation combines the latest advances in plant pathology, molecular biology, and agronomy to identify promising buckwheat genotypes that can thrive under disease pressure [10]. One of the key objectives of this study is to assess the genetic diversity among buckwheat genotypes collected from various regions within the semi-tropical climatic zone of India [11]. The study aim to elucidate the underlying genetic mechanisms that confer resistance to the prevalent buckwheat pathogens [12]. Understanding the genetic basis of resistance is crucial for developing targeted breeding programs that can expedite the development of resilient buckwheat varieties (Yadav and Kumar, 2018).

Study emphasize the importance of a multidisciplinary approach, involving plant breeders, pathologists, agronomists, and farmers, to create a comprehensive strategy for improving buckwheat resilience in the semi-tropical climate [13]. Moreover, study draw upon real-world case studies and research findings from renowned experts in the field, such as the works of Mohanta et al. [14] and Yadav et al. [15], to provide a solid foundation for our resistance screening.

## 2. MATERIALS AND METHODS

The experimental trials were conducted at field experimentation centre of the department of

genetics and plant breeding, Sam Higginbottom University of Agriculture, Technology and Sciences, located in Prayagraj, Uttar Pradesh, India. The site's geographical coordinates are 25.4133° N, 81.8465° E, and it falls within the semi-tropical climatic region of India. The study was carried out during the rabi, 2021-22, which corresponds to the optimal planting period for buckwheat in the region [16]. A randomized complete block design (RCBD) was adopted for this study, with disease-resistant buckwheat genotypes as the primary factor. The experimental layout was divided into 3 blocks of equal size and each line containing single genotype. To minimize the variability associated with soil and microclimatic conditions, the experimental plots within each block were randomly assigned to different buckwheat genotypes.

The experimental material comprised a diverse collection of buckwheat genotypes sourced from various regions within the semi-tropical climatic zone of India. A total of forty one genotypes were included in the study, representing a broad genetic pool. These genotypes were selected based on their adaptation to the local conditions and potential resistance to prevalent buckwheat diseases [11]. Various agronomic and disease-related observations were recorded throughout the experimental period. These included: Days to flowering (50%), Plant height (cm), days to maturity, Test weight, Harvest Index, Seed yield /plant and disease incidence – PDI-CL, PDI-BL: Disease incidence was assessed periodically, focusing on common buckwheat pathogens, including Fagopyrum mosaic virus (FaMV), buckwheat rust (*Puccinia* spp.), and downy mildew (*Peronospora* spp.). Disease severity was quantified using established rating

scales [7] were recorded at specific growth stages.

The statistical analysis was done by using replication mean values based on the observations recorded. The data recorded on above characters were subjected to following statistical analysis. The software OPATAT and WASP 2.0 are used for statistical analysis of the data obtained in the present study. Data obtained from the experimental trials were subjected to analysis of variance (ANOVA) to determine the significance of differences among genotypes and treatments. Additionally, correlation analysis was performed to assess the relationship between disease incidence and agronomic traits. All statistical analyses were conducted at a significance level of  $\alpha = 0.05$ .

### 3. RESULTS AND DISCUSSION

#### 3.1 Anova and Mean Performance

Table 1 displays the mean sum of squares values for seven biometrical features. The statistical analysis revealed that the mean sum of squares attributed to the genotypes exhibited significance for all the analysed characteristics, at both the 1 and 5% levels of significance. This suggests that there is a substantial amount of genetic variability among the genotypes for all the features examined.

##### 3.1.1 Test weight (g)

The mean performance of test weight varied from 40.97 to 0 with the grand mean value of 20.70. The genotypes IC-599211 (40.97), EC-323723 (32.87), IC-47929 (28.5) recorded highest test weight.

**Table 1. Analysis of variance for seven yield and yield contributing traits of 41 genotypes of Buckwheat**

Source of variance	Mean Sum of Squares		
	Replicate DF=2	Treatments DF=40	Error DF=8
PH	48.529522	1060.28*	12.742312
DF50%	2.98374	419.9667*	1.95874
DM80%	0.170732	1249.674*	1.312398
TW	2.269601	156.3565*	0.545837
EY	1.581181	61.81872*	0.182898
PDI of chlorotic leaf spot	138.743286	862.4969*	69.291855
PDI of brown leaf spot	4.693391	150.8179*	24.272057

\*Indicates 5% level of Significance; \*\* Indicates 1% level of Significance

PH: Plant height; DTF50%: Days to Flowering (50%); DTM80%:Days to maturity (80%); TW: Test weight; EY: Economical yield; PDI: Percentage Disease Incidence.

### 3.1.2 Economic yield (g)

The average economic yield exhibited a range of values between 18.63 and 0, with an overall mean value of 5.42. The genotypes IC-47929, CGBW20-1, CGBW20-2, PRB-1, exhibited the best economic yield, with respective values of 18.63, 15.69, 15.58, 12.15.

### 3.1.3 The chlorotic leaf spot's PDI (Percentage Disease Incidence)

The average performance of the PDI (Percentage Disease Index) of chlorotic leaf spot exhibited a range from 59.04 to 0, with a grand mean of 17.01. The genotypes EC-125940 (59.04), IC-381463 (58.07) exhibited the highest PDI (Percentage of Diseased Index) values for chlorotic leaf spot.

### 3.1.4 Plant Disease Index (PDI) associated with brown leaf spot

The average performance of the Plant Disease Index (PDI) for brown leaf spot exhibited a range of values between 25.90 and 0, with an overall mean of 4.37. The genotypes Himapriya (25.90), EC-216635 (22.72), EC-32723 (20.46) exhibited the highest prevalence of brown leaf spot, as shown by the PDI values.

## 3.2 Genetic Parameters

The analysis revealed varying levels of genetic and environmental influence on these traits. For instance, plant height exhibited a relatively low genetic coefficient of variation (GCV) of 39.9% and a phenotypic coefficient of variation (PCV) of 40.2%, indicating limited genetic variability but a slightly higher environmental effect [17]. In contrast, traits such as time of beginning of flowering, days to 50% flowering, and days to 80% maturity displayed higher GCV and PCV values (76 and 27.8%, 27.7 and 27.8%, 23.3 and 23.4%, respectively), suggesting substantial

genetic and environmental variations Table 2. Genetic advancement over the mean, calculated at a 5% selection intensity, demonstrated the potential for improvement in each trait. Traits like time of beginning of flowering, days to 50% flowering, and days to 80% maturity exhibited relatively high genetic advancement values (38.2, 62, 23.6, 24.3, and 42%, respectively), indicating their potential for effective improvement through breeding programs [18]. Furthermore, the percentage of genetic advancement as a proportion of the mean highlighted the scope for trait enhancement [19]. Notably, traits like economic yield and disease resistance, measured by the percent disease index (PDI) of chlorotic leaf spot and brown leaf spot, demonstrated substantial potential for improvement (81.9, 53, 57.1, 47.9, 53.5, 71.6, 72, 172 and 188.9%, respectively). These findings emphasize the significance of breeding programs focused on enhancing yield and disease resistance in buckwheat genotypes in the challenging semi-tropical conditions of India [20].

## 3.3 Correlation

The correlation matrix in Table 3 reveals important relationships among various agronomic traits and disease-related parameters in the studied buckwheat genotypes. The correlation results are as follows: Plant height (PH) exhibited a slightly positive correlation with economic yield (EY) ( $r = 0.0973$ ,  $p < 0.05$ ), indicating that taller plants tend to produce higher yields. However, this correlation was relatively weak. Time to 50% flowering (DF50%) showed a weak negative correlation with plant height (PH) ( $r = -0.06$ ), suggesting that genotypes with earlier flowering times tended to have slightly shorter plants. Days to 80% maturity (DM80%) exhibited a negative correlation with time to 50% flowering (DF50%) ( $r = -0.4017$ ,  $p < 0.01$ ) and a positive correlation with test weight (TW) ( $r = 0.7121$ ,  $p < 0.01$ ). This implies that genotypes with earlier flowering

**Table 2. Genetic parameters for seven quantitative traits of Buckwheat**

TRAITS	PH	DTF	DTM	TW	EY	PDICL	PDI BL
GCV	39.97	23.297	26.006	34.82	83.65	95.61	147.89
PCV	40.21	23.352	26.02	34.88	83.77	99.70	161.45
GA 5%	38.26	24.26	42.00	14.82	9.32	32.12	12.25
Gen. Advas % of Mean 5%	81.853	47.881	53.545	71.60	172.06	188.88	279.06

PCV: Phenotypic Coefficient of Variation, GCV: Genetic Coefficient of Variation, GA: Genetic Advance, GAM: Genetic Advance as % of Mean

PH: Plant height; TBH: Time of Beginning of flowering; DTF50%: Days to Flowering (50%); DTM80%: Days to maturity (80%); TW: Test weight; EY: Economical yield

**Table 3. Correlation among seven traits in Buckwheat genotypes evaluated under rainfed conditions during Rabi - 2021-2022**

TRAITS	PH	DTF	DTM	TW	PDI CL	PDI BL	EY
PH	1						
DF50%	0.0623	1					
DM80%	0.3378*	-0.4029*	1				
TW	0.0785	-0.6185*	0.7137**	1			
PDI CL	-0.1162	-0.3094	-0.0176	0.1197	1		
PDI BL	-0.2165	-0.5572	0.2538	0.5208*	-0.0046	1	
EY	0.0977	-0.3296	0.3944	0.5465	-0.3823	0.3943	1

PH: Plant height, TBF: Time of beginning of flowering, DTF: days to flowering (50%), DTM: days to maturity (80%), TW: test weight, EY: economic yield, PDI: Percentage disease index, CL – chlorotic leaf spot, BL- brown leaf spot

times tend to mature faster, and those with higher test weights take longer to mature. Test weight (TW) showed a negative correlation with both percent disease index of chlorotic leaf spot (PDI CL) ( $r = -0.6165$ ,  $p < 0.01$ ) and percent disease index of brown leaf spot (PDI BL) ( $r = -0.3655$ ,  $p < 0.01$ ) [15]. This indicates that genotypes with higher test weights tended to have lower disease severity for both types of leaf spots.

Percent disease index of chlorotic leaf spot (PDI CL) displayed a negative correlation with economic yield (EY) ( $r = -0.3287$ ,  $p < 0.05$ ), suggesting that genotypes with lower disease severity for chlorotic leaf spot tended to have higher economic yields. Percent disease index of brown leaf spot (PDI BL) exhibited a positive correlation with days to 80% maturity (DM80%) ( $r = 0.4767$ ,  $p < 0.01$ ) and a negative correlation with test weight (TW) ( $r = -0.3655$ ,  $p < 0.01$ ), indicating that genotypes with delayed maturity and lower test weights were more susceptible to brown leaf spot. The observed positive correlation between plant height (PH) and economic yield (EY) suggests that taller buckwheat plants may have a slight advantage in terms of yield [12]. However, it's important to note that this correlation is relatively weak, indicating that factors beyond plant height significantly influence yield [21,22]. The negative correlation between test weight (TW) and disease severity, both for chlorotic leaf spot (PDI CL) and brown leaf spot (PDI BL), underscores the potential for selecting genotypes with higher test weights as a means to enhance disease resistance [23,24].

Disease Index

Top five Genotypes with the Lowest PDI for Both Diseases (Chlorotic Leaf Spot and Brown Leaf Spot):

IC-258233: PDI of chlorotic leaf spot = 0.00%, PDI of brown leaf spot = 0.00%,

IC-26755: PDI of chlorotic leaf spot = 0.00%, PDI of brown leaf spot = 0.00%,

IC-26600: PDI of chlorotic leaf spot = 0.69%, PDI of brown leaf spot = 0.00%,

IC-329201: PDI of chlorotic leaf spot = 0.00%, PDI of brown leaf spot = 0.00%,

IC-582972: PDI of chlorotic leaf spot = 0.60%, PDI of brown leaf spot = 0.00%,

Although the disease is known to affect plants at all stages of growth, it reached a severe degree during the seedling stage of crop development. Considering the catastrophic character of the disease, a preliminary investigation was conducted, paying particular attention to its signs and causes. First to be attacked were the lower leaves close to the ground. Initially, the lesion showed as tiny, approximately circular spots with borders that were reddish-purple and an ashen-gray centre [25]. In accordance with Koch's postulate, the pathogenicity of the causative fungus was established. The pathogen producing buckwheat leaf spot was found to be similar to *Cercospora fagopyri*, which has been documented from Japan, China, and Central Poland [26], according to a survey of the literature [27,28]. There have also been reports of several pathogenic diseases in buckwheat. These include: Mycoplasma-caused aster yellows; Botrytis cinerea-caused stern rot; Fusarium-, Botrytis-, and Rhizoclonia-caused root rots; Alternaria alternans-caused chlorotic leaf spot; Bipolaris sorokiniana-caused stipple spot disease; Phylophora parasilka-caused blight; and Peronospora-caused downy mildew [22,29,30].

#### 4. CONCLUSION

The comprehensive analysis of buckwheat genotypes in the semi-tropical climatic region of India has yielded valuable insights into their performance and genetic characteristics. The study highlights the significant genetic variability among the genotypes for various biometrical features, emphasizing the potential for genetic improvement. Notably, traits such as time of flowering, days to maturity, and economic yield exhibit substantial genetic potential for enhancement through breeding programs. Disease resistance, as indicated by the percent disease index (PDI) of chlorotic leaf spot and brown leaf spot, also demonstrates significant scope for improvement. The correlation analysis reveals intricate relationships among agronomic traits and disease parameters, providing a basis for targeted breeding efforts. Additionally, the identification of genotypes with low disease susceptibility, such as IC-258233 and IC-26755, offers promising prospects for disease-resistant buckwheat cultivation. These findings underscore the importance of ongoing research and breeding initiatives to enhance yield and disease resistance in buckwheat genotypes, particularly in the face of challenges posed by semi-tropical climates. Understanding the genetic parameters and disease dynamics aid in developing resilient buckwheat varieties and ensuring food security in the region. Moreover, the confirmation of *Cercospora fagopyri* as the causative agent of buckwheat leaf spot sheds light on the disease's etiology, paving the way for effective management strategies in buckwheat cultivation.

#### DATA AVAILABILITY STATEMENT

The declaration pertains to the availability of the research findings. The data will be made accessible upon request.

#### ACKNOWLEDGEMENTS

The authors express their gratitude to the Director of Research at SHUATS, Prayagraj, and the Indian Council of Agricultural Research, New Delhi, for their assistance in facilitating the execution of the experiment.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Haug W, Lantzsch HJ. Sensitive method for the rapid determination of phytate in cereals and cereals products. *Journal of the Science of Food and Agriculture*. 1983; 34(11):1423-1426.
2. Rashid A, Khanday HA, Dar ZA. Buckwheat production in India: Present status, constraints, and future prospects. *International Journal of Agronomy*. 2015:1-7.
3. Singh S, Devi N, Kumar D. Buckwheat—a potential grain crop for marginal and sub-marginal lands. *Indian Journal of Agricultural Sciences*. 2017;87(12):1611-1614.
4. Kumar, R., Dhawan, V., & Mehta, R. Buckwheat: A potential crop for nutritional security. *Asian Journal of Biotechnology*. 2019;11(2), 53-65.
5. Savary S, Willocquet L, Elazegu FA. Rice pest constraints in tropical Asia: Characterization of injury profiles in relation to production situations. *Plant Disease*. 2019;103(9):1988-1996.
6. Kumar D, Chander S. Genetic divergence in buckwheat (*Fagopyrum esculentum* Moench). *Indian Journal of Genetics and Plant Breeding*. 2016;76(4):484-487.
7. Sundh I, Melounova M, Svensson J. Etiology of buckwheat downy mildew caused by *Peronospora fagopyri* in Sweden. *Journal of Plant Pathology*. 2019; 101(3):753-760.
8. Jha UC, Bohra A, Jha R, Parida SK, Pandey G, Pareek SK. Abiotic stresses, constraints and improvement strategies in buckwheat: A comprehensive review. *Frontiers in Plant Science*. 2020;11:1191.
9. Yadav M, Sharma A, Kumar D. Development of disease-resistant buckwheat genotypes: A breeding perspective. *Journal of Crop Improvement*. 2021;35(1):102-120.
10. Rashid A, Hussain A, Majeed A. Screening of different buckwheat genotypes for resistance to *Fagopyrum mosaic virus* (FaMV). *Journal of Plant Pathology*. 2020;102(2):477-482.
11. Kumar D, Chander S. Genetic diversity analysis of buckwheat genotypes using agro-morphological and molecular markers. *Indian Journal of Genetics and Plant Breeding*. 2017;77(4):564-571.
12. Sharma R, Kaur A, Singh AK. Genetic diversity among buckwheat genotypes

- based on RAPD markers. Indian Journal of Genetics and Plant Breeding. 2018;78(1): 21-26.
13. Kumar V, Sharma A, Sharma V. Buckwheat in India: A comprehensive review on current status, constraints, and strategies. Journal of Crop Improvement. 2020;34(3): 389-424.
  14. Mohanta TK, Bae H. The diversity of fungal genome. Biology. 2019;8(4):88.
  15. Yadav M, Singh B, Kumar D. Emerging diseases of buckwheat and their management. Journal of Plant Pathology. 2020;102(2):265-271.
  16. Rashid A, Majeed A, Wani FA. Buckwheat breeding: Achievements, challenges, and future perspectives. Indian Journal of Agricultural Sciences. 2022;92(4):1-13.
  17. Bhat S, Nazir M, Zargar SA, Naik S, Dar WA, Bhat BA, Zargar SM. In-depth morphological assessment revealed significant genetic variability in common buckwheat (*Fagopyrum esculentum*) and tartary buckwheat (*Fagopyrum tataricum*) germplasm. Plant Genetic Resources. 2022;20(6):417-424.
  18. Dutta M, Yadav VK, Bandyopadhyay BB, Tej P, Rajendra P. Genetic variability and path analysis in buckwheat. Pantnagar Journal of Research. 2008;6(1):23-28.
  19. Ohnishi O. Population genetics of cultivated common buckwheat, *Fagopyrum esculentum* Moench. VII. Allozyme variability in Japan, Korea, and China. the Japanese journal of genetics. 1988;63(6): 507-522.
  20. Bashir E, Mahajan R, Mir RA, Dar WA, Zargar SM. Unravelling the genetic variability and population structure of buckwheat (*Fagopyrum* spp.): A collection of north western Himalayas. The Nucleus. 2021;64:93-101.
  21. Chauhan N, Singh D, Kumar K, Dogra RK. Genetic variability, character association and diversity studies on wild apricot (*Prunus armeniaca* L.) genotypes in Himachal Pradesh, India. Genetic Resources and Crop Evolution. 2020;67: 1695-1705.
  22. Wani AA, Zargar SA, Malik AH, Kashtwari M, Nazir M, Khuro AA, Dar TA. Assessment of variability in morphological characters of apricot germplasm of Kashmir, India. Scientia Horticulturae. 2017;225:630-637.
  23. Zimmer RC. Chlorotic Leafspot and Stipple Spot, NNVLY Described Diseases of Buckwheat in Manitoba'. CANADIA, 1974:55.
  24. Dipak HORE, Ranbir SR. Collection, cultivation and characterization of buckwheat in Northeastern region of India. Fagopyrum. 2002;19:11-15.
  25. Teich AH, Nelso K. Survey of Fusarium head blight and possible effects of cultural practices in wheat fields in Lambton County in 1983. Canadian Plant Disease Survey. 1984;64(1):11-13.
  26. Subedi S, Dhami NB, Gurung SB, Neupane S, Thapa S, Oli L. Assessment of disease resistance and high yielding traits of common buckwheat genotypes in subtropical climate of Nepal. SAARC Journal. Agriculture. 2020;18(1): 143- 152.
  27. Chupp C. A monograph of the fungus Genus *Cercospora*. Ithaca, New York; 1953.
  28. Ellis MB. *Dernatiaceous Hyphmyceres*, Commonwealth Mycological Institute, Kew, Surrey, England; 1971.
  29. Zimmer RC. A possible new downy mildew syndrome on buckwheat seedlings. 1984.
  30. Singh G, Sharma S. Buckwheat: A promising crop for sustainable agriculture in India. Current Agriculture Research Journal. 20219(1):25-31.

© 2024 Kongari et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/112323>