



# Evan's Syndrome Hemolytic Anemia with Thrombocytopenia in a Labrador Dog: A Case Report

Bhand Akshata Chandrakant <sup>a\*</sup>, Ravi Mohan Shukla <sup>b</sup>,  
Siddharth Chaudhary <sup>a</sup>, Garima Rathore <sup>a</sup>, Chand Tanwar <sup>a</sup>,  
Syed A Arif <sup>a</sup> and Indu Yadav <sup>c</sup>

<sup>a</sup> Division of Medicine, Indian Veterinary Research Institute, Izatnagar, Uttar Pradesh-243122, India.

<sup>b</sup> Department of Pathology, Veterinary College and Research Institute, Orathanadu, TANUAVAS- 61462, India.

<sup>c</sup> Division of Parasitology, Indian Veterinary Research Institute, Izatnagar, Uttar Pradesh-243122, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.56557/UPJOZ/2024/v45i73968

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.mbimph.com/review-history/3320>

**Case Report**

**Received: 12/01/2024**  
**Accepted: 16/03/2024**  
**Published: 22/03/2024**

## **ABSTRACT**

A seven-year-old Labrador retriever intact male dog was presented to the Referral Veterinary Polyclinic, Indian Veterinary Research Institute, Izatanagar with a history of inappetence, melena, and petechial hemorrhages on the ventral abdomen and treated by a local veterinarian for a prolonged period of time. Clinical examination showed pale mucous membranes, peripheral lymphadenopathy, and mild hepatomegaly. A Complete blood count revealed anemia, leukopenia, and thrombocytopenia and serum biochemistry showed hypoalbuminemia and hypoproteinemia. The dog was suspected of having a haemoprotozoan infection, and a thin peripheral blood smear was submitted to the Division of Parasitology, Indian Veterinary Research Institute, Izatanagar,

\*Corresponding author: Email: akshatabhand0807@gmail.com;

Bareilly, for examination. A blood smear was screened for haemoprotozoan examination using Giemsa's stain, which showed positivity for *Babesia vogeli*. An abdominal ultrasound confirmed hepatic enlargement. The blood clotting profile reveals an increase in activated partial thromboplastin time and prothrombin time. The dog's blood sample tested negative for *Babesia vogeli*, *B. gibsoni*, *E. canis*, and *H. canis* by multiplex PCR; this could be because of prior initiation of antibiotics. A perihelal blood smear showed elevated spherocytes count and saline agglutination test was positive. Based on history, clinical examinations, laboratory findings and cytological examinations the case was tentatively diagnosed as chronic case of tick fever with concomitant Evans syndrome. A blood transfusion was performed and around 300 ml of whole blood was infused. Treatment was initiated with triple antibiotic therapy and an immunosuppressive dose of steroids, but after 4 weeks, the dog succumbed to the condition. Further research is required to optimize treatment regimens for dogs with Evan's syndrome, as the prognosis is uncertain.

**Keywords:** Evans syndrome; dog; hemolytic anemia; Thrombocytopenia; etc.

## 1. INTRODUCTION

One of the most fatal haemoprotozoan infections in dogs that is spread by ticks is babesiosis. The most common Babesia species in dogs are *B. canis*, *B. vogeli*, *B. rossi* and *B. gibsoni*, which are intracellular protozoan parasites of canine erythrocytes [1]. *B. canis* is indigenous to Southern Europe, America, Asia, and South Africa, but *B. gibsoni* is found in the Middle East, Northern Africa, and South Asia [2]. Babesia infection produces autoimmune disorders in dogs, such as immune-mediated haemolytic anaemia (IMHA) and immune-mediated thrombocytopenia (ITP), which can develop separately or concurrently. If they present together then it is called as Evans Syndrome [3]. Evan's syndrome is named after Robert Evans, who originally documented it in humans in 1951 [4]. It appears similarly in dogs, although with variable clinical presentations and findings. Evans syndrome with Babesia in canines destroys RBCs, platelets, and causes oxidative stress, and endothelial damage [5].

*B. vogeli* in a dog is spread by brown dog tick that is *Rhipicephalus Sanguineus* and is indigenous to Africa, America, Asia, Australia, and Europe [6]. *B. vogeli* is considered to be mildly pathogenic, causing subclinical or mild disease [7]. Clinical findings include depression, weakness, anorexia, fever, lymphadenomegaly, splenomegaly, anaemia, thrombocytopenia, jaundice. These symptoms are similar to those observed with other forms of Babesia [8]. Although severe infection can develop in dogs infected with other species, systemic inflammatory response syndrome (SIRS), multiple organ dysfunction syndrome (MODS), refractory hypotension and septic shock are common findings [9]. In this study, a dog infected

with *Babesia vogeli* was diagnosed conventionally, and the prevalence and control methods in India were discussed.

### 1.1 Clinical Presentation of Animal


A seven-year-old Labrador retriever intact male dog was presented with a history of inappetance, melena, lethargy and petechial hemorrhages on the ventral abdomen to the Referral Veterinary Polyclinic, Indian Veterinary Research Institute, Izatanagar and treated by a local veterinarian for a prolonged period of time. Thorough clinical examination of the dog showed enlargement of peripheral lymph node, pale conjunctival and oral mucous membrane (1A and 1B), petechial hemorrhages on the body (Fig. 1C) and hepatomegaly evident on abdominal palpation. Due to suspicion of tick fever, thin blood smear was sent of Division of Parasitology, Indian Veterinary Research Institute, Izatanagar, Bareilly for haemoprotozoan examination.

### 1.2 Laboratory Examination

A 1mL blood sample was obtained in an EDTA vial and 2mL in serum vial and sent to the laboratory. A routine haematology revealed anaemia (Hb: 2.6mg/dL, Erythrocytes:  $0.96 \times 10^6$ ) and thrombocytopenia (25,000/cmm). Serum biochemistry examination showed decrease in total protein (3.94g/dL) count while kidney values were all within normal range. Blood clotting profile revealed increase in activated partial thromboplastin time (20 second) and prothrombin time (12.2second).

### 1.3 Diagnostic Imaging

Ventral abdominal area was shaved and prepared for abdominal ultrasound. Abdominal



**Fig. 1. (A) Pale conjunctiva (B) Pale mucous membrane (C) Petechial hemorrhages on the ventral abdomen**

ultrasound examination was carried out with the help of Curvilinear transducer (2.5-5MHz frequency) showed enlargement in liver parenchyma. No other significant findings were reported on abdominal ultrasound.

## 2. MATERIALS AND METHODS

Following one minute of immersion in methanol, the thin blood smear was subjected to Giemsa's stain, diluted at a ratio of 1:10 with water to create a working solution, and left for 40 minutes. Subsequently, the slide underwent rinsing with tap water, drying, and focusing under a compound binocular light microscope with oil immersion. Microscopic examination of thin blood smear stained with Giemsa stain showed presence piroplasms of *Babesia vogeli* in the erythrocytes (Fig. 2). Spherocytes were also observed in the smear indicating autoimmune hemolytic anemia (Fig. 3).


### 2.1 Treatment Protocol

Therapy was started with Diminazene aceturate at the dose rate of 5mg/kg body weight, deep


intramuscularly and repeated after 14 days. Also triple antibiotic therapy was initiated using Doxycycline at the dose rate of 10 mg/kg body weight once daily orally for 21days, Clindamycin at the dose rate of 11 mg/kg body weight twice daily orally for 21days, and metronidazole at the dose rate of 10 mg/kg body weight twice daily orally for 21days along with Pantoprazole at the dose rate of 1 mg/kg body weight once daily orally for 21 before food. Immunosuppressive therapy with prednisolone at the dose rate of 1 mg/kg body weight once daily orally at tapering dose was initiated. Fluid therapy, styptics, hematinics, and carioca papaya extract were all used as supportive treatments. To get rid of tick infestations, Fipronil and S-Methoprene were administered topically.

### 2.2 Blood Transfusion

Heparin (5–10 units/ml of blood) was used as an anticoagulant to draw 300 ml of whole fresh blood from the donor. Blood samples from donors and recipients were compatible, as demonstrated by major and minor cross-



**Fig. 2. Piroplasm of *Babesia vogeli* in erythrocyte of dog**



**Fig. 3. Spherocytes in Giemsa stained blood smear**

matching. The recipient dog received 300 milliliters of blood over three hours. No adverse transfusion reaction was seen.

### **3. RESULTS AND DISCUSSION**

Immune-mediated thrombocytopenia (ITP), alternatively termed Evans syndrome or IMP, arises from various origins, either primary or secondary. Primary immune thrombocytopenia (IMTP) lacks a discernible underlying cause, while secondary IMTP can arise from various potential triggers, such as neoplasia, tick-borne

illnesses (like ehrlichiosis and babesiosis), certain medications (including sulfa antibiotics), vaccination, and blood transfusions. Unlike immune-mediated hemolytic anemia (IMHA) which targets red blood cells, ITP manifests with the body's immune system excessively targeting platelets, leading to their depletion at a pace surpassing their production in the bone marrow. Evan's syndrome is a rare autoimmune disorder that poses significant diagnostic and therapeutic challenges in veterinary medicine [10]. Researchers claim that 30% of animals with IMTP also develop IMHA at the same time [11].

The conclusive diagnosis of IMTP depends on the detection of anti-platelet antibodies, but these tests are not commonly accessible and cannot differentiate between primary and secondary IMTP. Hence, treatment may begin based on a presumptive diagnosis.

Immunosuppressive medications are the mainstay of IMTP treatment. The most commonly prescribed class of immunosuppressive medications is corticosteroids, predominantly prednisone at a tapering dosage of 1 to 1.5 mg/kg, every 12 hours [11]. It functions by hindering macrophage activity in platelet destruction and can also enhance capillary resilience to bleeding. Apart from utilizing immunosuppressive medications for medical treatment, administering blood transfusions could help replenish platelet levels in cases of IMTP.

Antiprotozoal therapy with triple antibiotics was used for treating underlying haemoprotozoan infection with *Babesia vogeli* [12,13]. The treatment regimen employed use of three antibiotics—clindamycin, metronidazole, and doxycycline—which has proven to be more efficacious in treating canine babesiosis. Clindamycin enhances both cellular and humoral immunity by targeting *Babesia* sp., demonstrating effectiveness against babesiosis. Doxycycline has been noted for its preventive action against *Babesia canis* infection, while metronidazole, an anti-trichomonal agent, exhibits therapeutic efficacy against *Babesia gibsoni* infection. As Evans syndrome has poor prognosis [11] animal succumb to the condition after one month.

#### 4. CONCLUSION

Despite initiation of treatment consisting of triple antibiotic therapy and immunosuppressive doses of steroids along with a blood transfusion, the dog's condition deteriorated, and it ultimately succumbed to the illness after four weeks. This case underscores the complexity and challenges associated with managing canine Evans syndrome, highlighting the need for further research to optimize treatment regimens and improve prognostic outcomes in affected dogs.

#### ACKNOWLEDGEMENT

We acknowledge the Head, Division of Medicine and Director of IVRI, Izatnagar for providing necessary facilities for disease diagnostics and treatment.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Breitschwert E. Babesiosis. Clinical Microbiology and Infectious Diseases of the Dog and Cat. Philadelphia: WB Saunders. 1984;796:805.
2. Salem NY, Farag HS. Clinical, hematologic, and molecular findings in naturally occurring *Babesia canis vogeli* in Egyptian dogs. Veterinary Medicine International; 2014.
3. Norkus CL, editor. Veterinary technician's manual for small animal emergency and critical care. John Wiley & Sons; 2018.
4. Evans RS, Takahashi K, Duane RT, Payne R, Liu CK. Primary thrombocytopenic purpura and acquired hemolytic anemia: evidence for a common etiology AMA Arch. Intern. Med. 1951; 87(1):48-65.
5. Di Cicco MF, Birkenheuer AJ. Canine Babesiosis - Peer Reviewed. NAVC Clinician's Brief. 2012:31-35.
6. Uilenberg G. Babesia—A historical overview. Vet. Parasitol. 2006;138(1-2):3-10.
7. Koster LS, Lobetti RG, Kelly P. Canine babesiosis: a perspective on clinical complications, biomarkers, and treatment. Vet. Med. Res. 2015;119-28.
8. Wang J, Zhang J, Kelly P, Zheng X, Li M, You J, Huang K, Qiu H, Wang Y, Zhang R, Li J. First description of the pathogenicity of *Babesia vogeli* in experimentally infected dogs. Vet. Parasitol. 2018;253:1-7.
9. Matijatko V, Kiš I, Torti M, Brkljačić M, Kučer N, Rafaj RB, Grden D, Živičnjak T, Mrljak V. Septic shock in canine babesiosis. Vet. Parasitol. 2009;162(3-4): 263-70.
10. Momin M, Aluri A, Reddy S, Pasupala NK. Evans' syndrome-haemolytic anaemia with thrombocytopenia—a rare autoimmune disorder. J. clin. sci. res. 2017;6(4):237-40.
11. Apps C. A case of immune-mediated thrombocytopenia, possibly Evans syndrome, in a neutered Rottweiler dog. Can Vet J. 2023;64(4):379-382.
12. Nandini MK, Poonam Vishwakarma and Ansar Kamran. New therapeutic protocol for canine babesiosis: a case report. J. Dairy Vet Anim. Res. 2016;3(3):112–113

13. Swamy KP, Mohanapriya T, Enbavelan PA, Sundararajan RC, Saravanan S, Ramprabhu R. Triple therapy in Canine Babesiosis-a case report. Int. J. Curr. Microbiol. App. Sci. 2019;8(12): 964-7.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://prh.mbimph.com/review-history/3320>