



Optimizing Nutrient Recommendations for Bottle Gourd Crop by Formulation of Targeted Yield Equations in an *Inceptisol* of Odisha, India

Badireddi Revathi ^a, Subhashis Saren ^a,
Prava Kiran Dash ^{a,b*}, Antaryami Mishra ^a,
Srikanta Kumar Sethy ^{a,c}, Truptimayee Pattnaik ^{a,d}
and Pradip Dey ^e

^a Department of Soil Science, College of Agriculture, Odisha University of Agriculture and Technology, Bhubaneswar-751003, Odisha, India.

^b Regional Research and Technology Transfer Station, Mahisapat, Dhenkanal-759013, Odisha, India.

^c Directorate of Horticulture, Department of Agriculture and Farmers' Empowerment, Government of Odisha, Bhubaneswar-751001, Odisha, India.

^d Department of Basic Science and Humanities, Gandhi Engineering College, Bhubaneswar, Odisha-752054, India.

^e ICAR-Agricultural Technology Application Research Institute, Kolkata-700097, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJSSPN/2024/v10i2279

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115651>

Original Research Article

Received: 10/02/2024
Accepted: 14/04/2024
Published: 22/04/2024

ABSTRACT

A field experiment was conducted to formulate targeted yield equations for bottle gourd under a rice-vegetable cropping system in an *Inceptisol* of Odisha. The experiment started with the creation of three soil fertility gradient stripes by applying no fertilizer, recommended dose and double the

*Corresponding author: Email: dashprava111@gmail.com;

recommended dose of fertilizer in rice (cv. *Lalat*) during the *kharif* season. Each fertility gradient strip was divided into 24 sub-plots and superimposed with 21 different combinations of nutrients containing N, P, and K; FYM in two plots at 5t and 10t ha⁻¹ and one plot was kept as absolute control following which, bottle gourd was grown during the *rabi* season. The highest yield (116.3q ha⁻¹) of bottle gourd was achieved in S-III with an application of 70, 100, and 100 kg N, P₂O₅, and K₂O per hectare, respectively. The nutrient requirement (NR) for producing one quintal of bottle gourd yield was 0.30, 0.20, and 0.30 kg N, P₂O₅ and K₂O, respectively. The effect of graded doses of fertilizers on nutrient requirement, yield and nutrient uptake of bottle gourd were studied and subsequently, fertilizer prescription equations were derived for targeted yield of bottle gourd. A ready reckoner chart has also been prepared for facilitating farmers to achieve the desired yield target of bottle gourd by applying the required quantity of plant nutrients in the existing soil fertility level. The equations provide a basis for site-specific nutrient management based on desired yield targets under varying soil fertility conditions.

Keywords: Bottle gourd; targeted yield; fertilizer prescription equations; inceptisols; STCR-IPNS.

1. INTRODUCTION

Bottle gourd [*Lagenaria siceraria* (Molina) Standl.] is a fast-growing *monoecious* annual climber in the family *Cucurbitaceae*. Bottle gourd fruit is rich in carbohydrates, protein, fat, and minerals including essential elements such as calcium and phosphorus. This nutritionally rich vegetable needs to be cultivated with balanced fertilization so that maximum yield can be achieved with optimum use of externally applied nutrients [1,2]. Moreover, the site-specific application of nutrients has become increasingly important in present-day precision agriculture, which is based on the need-based precise application of agricultural inputs [3-4].

Application of site-specific balanced nutrients on crops can be achieved with fertilizer prescription equations through a targeted yield approach as described by Ramamoorthy et al. [5] The targeted yield approach considers not only the nutrient status of the soil but also the targeted yield for the determination of the amount of nutrients to be added through external sources [6]. Fertilizer application based on the targeted yield equation approach is an effective way to obtain higher yield, net benefit, enhanced nutrient use efficiencies, and fertilizer savings over general fertilizer recommendation [7].

Bhatt et al. [8] observed that the targeted yield model for brinjal was better than the general recommendations in terms of net savings on fertilizers. Recently, Murthy et al. [9] concluded that the targeted yield equation approach was the most suitable and effective approach for fertilizer recommendation for finger millet crop. Eunice et al. [10] while generating targeted yield equations for the amaranthus crop concluded

that fertilizer and manure recommendations based on STCR (Soil Test Crop Response)-targeted yield equations not only improved crop yield but also improved the soil quality. Further, Goyal et al. [11] validated the targeted yield equations for Bt. Cotton in farmers' fields and observed the said approach to be superior to farmer's practice and general recommended dose. Keeping the above facts in view, a field experiment was conducted to formulate targeted yield equations for bottle gourd under a rice-vegetable cropping system in an *Inceptisol* of Odisha.

2. MATERIALS AND METHODS

2.1 The Study Area

The experiment was conducted at E block of the Central Research Farm of OUAT, Bhubaneswar, which comes under the East and Southeastern Coastal Plain agroclimatic zone of Odisha (latitude: 20.266829°, longitude: 85.795856°, elevation: 175 ft above mean sea level). The climate of the study site is hot and humid characterized by hot summer and dry winter with mean annual rainfall, maximum and minimum temperatures of 1,657.8 mm, 33°C, and 22.4°C, respectively.

2.2 Soil Characteristics

Before the development of the fertility gradient, composite soil samples were taken and analyzed for initial soil properties using standard procedures. The experimental site was characterized by medium land, sandy loam in soil texture, acidic (pH 5.67) in soil reaction and medium (5.5 g kg⁻¹) in soil organic carbon

content. The Cation Exchange Capacity of the surface soil was 4.5 cmol (p⁺) kg⁻¹ with 65 percent of base saturation. The experimental site was low (117 kg ha⁻¹) in average soil available nitrogen (N), medium (36 kg ha⁻¹) in average available phosphorus (P), and low (87 kg ha⁻¹) in average available potassium (K). The soil has been classified as fine, mixed, hyperthermic, *Vertic Ustochrepts* as per USDA soil taxonomy.

2.3 Creation of Fertility Gradient

The experiment started with the creation of three soil fertility gradient strips in *Kharif*, 2019. Three strips were created by applying no N, P, K fertilizers in S-I, the recommended dose of fertilizers (N:P: K::80:40:40) in S-II and double of the recommended dose (N: P: K::160:80:80) in S-III strip.

Rice (cv. *Lalat*) crop was grown so that fertilizer could interact with soil constituents, plant rhizosphere and thus become part of the soil system. After the harvest of paddy, grain and straw yields were recorded.

2.4 Test Crop Experiments

After the harvest of the paddy crop, each of the three fertility gradient strips was subdivided into 24 sub-plots. In each strip, out of the 24 sub-plots, 21 sub-plots were superimposed with different graded doses of N, P, and K fertilizers (Table 1) in different combinations; two sub-plots (22nd and 23rd) were applied with FYM at 5 t and 10 t ha⁻¹, respectively and the 24th plot was kept as absolute control. The subplots were created in such a way that all 24 treatments were present in all three strips. Thereby, a total of 72 numbers of subplots were created, and 72 numbers of soil samples from the subplots were collected and analyzed in the laboratory to determine the soil test values. Bottle gourd (cv. *Kaveri*) was grown as the test crop during the following *rabi* season. Sources of nutrients were fertilizer urea, diammonium phosphate and muriate of potash. Nitrogen was applied in split doses i.e., 50% N was applied as basal dose and the rest 50% was applied at the time of flowering. Full doses of P, K and FYM were applied at the time of sowing as basal doses. Standard agricultural practices of bottle gourd were adopted during its cultivation.

2.5 Soil and Plant Analysis

The soil samples collected after the harvest of paddy crop and before planting of bottle gourd

crop were analysed to determine soil organic carbon [12], available nitrogen [13], phosphorus [14], and potassium [15] as outlined by Jackson [16]. After completion of the bottle gourd crop cycle, yield information was recorded, and laboratory analysis of post-harvest soil samples, fruit and vine samples was performed. For analysis of total N, P and K content in fruit and vine, representative plant samples were collected and after proper processing, wet oxidation of samples was done by using the diacid digestion mixture (HNO₃:HClO₄::9:4) for determination of total P and K content. Total N content was determined using the micro-Kjeldahl's method [16]. Uptakes of N, P and K were computed by using yield data and the total N, P and K contents of the plant samples.

2.6 Formulation of Targeted Yield Equations

The factors required for targeted yield equations such as nutrient requirement (NR), soil efficiency (Cs), fertilizer efficiency (Cf), and organic matter efficiency (Co) were calculated following Ramamoorthy's inductive cum targeted yield model as given below.

$$NR \text{ (kg/ q)} = \text{Uptake of nutrient by bottle gourd (kg/ ha)} / \text{Yield of bottle gourd (q/ ha)}$$

$$Cs \text{ (\%)} = \text{Uptake of nutrient in absolute control plot (kg/ha)} / \text{Initial soil test value of a particular nutrient in control plot (kg/ha)} * 100$$

$$Cf \text{ (\%)} = \frac{\text{Uptake of nutrient in fertilizer treated plot (kg/ ha)} - [\text{Initial soil test value} \times Cs/100]}{\text{Nutrient applied through fertilizer (kg/ha)}} * 100$$

$$Co \text{ (\%)} = \frac{\text{Uptake of nutrient in organic matter treated plot} - (\text{Initial soil test value of control plot with FYM} \times Cs/100)}{\text{Nutrient applied through organic matter (kg/ha)}}$$

All the required parameters were calculated and combined for formulating the targeted yield equations with and without FYM as follows:

$$FD \text{ (without FYM)} = NR \times 100 \times T / Cf - Cs \times STV / Cf$$

$$FD \text{ (with FYM)} = \frac{NR \times 100 \times T}{Cf - Cs \times STV / Cf - Coxnutrient \text{ (kg/t) in FYM} \times FYM \text{ (t/ha)}} / Cf$$

Where, FD = fertilizer dose (kg ha⁻¹), T = targeted yield (q ha⁻¹), and STV = soil test value.

Table 1. Combinations of nitrogen, phosphorus, potassium and FYM used in the experiment

T ₁ - N ₀ P ₂ K ₂	T ₁₃ - N ₂ P ₃ K ₂
T ₂ - N ₁ P ₁ K ₁	T ₁₄ - N ₂ P ₃ K ₃
T ₃ - N ₁ P ₁ K ₂	T ₁₅ - N ₁ P ₃ K ₂
T ₄ - N ₁ P ₂ K ₁	T ₁₆ - N ₃ P ₂ K ₁
T ₅ - N ₁ P ₂ K ₂	T ₁₇ - N ₃ P ₂ K ₂
T ₆ - N ₂ P ₀ K ₂	T ₁₈ - N ₃ P ₂ K ₃
T ₇ - N ₂ P ₁ K ₁	T ₁₉ - N ₂ P ₁ K ₃
T ₈ - N ₂ P ₁ K ₂	T ₂₀ - N ₃ P ₃ K ₂
T ₉ - N ₂ P ₂ K ₀	T ₂₁ - N ₃ P ₃ K ₃
T ₁₀ - N ₂ P ₂ K ₁	T ₂₂ - N ₀ P ₀ K ₀ + FYM @ 5 t ha ⁻¹
T ₁₁ - N ₂ P ₂ K ₂	T ₂₃ - N ₀ P ₀ K ₀ + FYM @ 10 t ha ⁻¹
T ₁₂ - N ₂ P ₂ K ₃	T ₂₄ - N ₀ P ₀ K ₀ (absolute control)

(where, N₀, N₁, N₂, N₃ corresponds to 0, 30, 50, 70 kg N/ha; P₀, P₁, P₂, P₃ corresponds to 0, 60, 80, 100 kg P₂O₅/ha; K₀, K₁, K₂, K₃ corresponds to 0, 60, 80, 100 kg K₂O /ha)

3. RESULTS AND DISCUSSIONS

3.1 Effect on Yield of Paddy Crop

Three fertility gradient stripes were created while cultivating rice crop by applying no fertilizer in the S-I block (N₀: P₀: K₀), the recommended dose of fertilizer in S-II (N₈₀: P₄₀: K₄₀) and double of the recommended dose of fertilizer in S-III block (N₁₆₀: P₈₀: K₈₀). Data showed that the highest grain yield (45.2 q ha⁻¹) of rice was achieved in S-II block followed by S-III (42.9 q ha⁻¹) and S-I (29.6 q ha⁻¹) blocks (Table 2). On the other hand, the highest straw yield was recorded (53.4 q ha⁻¹) in the S-III block followed by the S-II (51.7 q ha⁻¹) and S-I (33.8 q ha⁻¹) blocks. Both grain and straw yield of rice were lowest in S-I (control). Data clearly indicated that the application of a recommended dose of fertilizers (S-II) resulted in 52% more grain yield and 53% more straw yield compared to the control strip (N₀P₀K₀). However, the S-III strip showed 44% more grain yield and 57% more straw yield than the control strip. The results clearly showed that higher doses of fertilizer in S-III increased the vegetative growth of rice which was reflected in straw yield. Excess use of fertilizers often delays the maturity of the crop [17,18].

3.2 Effect on Soil Fertility Status

The experimental result showed that the highest soil nutrient was built up in the S-III strip among

the three fertility gradient stripes. The mean values of soil available NPK increased with an increase in fertilizer doses from the S-I to S-III strip. The mean available soil N was found to be 117.3, 135.9 and 145.6 kg ha⁻¹, that of P₂O₅ was 34.2, 41.8 and 64.5 kg ha⁻¹ and the mean available K₂O was 87.1, 66.8 and 99.3 kg ha⁻¹ in S-I, S-II, and S-III stripes, respectively (Table 3). Higher soil fertility status was observed in the S-III strip as the highest quantities of fertilizers were applied for rice during *kharif* and a large amount of applied nutrients might have remained unutilized after the harvest of the crop [19,20]. Similar results have recently been reported by Pandey et al., [21] while formulating targeted yield equations for Sesame.

3.3 Effect on Fruit Yield

Results indicated that the fruit yield of bottle gourd in S-I, S-II and S-III was found to vary from 52.9 to 92.5, 73.6 to 106.3 and 95.1 to 116.3 q ha⁻¹ with the average yield of 77.5, 87.3 and 103.5 q ha⁻¹, respectively (Table 3). The lowest yield of bottle gourd was observed in the absolute control plot in all three fertility gradient stripes as it did not receive any fertilizer. The highest fruit yield was achieved in the S-III strip with a fertilizer dose of 70:100:100 kg N: P₂O₅: K₂O and this graded dose of fertilizer recorded the highest yield in all the stripes (supplementary material). In contrast, the lowest yield was found in the S-I as no fertilizer was applied to rice during *kharif*. The percentage increase of bottle gourd fruit yield in S-II and S-III as compared to S-I was observed to be 12% and 33%, respectively. Similar observations have also been reported for French bean [22], green gram [23], coriander [24], and rice [25]. Thus, higher nutrient content in soil resulted in higher yield of bottle gourd.

3.4 Effect on Biomass (vine) Yield

The biomass (leaves and vine) yield in S-I, S-II and S-III was found to range between 32.8 to 55.8, 41.0 to 60.2 and 53.7 to 68.4 q ha⁻¹ with average values of 44.7, 49.9 and 58.9 q ha⁻¹, respectively (Table 3; Fig.1). The highest biomass production also followed the same trend of fruit yield i.e., biomass yield also increased from S-I to S-III. Similar observations were also recorded in different crops by the earlier workers [17,18].

Table 2. Effect of graded doses of fertilizer on rice

Strip-I N ₀ P ₀ K ₀		Strip-II N ₈₀ P ₄₀ K ₄₀		Strip-III N ₁₆₀ P ₈₀ K ₈₀	
	Yield (q ha ⁻¹)		Yield (q ha ⁻¹)		Yield (q ha ⁻¹)
Grain	29.6	Grain	45.2	Grain	42.9
Straw	33.8	Straw	51.7	Straw	53.4
			% increase		% increase
			52		44
			53		57

Table 3. Range and average yield of fruit and biomass of bottle Gourd (cv. Kaveri), soil test values and NPK uptake in different fertility gradient stripes

Parameter	Strip-I		Strip -II		Strip-III	
	Range	Mean ±SEm	Range	Mean ±SEm	Range	Mean ±SEm
Fruit yield (q ha ⁻¹)	52.9-92.9	77.5±1.38	73.6-106.3	87.3±1.35	95.1-116.3	103.5±0.61
Vine yield (q ha ⁻¹)	32.8-55.8	44.7±1.11	41.0-60.2	49.9±1.0	53.7-68.4	58.9±0.51
Av. N (kg ha ⁻¹)	106.7-128.6	117.3±0.48	123.8-143.8	135.9±0.42	137.9-152.3	145.6±0.36
Av. P ₂ O ₅ (kg ha ⁻¹)	27.6-38.8	34.2±0.62	31.9-49.1	41.8±0.93	45.3-75.9	64.5±1.23
Av. K ₂ O (kg ha ⁻¹)	66.1-126.7	87.1±1.75	42.6-102.8	66.8±1.72	65.2-195.6	99.3±1.13
N uptake (kg ha ⁻¹)	12.2-43.8	26.6±1.61	20.2-47.6	30.8±1.81	27.1-61.0	42.1±1.61
P uptake (kg ha ⁻¹)	5.0-26.9	13.8±1.93	10.5-34.8	19.4±1.85	16.2-45.0	30.3±1.79
K uptake (kg ha ⁻¹)	9.6-42.4	24.5±1.63	15.1-53.5	32.6±1.96	25.1-68.4	44.3±1.60

Table 4. Basic data required for formulating fertilizer adjustment equations for bottle gourd crop

Basic data	N	P ₂ O ₅	K ₂ O
Nutrient requirement (kg q ⁻¹)	0.3	0.2	0.3
Soil efficiency (Cs, %)	15	21	21
Fertilizer efficiency (Cf, %)	28	15	22
Organic matter efficiency (Co, %)	7.6	9.8	7.8

Table 5. Targeted yield equations for bottle gourd

Fertilizer dose	Without FYM	With FYM (IPNS)
N (kg ha ⁻¹)	FN= 1.26T – 0.54 SN	FN= 1.26T – 0.54 SN – 0.26 O N
P ₂ O ₅ (kg ha ⁻¹)	FP ₂ O ₅ = 1.6T- 1.34 S P ₂ O ₅	FP ₂ O ₅ = 1.6T- 1.34 S P ₂ O ₅ – 0.62 O P ₂ O ₅
K ₂ O (kg ha ⁻¹)	FK ₂ O= 1.6T- 0.94 S K ₂ O	FK ₂ O= 1.6T- 0.94 S K ₂ O – 0.34 O K ₂ O

(FN, FP₂O₅ and FK₂O= Nitrogen, phosphatic and potassic fertilizers required (kg ha⁻¹); T= Yield target (in quintals); SN, S P₂O₅ and S K₂O = Soil testing values of N, P₂O₅ and K₂O (in kg ha⁻¹); ON, OP₂O₅ and OK₂O = Nutrients N, P₂O₅ and K₂O supplied through organic source (in kg ha⁻¹))

Table 6. Ready reckoner chart of fertilizer doses for different yield targets of bottle gourd under varying fertility status

Initial soil status (kg ha ⁻¹)			Yield Target (q ha ⁻¹)											
			80			100			120			140		
SN	SP ₂ O ₅	SK ₂ O	FN	FP ₂ O ₅	FK ₂ O	FN	FP ₂ O ₅	FK ₂ O	FN	FP ₂ O ₅	FK ₂ O	FN	FP ₂ O ₅	FK ₂ O
100	20	70	47	101	62	72	133	94	97	165	126	122	197	158
110	25	80	41	94	53	66	126	85	92	158	117	117	190	149
120	30	90	36	88	43	61	120	75	86	152	107	112	184	139
130	35	100	31	81	34	56	113	66	81	145	98	106	177	130
140	40	110	25	74	20	50	106	57	76	138	89	101	186	121
150	45	120	20	68	15	12.5	100	20	70	132	79	95	163	111
160	50	130	14	61	20	40	93	20	65	125	70	90	157	102
170	55	140	9	54	20	34	86	28	59	118	60	85	150	92
180	60	150	4	48	20	29	80	19	54	112	51	79	144	79
190	65	160	12.5	41	20	23	73	10	49	105	42	74	137	74

(FN, FP₂O₅ and FK₂O= Nitrogen, phosphatic and potassic fertilizers required (kg ha⁻¹); SN, S P₂O₅ and S K₂O = Soil testing values of N, P₂O₅ and K₂O (in kg))




Fig.1. Average fruit and vine yields of bottle gourd crop in different fertility gradient strips

3.5 Effect on Nutrient Uptake

The uptake of N, P and K showed an increasing trend with an increase in artificially created fertility gradient stripes from S-I to S-III. The mean uptake of nitrogen was 26.6, 30.8 and 42.1 kg ha⁻¹, that of phosphorus was 13.8, 19.4 and 30.3 kg ha⁻¹ and mean potassium uptake was 24.5, 32.6 and 44.3 kg ha⁻¹ in S-I, S-II, and S-III stripes, respectively (Table 3). Thus, the uptake of major nutrients increased with the applied nutrients and nutrients present in the soil. Higher plant nutrient uptake corresponded to higher yields of both fruit and biomass. The uptake of N, P and K was found to be highest in the fertilizer dose of 70:100:100 kg N: P₂O₅: K₂O ha⁻¹ in all fertility gradient stripes (supplementary material). Among the 21 fertilizer-treated plots, one of each of the nutrients i.e., N, P and K was not applied (supplementary material) in some of the subplots (Table 1). Both nutrient uptake and yield drastically reduced where a single nutrient was not applied (supplementary material). Such type of treatment combinations in the present investigation showed the importance of any particular macronutrient on the yield and uptake of the crop.

3.6 Formulation of Targeted Yield Equation and Ready Reckoner for Bottle Gourd

The nutrient requirement (NR) for producing one quintal of bottle gourd was 0.3 kg N, 0.2 kg P₂O₅ and 0.3 kg K₂O. Soil efficiency (Cs) was found to

be 15, 21 and 21 per cent; fertilizer efficiency (Cf) was 28, 15 and 22 per cent and organic matter efficiency (Co) was 7.6, 9.8 and 7.8 per cent for N, P₂O₅ and K₂O, respectively (Table 4). Such low efficiencies of nutrients in both soil and fertilizer sources could be because of persistent high levels of soil acidity prevailed in the study area since soil acidity is associated with poor nutrient availability and poor nutrient use efficiencies [26]. Soil acidity has been reported in several other studies in the state of Odisha [27-31].

Targeted yield equations for bottle gourd (*cv. Kaveri*) were thus formulated (Table 5). In the equations, the yield target (T) was fixed based on the yield potential of the crop. In the equations, SN, SP₂O₅ and SK₂O values stand for available soil nitrogen, soil phosphorus and soil potassium, respectively. Similarly, ON, OP₂O₅ and OK₂O values stand for nitrogen, phosphorus, and potassium through organic sources. The amount of fertilizer to be added is dependent on the status of soil nutrients and the yield target. The equations show that if a part of the nutrient requirement is filled up by the application of FYM, there will be a net saving on the cost of fertilizers. Similar findings were also reported by Singh et al. [32-34]. A ready reckoner for fertilizer doses was prepared to facilitate farmers to achieve the desired yield target of bottle gourd by applying the required quantities of plant nutrients in different existing soil fertility levels (Table 6). A higher yield target of the crop suggested more fertilizer requirements. On the other hand, lower

nutrient content in the soil needed higher fertilizer doses to achieve the desired yield target. In case, when soil nutrient content is high and the yield target is low, the required fertilizer as per the equations will be very low or in some cases, values might be negative. In that case, only a maintenance dose of 25% of the recommended dose of fertilizer is recommended to avoid nutrient mining from the soil. Thus, in soils with higher initial nutrient content, unnecessary application of fertilizers can be checked by using the targeted yield equation approach.

4. CONCLUSION

Fertilizer recommendation based on targeted yield equations is a scientific way of precise and need-based nutrient management. This study has led a path towards adopting site-specific nutrient management in bottle gourd crop. This will not only supply the required quantity of nutrients to achieve a specific yield target but also will help in maintaining soil fertility status over the long run. For further validation, in future, the equations need to be tested at farmers' fields at multiple locations with varying levels of soil fertility status and yield targets. For effective adaptation of these equations among the farmers, initial soil testing must be promoted as a first step. The targeted yield equations prepared in this study will especially be useful to increase the production and productivity of bottle gourd crop in red, laterite and yellow soils of Odisha (*Inceptisols* and *Alfisols*) and similar soils at other places. Integrating the application of FYM along with inorganic sources of nutrients will help curtail the amount of inorganic nutrients to be applied. Thus, the targeted yield equation approach can be effectively utilized for improving bottle gourd yield and saving costs on fertilizers, besides having environmental benefits by avoiding the over-application of chemical fertilizers.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the Indian Council of Agricultural Research (ICAR) for the support through AICRP on Soil Test Crop Response (STCR) project. The physical facility provided by the Odisha University of Agriculture and Technology, Bhubaneswar, Odisha is also gratefully acknowledged.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lokya T, Mishra A., Saren S, Dash PK. GPS and GIS based soil fertility maps of KVK farm, Nayagarh located in the North-Eastern Ghat Agro-Climatic zone of Odisha, India. *Journal of Pharmacognosy and Phytochemistry*. 2020;9:599–605.
2. Kishore MA, Mishra A, Dash PK, Saren S, Das TK, Priyadarsini S, Sethy H, Sardar MS. Vertical distribution of available nutrients in boudh block of boudh District Catena. *International Journal of Plant & Soil Science*. 2022;34:1681–1686.
3. Dash PK, Panigrahi N, Mishra A. Identifying opportunities to improve digital soil mapping in India: A systematic review. *Geoderma Regional*. 2022a;e00478.
4. Dash PK. Chapter 22 - Remote sensing as a potential tool for advancing digital soil mapping, in: Dharumarajan, S., Kaliraj, S., Adhikari K, Lalitha M, Kumar N. (Eds.), *Remote Sensing of Soils*. Elsevier. 2024;357–370. Available: <https://doi.org/10.1016/B978-0-443-18773-5.00021-1>
5. Ramamoorthy B, Narasimhan RL, Dinesh RS. Fertilizer application for specific yield targets on Sonora 64 (wheat). *Indian Farming*. 1967;17(5):43-45.
6. Ranjan KR, Singh Y, Singh S, Dey P. Fertilizer recommendations developed through soil test crop response studies with integrated plant nutrient management system for field pea in an Inceptisol. *Journal of the Indian Society of Soil Science*. 2018;66:318–323.
7. Pande J, Singh S. Fertilizer recommendations based on targeted yield concept for cabbage grown in a Mollisol of Uttarakhand. *Journal of the Indian Society of Soil Science*. 2016;64 (3):265-270.
8. Bhatt P, Singh S, Pant PK., Chandra R, Gautam P, Bhatt L, Arya A. Assessment of nutrient requirement for spring brinjal (*Solanum melongena* L.) through soil test crop response approach on north Indian Mollisol. *Journal of the Indian Society of Soil Science*. 2021;69:162–170.
9. Murthy RK, Bhavya N, Govinda K., Kumar SU, Basavaraja P. Validation of Targeted yield equations for finger millet under southern dry zone of Karnataka. *Int. J. Plant Soil Sci*. 2023;35:84–91.
10. Eunice D, OP RR, Gopinath PP. Evaluation of STCR Based targeted yield equations of amaranthus (*Amaranthus*

- tricolor* L.) in southern laterite soils (AEU-8) of Kerala. International Journal of Plant & Soil Science. 2023;35:53–58.
11. Goyal V, Bhardwaj K, Dey P. Refinement of fertilizer prescriptions equations of Bt. cotton hybrids under integrated plant nutrient supply in semi-arid inceptisols of Haryana. Journal of the Indian Society of Soil Science. 2023;71:87–98.
 12. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. Soil Science. 1934;37(1):29-38.
 13. Subbiah BV, Asija GL. A rapid procedure for the determination of available nitrogen soils. Current Science. 1956;25:259-260.
 14. Bray RH, Kurtz LT. Determination of total organic and available forms of phosphorus in soils. Soil Science. 1945;59:39-45.
 15. Hanway JJ, Heidel H. Soil analyses methods as used in Iowa State college soil testing laboratory. Iowa Agric. 1952;57:1-31.
 16. Jackson ML. Soil chemical analysis. Prentice Hall of India (Pvt.) Ltd., New Delhi; 1973.
 17. Gogoi A, Mishra A, Jena B. Soil test-based fertilizer recommendation for targeted yield of pumpkin (*Cucurbita moschata*) under rice-pumpkin cropping system in an Inceptisol of Odisha. Environment and Ecology. 2011;29(2): 574-576.
 18. Mishra A, Saren S, Dey P. Integrated nutrient management and formulation of targeted yield equations for Cowpea [*Vigna unguiculata* (L.) Walp.]. Environment and Ecology 2016;35(2A):867-872.
 19. Ammal UB, Venkatakrishnan L, Vetrivel M, Sankar R, Coumaravel K, Dey P. Soil test based fertilizer prescriptions under IPNS for rice-rice cropping sequence in Inceptisol of Puducherry. Journal of the Indian Society of Soil Science 2020;68: 408–414.
 20. Singh J, Mavil M, Saini SP. Fertilizer prescription for target yield of wheat (*Triticum aestivum*) in alluvial soils. Indian Journal of Agricultural Sciences. 2022;92: 1502–1505.
 21. Pandey R, Trivedi A, Sharma V, Chobhe K, Dey P, Chandra S. Soil test-based fertilizer prescription for targeted yield of sesame (*Sesamum Indicum* L.). Communications in Soil Science and Plant Analysis 2023;54:2572–2578.
 22. Pogula S, Truptimayee Mishra A, Saren S, Dey P. Soil test based fertilizer recommendation for targeted yield of test-basedan (*Phaseolus vulgaris*) under Rice-French bean cropping system. International Journal of Bio-resource and Stress Management. 2016;7(5):1128-1130.
 23. Saren S, Mishra A, Dey P. Site specific nutrient management through targeted yield equations formulated for green gram (*Vigna radiata* L. Wilczek). Legume Research. 2018;41(3):436-440.
 24. Singh YV, Singh SK, Meena RN, Jatav HS.. Soil test crop response approach for optimizing integrated plant nutrient management supply for coriander in an Inceptisol of eastern Uttar Pradesh. Journal of the Indian Society of Soil Science. 2020;68(1):114-119.
 25. Mondal S, Gorai D, Saha N, Dasgupta S, Bhattacharyya K, Dey P. Integrated fertilizer prescriptions through targeted yield model and prediction of post-harvest soil test values for kharif rice (*Oryza sativa*). Journal of the Indian Society of Soil Science. 2020;68 (3):330-336.
 26. Digal M, Saren S, Mishra A, Dash PK, Swain N, Acharya BP.. Soil fertility status of some villages in Phiringia block of Kandhamal district under North-Eastern Ghat agro climatic zone of Odisha, India. Journal of Pharmacognosy and Phytochemistry. 2018;7:658–662.
 27. Dash PK, Mishra A, Saren S. Characterization and taxonomic classification of soils under a toposequence located in Eastern India. Environment and Ecology. 2019;37(4): 1240—1249.
 28. Sathy SK, Mishra A, Dash PK, Saren S, Dey P. Geo-Information based soil fertility Status of Deogarh District of Odisha, India. Int. J. Curr. Microbiol. App. Sci. 2019;8: 255–262.
 29. Singh RP, Mishra A, Dash PK, Saren S, Mallik H, Mishra BB. Soil Fertility Assessment of Some Villages in Kankadahad Block of Odisha. Environment and Ecology. 2021;39:250–255.
 30. Dash PK, Mishra A, Patro H, Nayak RK, Jena B, Satapathy M.. Soil characterization along a toposequence located in North-Central Plateau Agro-climatic zone of Odisha. Ecology Environment and Conservation. 2022b;28:471—476.

31. Pattanaik T, Dash PK, Mishra A, Saren S, Phonglosa A, Kishore MA. Vertical distribution of available plant nutrients in soils of mid-central tableland agro-climatic zone of Odisha, India, Environment and Ecology. 2023;41(4D): 2982—2988.
32. Singh Y, Parveen A, Singh S, Dey P. Soil test-based fertilizer prescriptions under integrated plant nutrient management system for barley in an Inceptisol of eastern plain zone of Uttar Pradesh. Journal of the Indian Society of Soil Science. 2017;65:423–427.
33. Gogoi Jhinkwan S, Singh S, Pareek N, Gautam P. Optimization of integrated nutrient recommendations through soil test crop response approach for forage oats (*Avena sativa* L.) in a mollisol. Journal of the Indian Society of Soil Science. 2021; 69:156–161.
34. Mishra A, Dash BB, Nanda SK, Das D, Dey P. Soil test based fertilizer recommendation for targeted yield of tomato (*Lycopersicon esculentum*) under rice-tomato cropping system in an ustochrept of Odisha. Environment and Ecology. 2013;31(2A): 655-658.

SUPPLEMENTARY MATERIALS

Table 1. Initial soil test values, fertilizer dose, yield and uptake of nutrients in Bottle Gourd in Strip-I

Sl. No.	Initial soil test value (kg ha ⁻¹)				Fertilizer dose (kg ha ⁻¹)			Yield (q ha ⁻¹)		Uptake of nutrients (kg ha ⁻¹)		
	OC (%)	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	Fruit	Biomass	N	P	K
1	0.68	119.4	33.5	85.1	0	80	80	63.1	35.8	14.8	8.7	12.8
2	0.66	123.1	35.3	86.5	30	60	60	68.9	38.2	18.4	8.9	15.1
3	0.65	121.7	37.8	79.0	30	60	80	75.3	43.7	21.0	11.6	18.1
4	0.58	128.6	35.6	92.8	30	80	60	80.7	44.8	22.1	12.6	18.8
5	0.61	120.8	35.5	80.2	30	80	80	82.6	46.8	23.7	13.4	20.8
6	0.64	122.3	34.2	74.0	50	0	80	63.4	36.7	19.7	7.4	14.5
7	0.68	121.8	31.8	126.7	50	60	60	71.2	41.8	22.7	10.1	18.2
8	0.69	125.9	35.2	118.6	50	60	80	74.9	43.5	22.8	11.2	20.6
9	0.71	123.2	32.3	99.0	50	80	0	65.3	36.2	19.6	9.8	12.4
10	0.66	115.4	34.8	96.5	50	80	60	78.6	45.2	26.1	13.7	22.2
11	0.67	119.6	38.7	99.3	50	80	80	84.7	51.3	29.3	16.1	28.2
12	0.55	108.9	36.4	100.3	50	80	100	85.3	50.9	30.3	17.0	32.6
13	0.52	116.2	35.2	89.0	50	100	80	88.8	52.1	30.4	19.1	31.3
14	0.53	114.6	40.1	79.0	50	100	100	87.3	49.5	30.7	19.4	33.6
15	0.42	113.8	37.7	72.7	30	100	80	85.5	47.1	28.4	19.6	30.9
16	0.52	112.4	38.1	73.9	70	80	60	88.9	50.7	36.9	19.7	32.2
17	0.54	110.3	37.2	67.7	70	80	80	90.3	51.8	38.7	20.5	35.7
18	0.50	116.2	36.8	79.0	70	80	100	90.1	52.8	39.8	21.6	40.4
19	0.47	118.4	39.4	116.6	50	60	100	77.0	45.2	28.5	17.4	32.9
20	0.54	114.1	40.7	68.9	70	100	80	89.7	52.1	41.2	11.4	39.2
21	0.55	113.8	37.8	75.2	70	100	100	92.9	55.8	43.8	26.9	42.4
22	0.58	117.2	36.9	84.0	0	0	0	61.5	36.3	14.7	5.9	11.9
23	0.51	106.7	34.2	91.5	0	0	0	58.2	33.8	14.6	6.24	12.0
24	0.49	112.2	37.1	55.1	0	0	0	55.9	32.8	12.2	5.0	9.6

Table 2. Fertilizer dose, initial soil test values, yield of fruit and biomass and uptake of nutrients in Bottle Gourd in Strip-II

Sl. No.	Initial soil test value (kg ha ⁻¹)				Fertilizer dose (kg ha ⁻¹)			Yield (q ha ⁻¹)		Uptake of nutrients (kg ha ⁻¹)		
	OC (%)	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	Fruit	Biomass	N	P	K
1	0.57	140.7	43.8	61.4	0	80	80	75.7	44.7	20.9	10.8	17.1
2	0.42	136.2	44.3	67.7	30	60	60	76.1	45.1	23.1	10.5	18.6
3	0.49	129.8	48.4	89.0	30	60	80	78.5	44.1	24.4	12.2	20.1
4	0.53	134.3	46.2	91.5	30	80	60	81.3	45.7	25.3	14.5	22.7
5	0.54	128.4	45.0	102.8	30	80	80	78.0	43.4	26	12.8	25.2
6	0.51	131.2	44.8	79.0	50	0	80	77.4	43.9	22.9	12.7	20.3
7	0.58	139.1	42.2	67.7	50	60	60	80.2	46.8	27.2	15.7	24.1
8	0.48	137.9	45.7	65.2	50	60	80	79.8	45.4	25.9	16.1	27.5
9	0.42	135.4	41.9	60.2	50	80	0	79.1	45.5	22.4	13.5	24.3
10	0.47	141.2	42.4	60.2	50	80	60	79.6	46.4	27.5	20.2	30.5
11	0.50	137.8	47.9	55.1	50	80	80	80.5	46.2	28.7	20	33.6
12	0.46	142.1	45.8	60.2	50	80	100	93.6	52.7	34.3	25.7	44.1
13	0.55	135.6	44.1	79.0	50	100	80	98.5	56.2	37.1	27	42.4
14	0.49	143.8	48.0	85.2	50	100	100	101.4	58.6	39.7	28.8	48.8
15	0.46	132.9	44.9	62.7	30	100	80	100.1	57.1	36.5	26.4	43
16	0.42	140.6	44.7	56.4	70	80	60	102.0	58.4	38.8	24.9	44.2
17	0.47	137.8	46.0	42.6	70	80	80	100.9	57.5	40.5	24.4	46.3
18	0.43	134.5	45.9	56.4	70	80	100	101.3	57.2	43.1	24.3	49.8
19	0.49	135.2	48.3	70.2	50	60	100	101.0	58.1	40.8	24.3	48.6
20	0.52	142.1	49.0	51.4	70	100	80	102.0	58.4	44.5	32.2	50.1
21	0.56	137.9	47.6	60.2	70	100	100	106.3	60.2	47.6	34.8	53.5
22	0.52	135.0	44.2	55.1	0	0	0	73.6	42.5	21.5	11.6	16
23	0.54	129.7	39.8	61.4	0	0	0	74.1	41	22.5	12.1	16.9
24	0.56	123.8	40.9	63.9	0	0	0	75.6	42.8	20.2	11.1	15.1

Table 3. Fertilizer dose, initial soil test values, yield of fruit and biomass and uptake of nutrients in Bottle Gourd in Strip-III

Sl. No.	Initial soil test value (kg ha ⁻¹)				Fertilizer dose (kg ha ⁻¹)			Yield (q ha ⁻¹)		Uptake of nutrients (kg ha ⁻¹)		
	OC (%)	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	Fruit	Biomass	N	P	K
1	0.68	142.6	71.6	126.7	0	80	80	96.6	53.7	30.5	19.5	32.3
2	0.64	148.2	78.8	140.4	30	60	60	99.2	57.8	35.4	20.7	36.7
3	0.58	139.4	75.3	80.2	30	60	80	101.0	58.4	36.7	22.6	38
4	0.66	149.5	71.9	68.9	30	80	60	102.1	59.2	36.7	24	39.7
5	0.68	138.9	72.5	105.3	30	80	80	104.9	58.9	39.5	25.6	43
6	0.59	142.4	74.6	100.3	50	0	80	97.2	55.1	32.6	34.5	31.3
7	0.56	148.5	75.1	115.4	50	60	60	100.9	57.8	37	31	35.6
8	0.64	150.1	74.3	95.3	50	60	80	101.2	58.1	38.7	30.7	40.7
9	0.65	145.8	70.0	82.7	50	80	0	95.1	54.4	32	24.3	31.1
10	0.67	149.0	73.1	79.0	50	80	60	98.6	55.7	39.1	30.2	37.9
11	0.69	145.0	75.6	195.6	50	80	80	101.9	57.2	42.5	32.3	43.4
12	0.61	152.3	74.7	115.4	50	80	100	104.4	55.7	44.1	33.3	51.5
13	0.56	145.8	72.5	70.2	50	100	80	105.6	59.7	46.9	36.1	48.1
14	0.63	151.2	77.3	81.5	50	100	100	110.0	63.4	50.6	39.9	59.5
15	0.57	141.8	73.1	111.6	30	100	80	109.9	62.4	50.9	38.2	54.6
16	0.61	150.7	76.4	74.0	70	80	60	108.0	61.2	51	34.9	51.8
17	0.54	147.6	77.0	70.2	70	80	80	109.9	62.8	50.1	34.6	54
18	0.52	143.8	75.3	126.6	70	80	100	110.4	62.5	54	39.9	62.1
19	0.57	145.0	74.8	65.2	50	60	100	109.0	61.8	54.3	33.1	57.4
20	0.60	150.2	78.2	77.7	70	100	80	113.3	65.8	57.5	42.3	63.6
21	0.51	149.3	79.7	127.9	70	100	100	116.3	68.4	61	45	68.4
22	0.53	141.2	72.4	84.0	0	0	0	96.6	54.7	30.7	18.5	27.2
23	0.46	138.7	67.1	84.0	0	0	0	98.5	55.7	33.6	19.8	31.5
24	0.43	137.9	69.8	106.6	0	0	0	95.5	54.4	27.1	16.2	25.1

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/115651>