



Effect of Integrated Nutrient Management on Soil Health and Nutrient Balance Sheet of Soybean under Vertisols of Rajasthan

Neha Meena ^{a*}, M. K. Sharma ^a, Rajendra Kumar Yadav ^a, D. S. Meena ^a,
Seema Choudhary ^b, Kamlesh Bhil ^a, Rajesh Meena ^a, Suman Dhayal ^a,
Mamta Choudhary ^c and Narendra Danga ^a

^a Department of Soil Science and Agricultural Chemistry, College of Agriculture, Ummadganj, AU- Kota, Rajasthan-324 001, India.

^b Department of Soil Science and Agricultural Chemistry, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior, MP-474002, India.

^c Department of Soil Science and Agricultural Chemistry, Rani Lakshmi Bai Central Agricultural University, Jhansi, UP-284003, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i1831080

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87084>

Original Research Article

Received 05 March 2022

Accepted 09 May 2022

Published 13 May 2022

ABSTRACT

Background: A field experiment was conducted during kharif, 2019 at ARS farm, College of Agriculture, Ummadganj, Kota (Rajasthan). Effect of Integrated Nutrient Management on Soil Health and Nutrient Balance Sheet of soybean Under Vertisols of Rajasthan. The soil of the experimental field is clay loam having low to medium fertility status and good porosity.

Method: The experiment consisted of 10 treatments viz. T₀ - Control (Absolute), T₁ - RDF (20-40-40), T₂ - 75% RDF + FYM (2.0 t ha⁻¹), T₃ - 50% RDF + FYM (4.0 t ha⁻¹), T₄ - 75% RDF + FYM (2.0 t ha⁻¹) + *Rhizobium*, T₅ - 50% RDF + FYM (4.0 t ha⁻¹) + *Rhizobium*, T₆ - 75% RDF + Vermicompost (1.0 t ha⁻¹), T₇ - 50% RDF + Vermicompost (2.0 t ha⁻¹), T₈ - 75% RDF + Vermicompost (1.0 t ha⁻¹) + *Rhizobium*, T₉ - 50% RDF + Vermicompost (2.0 t ha⁻¹) + *Rhizobium*. The experiment was laid out in randomised block design and was replicated thrice.

Result: Among the nutrient management treatment, application of 50% RDF + Vermicompost (2.0 t ha⁻¹) + *Rhizobium* (T₉) significantly built up of the soil available nitrogen (258 kg ha⁻¹), available phosphorus (21.67 kg ha⁻¹) and available potassium (445.35 kg ha⁻¹), micronutrient and organic carbon (0.59) was recorded.

*Corresponding author: E-mail: neha1996jaipur@gmail.com;

Keywords: *Rhizobium japonicum*; FYM; vermicompost; micronutrients; soybean.

1. INTRODUCTION

Soybean (*Glycine max* L.) belongs to family *Leguminaceae* or *Fabaceae* and Sub- family *Papilionaceae*. Soybean [*Glycine max* (L.)] is predominantly cultivated under rainfed condition during kharif season at Hadoti region of Rajasthan. Soybean is one of the major kharif oilseed crops in India, mainly in semi-arid tropics of central India comprising the states of Madhya Pradesh, Maharashtra and Rajasthan. Estimated production of soybean in world is 364.33 million tonnes production and area is 127.19 million hectares whereas in India it was 12.10 million tonnes production from an area of 10.80 million hectares in Rajasthan is 1.16 million tonnes production and area is 0.93 million hectares [1]. Consecutive use of organic manures, biofertilizers and inorganic fertilizers to achieve sustained crop production and maintain the soil health is the basic need of the hour. Use of organic manures alone or in combination with chemical fertilizers will help to improve physico-chemical properties of the soils. Organic manures are used to supply both macro and micronutrients and sustain amount of humic substances particularly humic and fulvic acid that helps to maintain soil reaction. Organic sources of nutrients are derived from animal, human and agriculture wastes (e.g. composting and crop residues). Vermicompost is rich organic manure consist of macro and micronutrients, plant growth promoting substances, beneficial micro-organisms that are necessary for plant growth [2]. Seed inoculation with effective *Rhizobium* inoculant is recommended to ensure additional nodulation and N₂ fixation for maximum growth and yield of soybean crop. The organic manures along with bio fertilizers help in reducing the dose of inorganic fertilizer, which in turn reduces the cost of cultivation and help in improving the soil health. Therefore, this study was undertaken to analyze the effect of combined use of organic and inorganic sources of nutrients on growth and yield of soybean in the *Vertisols* of Rajasthan.

2. MATERIALS AND METHODS

The field experiment was conducted during kharif 2019 at ARS farm at College of Agriculture, Umedganj, Kota. The soil of the experimental site was medium black (*Vertisols*) having clay loam texture. The experiment consisted of 10 treatments viz. T₀ - Control (Absolute), T₁ - RDF (20-40-40), T₂ - 75% RDF + FYM (2.0 t ha⁻¹), T₃ -

50% RDF + FYM (4.0 t ha⁻¹), T₄ - 75% RDF + FYM (2.0 t ha⁻¹) + *Rhizobium*, T₅ - 50% RDF + FYM (4.0 t ha⁻¹) + *Rhizobium*, T₆ - 75% RDF + Vermicompost (1.0 t ha⁻¹), T₇ - 50% RDF + Vermicompost (2.0 t ha⁻¹), T₈ - 75% RDF + Vermicompost (1.0 t ha⁻¹) + *Rhizobium*, T₉ - 50% RDF + Vermicompost (2.0 t ha⁻¹) + *Rhizobium*. The experiment was laid out in randomised block design and was replicated thrice. Kota is characterized by sub-tropical, semi-arid climate, temperature often exceed 40°C during summer and sometimes touches 45°C, minimum temperature falls below 5°C during winter season. The average rainfall 575mm, contributed from south-west monsoon during July to September. Soybean crop sown using variety JS-335 with 80kg ha⁻¹ seed rate in kharif 2019. Soil samples were analysed for electrical conductivity (EC) of the supernatant liquid with the help of Solubridge or conductivity meter in 1: 2, soil water solution [3]. pH measured by Glass electrode pH meter in 1: 2, soil water solution [3]. Organic carbon in the soil was estimated by Walkley and black [4] Rapid titration method. Soil samples were analysed for available N using alkaline potassium permanganate, which oxidized and hydrolysed the organic matter present in the soil [5]. Available phosphorous was determined through the UV-spectrophotometer by extracting the soil with 0.5 N NaHCO₃ pH 8.5 [6] and measuring the intensity of blue colour developed by ammonium molybdate–stannous chloride. Available K in soil was determined by extracting with 1N neutral ammonium acetate [7]. The available micronutrient was determined using DTPA extractant as described by Lindsay and Norvell (1978). Use the filtrate for measurement of Fe, Mn, Zn and Cu on Atomic Absorption Spectrophotometry (AAS).

3. RESULTS AND DISCUSSION

3.1 Soil Physico-chemical Properties

The results indicated that soil parameters after harvest showed non- significant variation with pH and EC. Soil reaction (pH) in the experimental soil ranged from 7.52 to 7.60 and it was maximum for 100% RDF (T₁) and minimum for 50% RDF + Vermicompost (2.0 t ha⁻¹) + *Rhizobium* (T₉). Electrical Conductivity (EC) of the experimental soil ranged from 0.30 to 0.35 and it was maximum for 100% RDF (T₁) and minimum for 50% RDF + Vermicompost

Table 1. Initial physico-chemical properties of the soil of experimental field

Properties	Values	Method Employed
(A) Mechanical Composition		Hydrometer method [8]
Sand (%)	14.30	
Silt (%)	36.50	
Clay (%)	49.20	
Textural class	Clay loam	Texture Triangular diagram, [9]
(B) Physical properties		
Bulk density (Mg m^{-3})	1.38	Core Sampler method [10]
Particle density (Mg m^{-3})	2.66	Pycnometer or RD bottle method
Porosity (%)	48.2	Calculated by BD and PD
(C) Chemical properties		
Available N (kg ha^{-1})	214 kg ha^{-1}	Alkali permanganate method [5]
Available P (kg ha^{-1})	17 kg ha^{-1}	Olsen's method [6]
Available K (kg ha^{-1})	420 kg ha^{-1}	Neutral normal Ammonium acetate [7]
Organic carbon (%)	0.52	Walkley and Black [4] method
EC (dS m^{-1})	0.36	EC of soil samples was measured on Solubridge in soil:water solution (1:2) [3]
pH	7.6	Glass electrode pH meter in soil:water solution (1:2) [3]

(2.0 t ha^{-1}) + Rhizobium (T_9). Application of 50% RDF + Vermicompost (2.0 t ha^{-1}) + Rhizobium (T_9) found highest organic carbon (0.59%) content in soil after harvesting of crop which is significantly superior and at par with 75% RDF + Vermicompost (1.0 t ha^{-1}) + Rhizobium (T_8) and the value is 0.58%. The minimum organic carbon content was recorded in absolute control. Higher production of root biomass might have increased the organic carbon content Ram Lakshmi et al. [11]. These observations are in close agreements with the findings of Dhakal et al. [12] and Konthoujam et al. [13] who had indicated that the judicious integration of organic and inorganic sources of nutrition significantly improved the soil available N, P, K, and micronutrient and numerically change in soil pH, electrical conductivity and organic carbon content in soil.

3.2 Nutrient Availability in Soil after Harvesting the Crop and Nutrient Balance Sheet

The data on soil available nitrogen, phosphorus and potassium are presented in Table 2. It was observed that integrated nutrient management resulted into significant influence on available N, P, K in soil. The data showed that available nitrogen in soil was significantly improved with integrated application of chemical fertilizers, rhizobium and vermicompost with respect to the RDF alone and control plots during the experiment. As compared to the initial value of available nitrogen, the application of 75% RDF +

Vermicompost (1.0 t ha^{-1}) + Rhizobium improved its status after crop harvest. Although, available nitrogen under RDF after crop harvest remained almost same when compared to that of initial value. The nutrient management practices left the significant effect on available phosphorus and potassium in soil after harvest of soybean crop. The available nutrient status of the soil after harvesting the soybean crop showed considerable increment due to the soil application of different inorganic and organic sources. The highest soil available nitrogen (258 kg ha^{-1}), available phosphorus (21.22 kg ha^{-1}), available potassium (445.35 kg ha^{-1}) and micronutrients was recorded with the application of 50% RDF + Vermicompost (2.0 t ha^{-1}) + Rhizobium (T_9) after harvesting of crop which is significantly superior and at par with 50% RDF + FYM (4.0 t ha^{-1}) + Rhizobium (T_5). The increase in the available N, P, K and micronutrient content of soil might be due to release of those nutrients added through vermicompost with inorganic fertilizers to the soil after mineralization. Organic manures help to increase biological activity of soil microbes and improve soil structure, water holding capacity and other physico-chemical properties of soil Devi et al. [14]. These observations are in close agreements with the findings of Jadhav et al. [15], Dhakal et al. [12], Singh et al. [16], Konthoujam et al. [13], Meshram and Sapre [17], Bonde and Gawande [18], Shahina et al. [19] who had indicated that the judicious integration of organic and inorganic sources of nutrition significantly improved the soil available N, P, K, and micronutrient and

Table 2. Effect of integrated nutrient management on pH, EC, OC, N, P, K and DTPA extractable micronutrients in soil after harvesting the soybean crop

Treatment	pH	EC (dsm^{-1})	Organic carbon (%)	Nitrogen (kg ha^{-1})	Phosphorous (kg ha^{-1})	Potassium (kg ha^{-1})	Fe (mg kg^{-1})	Mn (mg kg^{-1})	Cu (mg kg^{-1})	Zn (mg kg^{-1})
T ₀ - Control (Absolute)	7.60	0.35	0.48	200.70	14.17	415.00	3.76	3.80	0.75	0.378
T ₁ - RDF (20-40-40)	7.65	0.36	0.50	213.19	18.22	421.67	4.33	4.33	0.76	0.53
T ₂ - 75% RDF + FYM (2.0 t ha^{-1})	7.59	0.34	0.51	225.29	19.57	426.67	4.47	4.47	0.76	0.55
T ₃ - 50% RDF + FYM (4.0 t ha^{-1})	7.59	0.34	0.51	221.11	19.17	425.67	4.37	4.43	0.76	0.54
T ₄ - 75% RDF + FYM (2.0 t ha^{-1}) + Rhizobium	7.57	0.31	0.57	248.99	20.64	434.00	5.37	4.90	0.77	0.92
T ₅ - 50% RDF + FYM (4.0 t ha^{-1}) + Rhizobium	7.53	0.33	0.57	250.00	20.35	436.33	5.00	4.73	0.77	0.79
T ₆ - 75% RDF+ Vermicompost (1.0 t ha^{-1})	7.58	0.33	0.53	233.33	19.98	430.33	4.83	4.70	0.77	0.61
T ₇ - 50% RDF + Vermicompost (2.0 t ha^{-1})	7.56	0.33	0.52	229.60	19.73	427.67	4.50	4.63	0.77	0.58
T ₈ - 75% RDF + Vermicompost (1.0 t ha^{-1}) + Rhizobium	7.53	0.30	0.58	254.77	21.53	441.33	5.43	5.10	0.77	0.93
T ₉ - 50% RDF + Vermicompost (2.0 t ha^{-1}) + Rhizobium	7.52	0.31	0.59	258.14	21.67	445.00	5.47	5.17	0.77	0.88
S.Em. \pm	0.05	0.01	0.01	5.65	0.43	1.61	0.12	0.07	0.01	0.02
CD at 0.05%	NS	NS	0.02	16.79	1.27	4.78	0.35	0.21	NS	0.05

Table 3. Nutrient balance sheet of nitrogen

Treatment	Nitrogen (kg/ha)						
	Initial soil N status (a)	N added (b)	N Uptake by crop (c)	Expected nutrient balance (d=(a+b)-c)	Actual nutrient balance (e)	Apparent gain/loss (f=e-d)	Actual difference of initial and final (g=e-a)
T ₀ - Control (Absolute)	210	0	51.06	158.94	200.7	41.76	-9.3
T ₁ - RDF (20-40-40)	210	20	61.25	168.75	213.19	44.44	3.19
T ₂ - 75% RDF + FYM (2.0 t ha ⁻¹)	210	25	71.99	163.01	225.29	62.28	15.29
T ₃ - 50% RDF + FYM (4.0 t ha ⁻¹)	210	30	71.02	168.98	221.11	52.13	11.11
T ₄ - 75% RDF + FYM (2.0 t ha ⁻¹) + Rhizobium	210	25	111.11	133.89	248.99	115.1	38.99
T ₅ - 50% RDF + FYM (4.0 t ha ⁻¹) + Rhizobium	210	40	96.61	153.39	250	96.61	40
T ₆ - 75% RDF+ Vermicompost (1.0 t ha ⁻¹)	210	25	85.15	149.85	233.33	83.48	23.33
T ₇ - 50% RDF + Vermicompost (2.0 t ha ⁻¹)	210	35	82.99	162.01	229.6	67.59	19.6
T ₈ - 75% RDF + Vermicompost (1.0 t ha ⁻¹) + Rhizobium	210	35	117.93	127.07	254.77	127.7	44.77
T ₉ - 50% RDF + Vermicompost (2.0 t ha ⁻¹) + Rhizobium	210	40	98.52	151.48	258.14	106.66	48.14

Table 4. Nutrient balance sheet of phosphorous

Treatment	Phosphorous (kg/ha)						
	Initial soil P status (a)	P added (b)	P Uptake by crop (c)	Expected nutrient balance (d=(a+b)-c)	Actual nutrient balance (e)	Apparent gain/loss (f=e-d)	Actual difference of initial and final (g=e-a)
T ₀ - Control (Absolute)	17	0	5.5	11.5	14.17	2.67	-2.83
T ₁ - RDF (20-40-40)	17	40	6.56	50.44	18.22	-32.22	1.22
T ₂ - 75% RDF + FYM (2.0 t ha ⁻¹)	17	34	7.69	43.31	19.57	-23.74	2.57
T ₃ - 50% RDF + FYM (4.0 t ha ⁻¹)	17	28	7.59	37.41	19.17	-18.24	2.17
T ₄ - 75% RDF + FYM (2.0 t ha ⁻¹) + Rhizobium	17	34	11.76	39.24	20.64	-18.6	3.64
T ₅ - 50% RDF + FYM (4.0 t ha ⁻¹) + Rhizobium	17	28	10.26	34.74	20.35	-14.39	3.35
T ₆ - 75% RDF+ Vermicompost (1.0 t ha ⁻¹)	17	48	9.06	55.94	19.98	-35.96	2.98
T ₇ - 50% RDF + Vermicompost (2.0 t ha ⁻¹)	17	56	8.83	64.17	19.73	-44.44	2.73
T ₈ - 75% RDF + Vermicompost (1.0 t ha ⁻¹) + Rhizobium	17	48	12.41	52.59	21.53	-31.06	4.53
T ₉ - 50% RDF + Vermicompost (2.0 t ha ⁻¹) + Rhizobium	17	56	10.47	62.53	21.67	-40.86	4.67

Table 5. Nutrient balance sheet of potassium

Treatment	Potassium (kg/ha)						
	Initial soil K status (a)	K added (b)	K Uptake by crop (c)	Expected nutrient balance (d=(a+b)-c)	Actual nutrient balance (e)	Apparent gain/loss (f=e-d)	Actual difference of initial and final (g=e-a)
T ₀ - Control (Absolute)	420	0	45.77	374.23	415	40.77	-5
T ₁ - RDF (20-40-40)	420	40	53.26	406.74	421.67	14.93	1.67
T ₂ - 75% RDF + FYM (2.0 t ha ⁻¹)	420	40	61.74	398.26	426.67	28.41	6.67
T ₃ - 50% RDF + FYM (4.0 t ha ⁻¹)	420	40	60.93	399.07	425.67	26.6	5.67
T ₄ - 75% RDF + FYM (2.0 t ha ⁻¹) + Rhizobium	420	40	90.57	369.43	434	64.57	14
T ₅ - 50% RDF + FYM (4.0 t ha ⁻¹) + Rhizobium	420	40	80.26	379.74	436.33	56.59	16.33
T ₆ - 75% RDF+ Vermicompost (1.0 t ha ⁻¹)	420	50	71.57	398.43	430.33	31.9	10.33
T ₇ - 50% RDF + Vermicompost (2.0 t ha ⁻¹)	420	60	69.8	410.2	427.67	17.47	7.67
T ₈ - 75% RDF + Vermicompost (1.0 t ha ⁻¹) + Rhizobium	420	50	92.92	377.08	441.33	64.25	21.33
T ₉ - 50% RDF + Vermicompost (2.0 t ha ⁻¹) + Rhizobium	420	60	81.78	398.22	445	46.78	25




Fig. 1. Effect of integrated nutrient management on nutrient content in soil after harvesting the soybean crop




Fig. 2. Effect of integrated nutrient management on micronutrient content (mg kg^{-1}) in soil after harvesting the soybean crop

and numerically change in soil pH, electrical conductivity and organic carbon content in soil.

3.3 Micronutrient Content Fe, Mn, Zn and Cu (mg kg^{-1})

The analysed data for micronutrient also indicated that after crop harvest, the available (DTPA-extractable) Zn and Fe in the soil remained affected by nutrient management practices. Application of 50% RDF + Vermicompost (2.0 t ha^{-1}) + Rhizobium (T_9)

recorded the highest Fe, Mn and Zn in soil, which was significantly superior to other treatments but found at par with treatment 75% RDF + Vermicompost (1.0 t ha^{-1}) + Rhizobium (T_8) (5.43 mg kg^{-1}), minimum Fe content in soil was recorded under absolute control (3.76 mg kg^{-1}). The effect of integrated nutrient management on Cu content in soil was found non-significant and DTPA- Cu ranged from 0.75 to 0.77 mg kg^{-1} and it was maximum for 75% RDF + Vermicompost (1.0 t ha^{-1}) + Rhizobium and minimum for absolute control. These observations are in close

agreements with the findings of Dhakal et al. [12], Singh et al. [16], Laharia et al. [20], Dadhich and Somani, [21].

4. CONCLUSION

The available nutrient status of the soil after harvesting the soybean crop showed considerable increment due to the soil application of different inorganic and organic sources. The highest soil available nitrogen (258 kg ha⁻¹), available phosphorus (21.67 kg ha⁻¹) and available potassium (445.35 kg ha⁻¹), micronutrient and organic carbon (0.59) was recorded with the application of 50% RDF + Vermicompost (2.0 t ha⁻¹) + Rhizobium (T₉). Hence, this combination of integrated nutrient management proved as productive, remunerative and beneficial for soil health in soils of ARS, Kota.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Anonymous 2017-18. Annual report. DES, Ministry of Agricultural & Farmer's Welfare, Government of India, Krishi Bhawan, New Delhi.
- Todkari AK, Dwivedi AK, Dikshit PR, Tiwari A. Vermicompost and organic matter consists of macro and micronutrients, plant growth promoting substance. necessary for plant growth. Journal of Indian Society of Soil Science. 2001;50(4):472-475.
- Jeckson ML. Soil chemical analysis. Prentice Hall of India, New Delhi; 1967.
- Walkley A, Black CA. Estimation of soil organic carbon by the chromic acid titration method. Soil Science. 1934;37(1):29-38.
- Subbiah BV, Asija GL. A rapid method for the estimation of available nitrogen in soil. Current Science. 1956;(25):259-260.
- Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimating of available phosphorus in soil by extraction with sodium bicarbonate USDA Circular. 1954;939.
- Hanway JJ, Heidel H. Soil analysis method as used in Iowa state college Soil testing Laboratory. Iowa State College of Agriculture Bulletin. 1952;57;1-3.
- Bouyoucos GJ. Hydrometer method for making particle size analysis of soil. Agronomy Journal. 1962;55:464-465.
- Brady, N.C. 1983. The nature and properties of soils. MacMillan Publishing Company, New York and Collier MacMillan Publisher, London pp. 750.
- Piper CS. Soil and plant analysis. Hans Publication, Bombay; 1950.
- Ram Lakshmi, Ch. S., Chandra Sekhar Rao, P. Sreelatha, T., Padmaja, G., Madhavi and Rao, P.V. Effect of integrated nutrient management on soil properties, yield and nutrient uptake in rice-green gram cropping system in an Inceptisol of Andhra Pradesh. Journal of the Indian Society of Soil Science. 2015;63(4):400-405.
- Dhakal Y, Meena RS, Kumar S. Effect of INM on nodulation, yield, quality and available nutrient status in soil after harvest of greengram. Legume Research. 2016;39(4):590-594.
- Konthoujam ND, Tensubam BS, Athokpam HS, Naorem BS, Shamurailatpam D. Influence of inorganic, biological and organic manures on nodulation and yield of soybean (*Glycine max* L. Merrill) and soil properties. Australian Journal of Crop Science. 2013;7(9):1407-1415.
- Devi KN, Singh TB, Athokpam HS, Singh NB, Shamurailatpam D. Influence of inorganic, biological and organic manures on nodulation and yield of soybean (*Glycine max* Merril L.) and soil properties. Australian journal of crop science. 2013;7(9):1407-1415.
- Jadhav PM, Rane SD, Ranshur NJ, Todmal SM. Effect of integrated nutrient management on quality and yield of soybean. The Asian Journal of Soil Science. 2007;2(1):68-73.
- Singh M, Kumar N, Das A, Rakshit R. Soybean Response to Organic and Inorganic Fertilization on Aquic Hapludoll Soil. Environment & Ecology. 2014;34(4B):1634-1638.
- Meshram N, Sapre N. Effect of integrated nutrient management on growth and yield of soybean [*Glycine max* (L.) Merrill]. Plant Archive. 2019;19(2):2933-2936.
- Bonde AS, Gawande SN. Effect of integrated nutrient management on soil properties and yield of soybean (*Glycine max*). Annals of Plant and Soil Research. 2017;19(2): 205–209.
- Shahina TK, Sammi UK, Singh VM, Biswas AK. Changes in organic and inorganic forms of nitrogen in a Typic Haplustert under soybean-wheat system

- due to conjoint use of inorganic fertilizers and organic manures. Journal of Indian Society of Soil Science. 2010;58(1):78-85.
20. Laharia GS, Patil DU, Damre PR. Effect of organic sources on soil fertility, nutrient uptake and yield of soybean. Crop Research. 2013;45(1to3):155-159.
21. Dadhich S, Somani L. Effect of integrated nutrient management in a soybean-wheat crop sequence on the yield, micronutrient uptake and post-harvest availability of micronutrients on typic ustochrepts soil. Acta Agronomica Hungarica. 2007; 55(2):205-216.

© 2022 Meena et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/87084>