



Magic Polygons and Combinatorial Algorithms

Daniel Dias Augusto ^{a*}

^aDepartamento de Matemática - Universidade de Brasília (UnB), Distrito Federal-70297-400, Brasil.

Authors' contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/ARJOM/2022/v18i1030424

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91120>

Received: 16 June 2022

Accepted: 22 August 2022

Published: 31 August 2022

Original Research Article

Abstract

In this work, we study the Magic Polygons of order 3 ($P(n, 2)$) and we introduce some properties that were useful to build an algorithm to find out how many Magic Polygons exists for the regular polygons up to 24 sides. The concepts of Equivalents Magic Polygons and Derivatives Magic Polygons which allowed to classify, and avoid ambiguity about the representations of such elements, are also introduced.

Keywords: Magic polygons; combinatorial algorithms; symmetric group; dihedral group.

2010 Mathematics Subject Classification: 53C25; 83C05; 57N16.

1 Introduction

Magic Polygons are objects that had its inspiration on the classics Magics Squares, and these are known from a long time ago, and there are algorithms to build Magic Squares of different orders, as we can see in [1, 2, 3, 4, 5, 6]. Furthermore, such objects has applications in fields like Computer Science Physics and Cryptography according to [7, 8, 9].

Magic Squares of order n are arrangement of n^2 squares in n rows and n columns and filled with numbers from 1 to n^2 by a way that the sum along the rows columns and diagonals is constant and called magic sum. The Magic Square of order 1 has only one element, therefore is trivial. There's no

*Corresponding author: E-mail: nieldiasguto@gmail.com, ddiasguto@protonmail.com;

Magic Square of order 2, because is impossible to build it without a repetition of values. It follows that the Magic Square of order 3 is the smaller non trivial Magic Square, and unless considering rotation and reflections there are only one possibility for the Magic Square of order 3. Ian Stewart in his book “*Incredible Numbers*”, claim that there are 880 possibles Magic Squares of order 4, and 275.305.224 for a Magic Squares of order 5. The number of possibilities for the Magic Squares of order 6 is unknown, but estimatives claim it is about 1.77×10^{19} .

However, we know that there are at least one Magic Polygon for all n even greater than six, according the algorithm provided in [11], except the magic square, is unknown how many Magic Polygons exists for n even equal or greater than six. This is the main goal of this work, show how many Magic Polygons exists from the Hexagon ($n = 6$) until the regular polygon of 24 sides.

To reach it, it was necessary a implementation of an algorithm that could find all possibles Magic Polygons for each regular polygon. Firstly, is provided the definition of a Magic Polygon, and then is showed how a Magic Polygon can be represented as an element of the Symmetric Group. Is also showed the properties that allow us to implement the algorithm with some basic optimizations.

Two relevant concepts are here introduced, *Equivalents Magic Polygons* and *Derivatives Magic Polygons*, these concepts avoid ambiguity in the count of the Magic Polygons, because, as shall be showed we can represent the Magic Polygons by many ways, and we can get infinities Magic Polygons from one, but with these two concepts, we can classify them.

By making an bond between the magic polygons and some elements of the symmetric groups and bring these two new elements, some possibilities of study appears.

In the last section, there are a table with the numbers of possibilities for all Magic Polygons up to 24 sides, and the main goal of this work is reached.

2 Magic Polygons - Definition and Properties

In this section, is adopted the definition of Magic Polygons provided in [10], which is a generalization of the Magic Polygon according [11], the propositions here not proved, can be viewed in these works.

Definition 2.1. Let Ω be a set of $\frac{k}{2}$ regular polygons on plane with n sides and corresponding parallel sides and centered in a central point C .

A magic polygon $P(n, k)$ of n sides and order $k + 1$ is a set of $\frac{k^2n}{2} + 1$ points satisfying the following conditions:

- (i) Points of magic polygon are labeled by distinct values from 1 to $\frac{k^2n}{2} + 1$;
- (ii) One point of a magic polygon is the central point C ;
- (iii) $\frac{kn}{2}$ points of magic polygon are vertices of the $\frac{k}{2}$ regular polygons of Ω ;
- (iv) The magic polygon has $k - 1$ intermediate points on each edge of regular polygons in Ω , which gives a total of $\frac{(k-1)nk}{2}$ intermediate points.
- (v) Segments with diametrically opposite ends of the larger polygon of Ω intersecting the central point contain $k + 1$ points of the magic polygon;
- (vi) Segments with ends at two adjacent vertices of a polygon of Ω contains $k + 1$ points of the magic polygon;
- (vii) The sum of values corresponding to the $k + 1$ points on each segment defined in (iv) and (v) is a fixed value u , called of **magic sum**.




Fig. 1. Examples of Magic Polygons $P(8, 2)$ and $P(4, 4)$ respectively

Theorem 2.1. A Magic Polygon $P(n, k)$ has the following properties:

- (i) the magic sum is: $(k + 1) \frac{k^2 n + 4}{4}$;
- (ii) the value of the central point is: $\frac{k^2 n + 4}{4}$
- (iii) the sum S_j of the values representing to j -th points partitioning each edge on magic polygon chosen clockwise is:

$$S_j = \frac{kn(k^2 n + 4)}{8}$$

These properties are proved in [10].

If we take $k = 2$, Magic Polygons $P(n, 2)$ are formed by one regular polygon, the vertex, and the edges middle points and its geometric center. Therefore has $2n + 1$ elements. Its magic sum is $3(n + 3)$, the central point has the value $n + 1$ and the sum S_j of all elements in the vertex is $n(n + 1)$ (the same for the sum of the elements in the edges middle points). We note that the magic sum u , is three time the central point c ($u = 3(n + 1)$).

In [10, 11] is proved that a Magic Polygon $P(n, 2)$ exist if, and only if, the regular polygon has a number of sides even and greater than six. All approach made in this work is valid only for the Magic Polygons of order 3 ($P(n, 2)$).

Is easy to see that Magic Polygons has a discrete and finite framework, therefore we can adopt an algebraic representation for such objects in order to be aided by algebraic concepts to explore its properties. Once the geometric framework of magic polygons has the symmetric properties natives in the regular polygons, and its elements are positives integers numbers along of the perimeter, the concepts of *dihedral group*, *finite groups* $\mathbb{Z}/n\mathbb{Z}$ and *symmetric groups*, shall be very important.

As mentioned, a Magic Polygon is about the natural numbers $1, 2, 3, \dots, 2n + 1$ along the perimeter of the regular polygon, so it is nothing beyond a permutation of such elements, therefore a Magic Polygon can be algebraically represented in the well known notation for symmetric groups:

$$\left(\begin{array}{cccccccc} 1 & 2 & \dots & n & n + 1 & n + 2 & \dots & 2n & 2n + 1 \\ f(1) & f(2) & \dots & f(n) & n + 1 & f(n + 2) & \dots & f(2n) & f(2n + 1) \end{array} \right)$$

where the numeration direction is counter-clockwise, and the first element is by convention the right vertex nearest the symmetric vertical axis of the polygon, and above the horizontal symmetric axis. As we can see, the image of $n + 1$ is $n + 1$, this is in fact the central point that stands in the polygon geometric center.

Definition 2.2. An element of the symmetric group of $(2n+1)!$ order, is an algebraically representation of the Magic Polygon of n if the following conditions are satisfied:

$$\left\{ \begin{array}{l} i) \quad f(n + 1) = n + 1; \\ ii) \quad \forall i, 1 \leq i \leq n, f(i) + f(i + (n + 1)) = 2c; \\ iii) \quad \forall i \text{ odd}, 1 \leq i < n - 1, f(i) + f(i + 1) + f(i + 2) = u; \\ iv) \quad f(n - 1) + f(n) + f(n + 2) = u; \\ v) \quad \forall i \text{ even}, n + 2 \leq i < 2n - 1, f(i) + f(i + 1) + f(i + 2) = u; \\ vi) \quad f(2n) + f(2n + 1) + f(1) = u; \end{array} \right. \quad (2.1)$$

The first condition is due to vertex root of the Magic Polygon, which value is fixed with $n + 1$, the second is due to the opposite elements in the perimeter of the polygon, once both stands in the same symmetric axis, that splits the polygon in two, they are the elements $f(i)$ and $f(i + n + 1)$, because the central point is the element $f(n + 1)$, thus $f(i) + f(i + n + 1) + c = u$, once $u = 3c$ we have $f(i) + f(i + n + 1) = 2c$. The next condition, is due to the sum of the elements along the edge of regular polygon, by definition, their sum must match u . Once $f(1)$ is by definition a vertex, for all i odd less than $n - 1$ it follows that $f(i) + f(i + 1) + f(i + 2) = u$. We have, $f(n + 1)$ as central point, thus $f(n - 1), f(n)$ e $f(n + 2)$ are in the same edge and their sum shall be the magic sum. From the element $n + 2$ beyond, if i is even, so $f(i), f(i + 1)$ and $f(i + 2)$ shall be in the same edge, and therefore their sum must be the magic sum. It give us the fifth condition. By last, the sixth condition refers the fact that the elements $f(2n), f(2n + 1)$ and $f(1)$ stands in the same edge which close the polygon.

By adopting this convention, we can represent the Magic Polygon $P(8, 2)$ on the Fig. 1 by the following way:

$$\left(\begin{array}{cccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \mathbf{9} & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 \\ 15 & 11 & 1 & 12 & 14 & 5 & 8 & 16 & \mathbf{9} & 3 & 7 & 17 & 6 & 4 & 13 & 10 & 2 \end{array} \right) \quad (2.2)$$

It follows that the Magic Polygons are elements of the symmetric group, but as a subset, they didn't make a subgroup, because it's obvious that the identity function, which is the neutral element in the symmetric group, does not represent a Magic Polygon. From this representation, to find all Magic Polygons for a given n , one option is by declaring $2n$ variables, and visiting all permutations of the set:

$$\{1, 2, 3, \dots, 2n, 2n + 1\} - \{n + 1\}$$

and verify, which ones satisfies the the conditions listed in Definition 2.2. However, such algorithm would return the same permutation many times, and this is not good for the computation.

Based in some Magic Polygons properties that shall be here approached, will be possible find all Magic Polygons for a given n by declaring only n variables, and reducing how many times the same permutation is returned. Simplifying and fastening the computational process.

Theorem 2.2. *The algebraic representation of a Magic Polygon is well defined by the values $f(1), f(2), \dots, f(n)$. Putting in other words, a Magic Polygon can be represented by a n -permutation of the set $\{1, 2, 3, \dots, 2n + 1\}$.*

Proof. Let $f : \{1, 2, \dots, n\} \mapsto \{1, 2, 3, \dots, 2n + 1\}$ an one-to-one function which has the following properties:

$$\begin{aligned} i) & f(i) \neq n + 1 \forall i \in \{1, 2, \dots, n\}; \\ ii) & f(i) + f(i + 1) + f(i + 2) = u \forall i \text{ odd} \in \{1, 2, \dots, n - 2\}; \\ iii) & f(i) + f(j) \neq 2c \forall i, j \in \{1, 2, \dots, n\}; \\ iv) & f(n - 1) + f(n) - f(1) = c; \end{aligned} \tag{2.3}$$

i.e., f defines a n -permutation of $\{1, 2, 3, \dots, 2n + 1\}$.

Let's extend the function f for the set $\{1, 2, 3, \dots, 2n + 1\}$ in order to generate an element belonging the symmetric group of order $(2n + 1)!$ this way:

- $f(n + 1) = n + 1$;
- $\forall i \in \{1, 2, \dots, n\}, f(i + n + 1) = 2c - f(i)$

so we have the σ element:

$$\sigma = \left(\begin{array}{cccccccccc} 1 & 2 & \dots & n & n + 1 & n + 2 & \dots & 2n & 2n + 1 \\ f(1) & f(2) & \dots & f(n) & n + 1 & f(n + 2) & \dots & f(2n) & f(2n + 1) \end{array} \right) \tag{2.4}$$

Affirmation: σ represents a Magic Polygon.

The injectivity of original function f with the properties i and iii ensures the bijectivity of the extended function for the set $\{1, 2, \dots, 2n + 1\}$. By property ii , for all i odd $1 \leq i < n - 1 \Rightarrow f(i) + f(i + 1) + f(i + 2) = u$ thus:

$$\begin{aligned} f(i + n + 1) + f(i + n + 2) + f(i + n + 3) &= \\ (2c - f(i)) + (2c - f(i + 1)) + (2c - f(i + 2)) &= \\ 6c - (f(i) + f(i + 1) + f(i + 2)) &= \\ 6c - 3c = 3c = u \end{aligned} \tag{2.5}$$

this way, for all i even $n + 2 \leq i \leq 2n - 2$, $f(i) + f(i + 1) + f(i + 2) = u$. By property iv , $f(n - 1) + f(n) - f(1) = c$ as $f(i + n + 1) = 2c - f(i) \Rightarrow -f(1) = 2c - f(n + 2)$ therefore:

$$\begin{aligned} f(n - 1) + f(n) - 2c + f(n + 2) &= c \\ f(n - 1) + f(n) + f(n + 2) &= 3c = u \end{aligned} \tag{2.6}$$

this way sum of the elements $f(n - 1)$, $f(n)$ and $f(n + 2)$ is the Magic Sum:

Now we have:

$$\begin{aligned} f(2n) &= 2c - f(n - 1) \\ f(2n + 1) &= 2c - f(n) \\ f(1) &= 2c - f(n + 2) \end{aligned} \tag{2.7}$$

As proved, $f(n - 1) + f(n) + f(n + 2) = u$, the sum $f(2n) + f(2n + 1) + f(1)$ shall be:

$$\begin{aligned} 2c - f(n - 1) + 2c - f(n) + 2c - f(n + 2) &= \\ 6c - (f(n - 1) + f(n) + f(n + 2)) &= \\ 6c - 3c = 3c = u \end{aligned} \tag{2.8}$$

Therefore, in conformity with the Definition 2.2 the sum of all elements in the edges of Regular Polygons satisfies the magic sum.

Now, it's necessary verify if the sums of the elements on the symmetric axis also satisfies the magic sum.

The sums in the symmetric axis, matches the pattern $f(i) + c + f(i + n + 1)$, where $1 \leq i \leq n$. Once $f(i + n + 1) = 2c - f(i)$ the result follows. □

The Definition 2.2 introduces the Magic Polygons as a permutation of the set $\{1, 2, \dots, 2n + 1\}$ which satisfies a few properties. In other words, the process of find Magic Polygons is a combinatorial issue. And based in the Theorem 2.2 is enough to declare only n variables in the algorithm.

3 Equivalent Magic Polygons

It's easy to see geometrically that rotations based on the geometric center of the polygon, and reflection on the symmetric axis, provide us an different way to represent what is the same Magic Polygon. A first look may induce to think that is another Magic Polygon, but in fact, is the same, represented by a distinct way.

This encourage us to define Equivalent Magic Polygons, distinct representations that we can obtain from a Magic Polygon by applying two operations, the rotation (r_o) and the reflection (r_e).

In this section, and in the one following, we adopt the notation:

- $a \leftrightarrow b$ means that the value a and the value b switches places;
- $a \rightarrow b$ means that the value a is replaced by the value b ;

Definition 3.1. Let σ an element of symmetric group that represents a Magic Polygon. We say that the element σ' of S_{2n+1} is an Equivalent Magic Polygon of σ if σ' can be obtained from σ applying the following operations a finite number of times:

ii) Rotation in the geometric center of polygon:

$r_o(\sigma)$:

$$\begin{pmatrix} 1 & 2 & \dots & n & n+1 & n+2 & \dots & 2n & 2n+1 \\ f(2n) & f(2n+1) & \dots & f(n-2) & n+1 & f(n) & \dots & f(2n-2) & f(2n-1) \end{pmatrix}$$

ii) Reflection applied in vertex:

$r_{ev}(\sigma)$:

$$\begin{pmatrix} 1 & 2 & \dots & n & n+1 & n+2 & \dots & 2n & 2n+1 \\ f(1) & f(2n+1) & \dots & f(n+3) & n+1 & f(n+2) & \dots & f(3) & f(2) \end{pmatrix}$$

If $i \oplus j = 2$ on group $\mathbb{Z}/(2n+1)\mathbb{Z} \Rightarrow f(i) \leftrightarrow f(j)$

iii) Reflection applied in the Edge Middle Point:

$r_{em}(\sigma)$:

$$\begin{pmatrix} 1 & 2 & \dots & n & n+1 & n+2 & \dots & 2n & 2n+1 \\ f(2n) & f(2n-1) & \dots & f(n) & n+1 & f(n-1) & \dots & f(2) & f(2n+1) \end{pmatrix}$$

If $i \oplus j = 0$ on group $\mathbb{Z}/(2n+1)\mathbb{Z} \Rightarrow f(i) \leftrightarrow f(j)$

In reflection operation applied on vertex, the elements $f(n+2)$ e $f(n+1)$ would switch their places, however, we know that the element $f(n+1)$ is a fixed element, thus, both remain in their places. In case $i = 1$, $f(i)$ would switch place with itself, for $1 + 1 = 2 \in \mathbb{Z}/(2n+1)\mathbb{Z}$, therefore it also remain without change. Geometrically, the line that contain the elements $f(1)$, $f(n+1)$ e $f(n+2)$ defines the symmetric axis by which we the Magic Polygon is reflected, and, as expect, such elements does not change their positions.

In reflection operation applied on edge middle point, something similar occurs, the elements $f(n+2)$ e $f(n+1)$ would switch their places, however, once again $f(n+1)$ is fixed, therefore both remain in their places. And the element $f(2n+1)$ would switch place with itself, therefore remain with no change. Again such elements stands in the symmetric axis by which the polygon is reflected.

Such operations are due to the Dihedral Group (D_n) that acts in the regular polygon where is built the Magic Polygon. We note that in a Magic Polygon, what really matters, is the adjacent elements, and the elements that lies in the end of the lines that defines the symmetric axis. In a Dihedral Group, such arrangement is not affected. Therefore, every element of a Dihedral Group gives us a distinct mode to represent exactly the same Magic Polygon. So we have the following proposition:

Proposition 3.1. *Every Magic Polygon of n sides can be represented by $2n$ distinct ways.*

Considering the way fo the numeration, defined above, and the definition of the r_o operation, we can see that such operation occurs in the counter-clockwise way. And the reflection is applied only in two axis, one of them is defined by two vertex, and the another by two edge middle points. However is known that a regular polygon has n symmetric axis, but there's no loss of generality here. Because the definition takes two axis that are like the "representatives" of the others, thus, if a composition of operations is made, we can reflect the Magic Polygon in any symmetric axis, by the following way:

$$r_o^{n-k}(r_e(r_o^k(\sigma))) \tag{3.1}$$

where r^k means the r_o applied k times, where k is the numbers of times necessary for the choose axis match one of those given in the definition. And r_o^{n-k} make the axis back to its original position. Because $r_o^{n-k}(r_o^k(\sigma)) = r_o^n = e$ (neutral element). Note that the Dihedral Group is not commutative, so the order of the operations in 3.1 cannot be changed.

Taking as example the Polygon $P(8,2)$ showed in the Figure 1, if we applied the r_o operation we shall have:

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \mathbf{9} & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 \\ 15 & 11 & 1 & 12 & 14 & 5 & 8 & 16 & \mathbf{9} & 3 & 7 & 17 & 6 & 4 & 13 & 10 & 2 \end{pmatrix}$$

$$r_o(\sigma) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \mathbf{9} & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 \\ 10 & 2 & 15 & 11 & 1 & 12 & 14 & 5 & \mathbf{9} & 8 & 16 & 3 & 7 & 17 & 6 & 4 & 13 \end{pmatrix}$$

The Fig. 2 show both representations:

Again, taking the Magic Polygon $P(8,2)$, if we applying in it the reflection in the vertex, we have:




Fig. 2. On the left σ , On the right $r_o(\sigma)$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \mathbf{9} & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 \\ 15 & 11 & 1 & 12 & 14 & 5 & 8 & 16 & \mathbf{9} & 3 & 7 & 17 & 6 & 4 & 13 & 10 & 2 \end{pmatrix}$$

$$r_{ev}(\sigma) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \mathbf{9} & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 \\ 15 & 2 & 10 & 13 & 4 & 6 & 17 & 7 & \mathbf{9} & 3 & 16 & 8 & 5 & 14 & 12 & 1 & 11 \end{pmatrix}$$




Fig. 3. On the left σ On the right $r_{ev}(\sigma)$

4 Derivatives Magic Polygons

By definition, the Magic Polygons, are built with the natural sequence from 1 to $2n + 1$, and the value $n + 1$ is fixed in the geometric center of the polygon. But is easy to see that if any constant k is added to every values in the Magic Polygon, the sum keeps constant, because, once $x + y + z = u$ so: $(x + k) + (y + k) + (z + k) = u + 3k$ and $k \in \mathbb{Z}$. Technically this is not a Magic Polygon, because according the definition, a Magic Polygon is strictly build with the values $1, 2, \dots, 2n + 1$.

An particular case is the null magic sum, this can be done by subtracting $n + 1$ from every element from a Magic Polygon, the Fig. 4 illustrates that.




Fig. 4. Null Magic Sum

More generally, it is possible to replace every value x in the Magic Polygon by $x - th$ value from an arithmetic progression that the sum keeps constant, this is proved in the Theorem below.

Theorem 4.1. Let σ an element of the symmetric group that represents a Magic Polygon, and be $a_1, a_2, a_3 \dots, a_{2n}$, a_{2n+1} an arithmetic progression where the first value is a_0 and common difference r . The replacement of every $f(i)$ for $a_{f(i)}$, change the magic sum to $3(a_1 + nr)$ and the value of the central point to $a_1 + nr$.

Proof. It is known that if a_1 is the first term of an arithmetic progression, the $n - th$ is: $a_1 + (n - 1)r$. Let $f(i) + c + f(i + n + 1) = u$ any sum along the symmetric axis of the polygon, due to the rotation (r_σ) , let's suppose, without loss of generality $f(i) > f(i + n + 1)$, where $1 < i < n$. So we have $f(i) = c + k$ for some k positive, and $f(i + n + 1) = c - k$, because as proved, $f(i) + f(i + n + 1) = 2c$. Once we have $c = n + 1$ so:

$$\begin{aligned} f(i) &= c + k = n + 1 + k \\ f(i + n + 1) &= c - k = n + 1 - k \end{aligned} \tag{4.1}$$

Therefore, the replacements shall be:

$$\begin{aligned} f(i) &\rightarrow a_1 + ((n + 1 + k) - 1)r \\ f(i + n + 1) &\rightarrow a_1 + ((n + 1 - k) - 1)r \\ c &\rightarrow a_1 + ((n + 1) - 1)r \end{aligned} \tag{4.2}$$

i.e. $c = a_1 + nr$, and the new sum shall be:

$$\begin{aligned} a_1 + ((n + 1) - 1)r + a_1 + ((n + 1 + k) - 1)r + a_1 + ((n + 1 - k) - 1)r &= \\ 3a_1 + nr + (n + k)r + (n - k)r &= \\ 3a_1 + (3n + k - k)r &= \\ 3a_1 + 3nr = 3(a_1 + nr) \end{aligned} \tag{4.3}$$

Now, we have to verify the sum along the edges. Let $i, j, k \in \{1, 2, 3 \dots 2n + 1\} - \{n + 1\}$ where $f(i), f(j)$ e $f(k)$ lies in the same edge, so: $f(i) + f(j) + f(k) = u$.

$$\begin{aligned} f(i) &\rightarrow a_1 + (f(i) - 1)r \\ f(j) &\rightarrow a_1 + (f(j) - 1)r \\ f(k) &\rightarrow a_1 + (f(k) - 1)r \end{aligned} \tag{4.4}$$

Therefore:

$$\begin{aligned} a_1 + (f(i) - 1)r + a_1 + (f(j) - 1)r + a_1 + (f(k) - 1)r = \\ 3a_1 + (f(i) + f(j) + f(k) - 3)r = \end{aligned} \quad (4.5)$$

By hypothesis: $f(i) + f(j) + f(k) = u = 3n + 3$, so we have:

$$3a_1 + (3n + 3 - 3)r = 3(a_1 + nr) \quad (4.6)$$

□

If $a_1 = 1$ and $r = 1$ we get the magic sum $3n+3$ for the natural sequence $\{1, 2 \dots 2n+1\}$ as we know.

Definition 4.1. The Magic Polygons built from an arithmetic progression that is not the natural, as procedure described in the Theorem 4.1, are called *Derivatives Magic Polygons*.

Corollary 4.1. Let, m, c' e r integers where $m = 3k$ for some $k \in \mathbb{Z}$ and $r \neq 0$, and be $n \geq 6$ even, so:

- i) Exists a Magic Polygon of n sides built by an arithmetic progression of common difference r and magic sum m ;
- ii) Exists a Magic Polygon of n sides built by an arithmetic progression of common difference r , and central point c' and magic sum $3c'$;

Proof. Based on the Theorem 2.1 the Magic Sum is $u = 3c \Rightarrow c = k$ and $u = 3k$, therefore according Theorem 4.1 is enough to get an arithmetic progression $a_1, a_2, \dots a_{2n+1}$ of common difference r where $a_{n+1} = k$. So we have:

$$a_{n+1} = a_1 + ((n + 1) - 1)r = k \Rightarrow a_1 = k - nr \quad (4.7)$$

This way, we have the common, r and the initial term a_1 given by the relation above. Now we just need to replace every value in a known Magic Polygon. The second case is the same, we just need to change k by c' .

□

5 Finding Magic Polygons

Based in the Theorem 2.2, it was implemented in the *C Programming Language* using loops and simple conditionals, programs to find all Magic Polygons up to 24 sides.

For each Magic Polygon of n sides, the algorithm has been implemented by declaring n integers variables in an array, (in the *C* language: `int f[n];`), and aided by the for loop, the value of each $f[i]$ variable was changed from 1 to $2n + 1$ and with the `if()` conditional, it was verified if the current permutation attended the restrictions showed in the Theorem 2.2. If so, the permutation was returned by the program and write in a plain text file. An auxiliary variable was initiated with value 0 and was increased by one each time a permutation was found, this way the number of magic polygons was counted.

Once each Magic Polygon can be represented in distinct ways, in virtue of the operation r_e and r_o here described, it would be desirable that the programs does not returned the same solutions many times.

By this way, the number 1 has been fixed as image of $f(1)$, i.e. as a vertex, and a second time as image of $f(2)$, i.e. as a middle point of a edge. This way, a permutation σ is returned, but the equivalents:

$$r_o(\sigma), r_o(r_o(\sigma)), r_o(r_o(r_o(\sigma))), \dots r_o^{n-1}(\sigma) \tag{5.1}$$

would not, for the value 1 won't be the image of the others vertex, and they are: $f(3), \dots, f(n-1), f(n+2) \dots f(2n)$. This way, all Magic Polygons where 1 is vertex are found. However, 1 can be middle point of an edge as well, thus running the program one more time, with the value 1 as image of $f(2)$, all Magic Polygons where 1 is middle point are found. Once 1 is necessarily a vertex or a edge middle point, all possible Magic Polygons for the current n are found.

However such procedure does not avoid the representations $r_{ev}(\sigma)$ and $r_{em}(\sigma)$ being returned, this way, all Magic Polygons are returned by the programs twice, being necessary to divide by 2 the quantity of Magic Polygons founded by the programs to know the real number. But by fixing 1 first as vertex and second as a edge middle point, is avoided that n Magic Polygons Equivalents being returned, because we can rotate the regular polygon n times until it back to its original position.

An example, for hexagon, the distinct solution are in number of 4, however, we can represent every by $2n$ distinct modes, as mentioned, due the action of dihedral group. If no element is fixed, so the program would return 48 permutations, with this optimization, only 8 is returned. This difference for polygons with a major number of sides became relevant, because as we shall see, the number of possibles Magic Possibles grows fast.

In order to provide examples, the tables below show all possibles Magic Polygons for the Magic Hexagon and the Magic Octagon, for both, they are in number of four. For the others Polygons, due the high number of possibles Magic Polygons, it would not be suitable to represent all in tables.

POSSIBILITIES FOR MAGIC HEXAGON

#	$f(1)$	$f(2)$	$f(3)$	$f(4)$	$f(5)$	$f(6)$	$f(7)$	$f(8)$	$f(9)$	$f(10)$	$f(11)$	$f(12)$	$f(13)$
1	1	8	12	4	5	3	7	13	6	2	10	9	11
2	1	9	11	4	6	2	7	13	5	3	10	8	12
3	8	1	12	5	4	11	7	6	13	2	9	10	3
4	9	1	11	6	4	12	7	5	13	3	8	10	2

POSSIBILITIES FOR MAGIC OCTAGON

#	$f(1)$	$f(2)$	$f(3)$	$f(4)$	$f(5)$	$f(6)$	$f(7)$	$f(8)$
1	1	10	16	5	6	14	7	3
2	1	11	15	2	10	13	4	6
3	14	1	12	10	5	15	7	16
4	16	1	10	14	3	11	13	12

According to Theorem 2.2, the possibilities for Magic Octagon are represented by a 8-permutation of the set $\{1, 2, 3 \dots 16, 17\}$.

The table below show the quantities for all Magic Polygons up to 24 sides. According Theorem 4.1 we can obtain infinities Magic Polygons, but they shall be just relabelling of these built with the sequence $\{1, 2, 3 \dots 2n+1\}$.

The column $f(1) = 1$ show all possibilities when 1 is a vertex, and the column $f(2) = 1$ show all possibilities when 1 is a edge middle point.

n	$f(1) = 1$	$f(2) = 1$	#	n	$f(1) = 1$	$f(2) = 1$	#
6	2	2	4	16	2.034	2.950	4.984
8	2	2	4	18	20.056	25.218	45.274
10	4	8	12	20	203.265	257.563	460.828
12	87	79	166	22	1.908.120	2.585.799	4.493.919
14	375	487	862	24	22.062.915	29.268.446	51.331.361

These programs has been executed on a Personal Computer in the Operational System *Debian GNU/Linux 11* with a 7.6 GB of RAM and the processor Intel Core *i5-6200U 2.3 GHz Quad Core*.

6 Conclusion

Basics but no trivials optimizations has been used in order to improve the algorithm. However, further properties of the Magic Polygons can be helpful to create algorithm with even better performance. By introducing the Magic Polygons as elements of the *symmetric groups*, some possibilities appears, like the aid of the concepts in group group theory and others algebraic structures to understand better such objects. However, such approach is still in an embryonic stage.

Competing Interests

Author has declared that no competing interests exist.

References

- [1] Chan CYJ, Mainkar MG, Narayan SK, Webster TD. A construction of regular magic squares of odd order, *Linear Algebra and its Applications*. 2014; 457: 293–302
- [2] Kim Y, Yoo J. An algorithm for constructing magic squares, *Discrete Applied Mathematics*. 2008; 156: 2804–2809.
- [3] Mattingly RB. Even order regular magic squares are singular, *Amer. Math. Monthly*. 2000; 107: 777–782.
- [4] Nordgren RP. New constructions for special magic squares, *Int. J. Pure Appl. Math*. 2012; 78.
- [5] Ollerenshaw K, Brée DS. *Most-Perfect Pandiagonal Magic Squares: Their Construction and Enumeration*, The Institute of Mathematics and its Applications, Southend-on-Sea, UK. 1998.
- [6] Planck C. Pandiagonal magic squares of orders 6 and 10 without minimal numbers, *Monist*. 1919; 29: 307–316.
- [7] Chu KL, Drury SW, Styan GPH, Trenkler G. Magic Moore–Penrose inverses and philatelic magic square with special emphasis on the Daniels–Zlobec magic square, *Croatian Oper*. 2011;Res. Rev. 2: 4–13.
- [8] Ganapathy G, Mani K. Add-on security model for public-key cryptosystem based on magic square implementation, in: *Proc*.

- [9] Loly PD. Franklin squares: a chapter in the scientific studies of magical squares, Complex Systems. 2007; 17: 143–161. World Congress on Engineering and Computer Science 1 WCECS; 2009.
- [10] Augusto D. D, Rocha, J. S, Magic Polygons and Degenerated Magic Polygons: Characterization and Properties; Asian Research Journal of Mathematics, Vol. 14;
- [11] Jakicic V, Bouchat R. Magic Polygons and their properties; 2018.
Available: <https://arxiv.org/abs/1801.02262arXiv:1801.02262v1>.
- [12] Stewart, Ian. Incredible Numbers; Profile Books Ltd, 2015.
- [13] Cammann S. The evolution of magic squares in China, J. Am. Oriental Soc. 1960; 80: 116–124.
- [14] Pickover CA. The Zen of Magic Squares, Circles, and Stars, second printing and first paperback printing, Princeton University Press, Princeton, NJ; 2003.
- [15] Andrescu T, Andrica D, Cucurezeanu I. Some Classical Diophantine Equations, First Online, Birkhäuser Boston; 2010.
- [16] Andress WR. Basic properties of pandiagonal magic squares, Amer. Math. Monthly. 1960; 67: 143–152.
- [17] Rosser B, Walker RJ. The algebraic theory of diabolic magic squares, Duke Math. J. 1939; 5: 705–728.

© 2022 Augusto; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/91120>