



Growth, Yield and Grain Arsenic Concentration of Rice Cultivars under Varying Levels of Soil Arsenic Contamination

M. Rafiqul Islam ^a, Samina Nasrin Borna ^b, M. R. K. Menon ^a
and Shofiqul Islam ^{a*}

^a Department of Soil Science, Bangladesh Agricultural University, Mymensingh - 2202, Bangladesh.

^b Department of Environmental Science, Bangladesh Agricultural University, Mymensingh - 2202, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJSSPN/2022/v8i330141

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91631>

Original Research Article

Received 15 June 2022
Accepted 09 August 2022
Published 14 September 2022

ABSTRACT

Arsenic (As) contamination in groundwater is a severe and widespread problem in Bangladesh and their exposure cause a serious health hazard in human history. A pot experiment was conducted in the net-house of the Department of Soil Science, Bangladesh Agricultural University, Mymensingh using fourteen cultivars of rice (Summer rice) to evaluate the effects of varying levels of As contamination on growth, yield and grain As contents. Arsenic contamination adversely affected tillering, filled grains, grain yield and straw yield of all rice cultivars, however, the effect varied among the cultivars. In As control treatment, the highest grain yield (75.66 g pot⁻¹) was observed in BR 11 while the lowest grain yield (8.73 g pot⁻¹) was in BRR1 dhan4. Application of 20 ppm As to the soil resulted in complete death of BR 11, BRR1 dhan30, BRR1 dhan33, BRR1 dhan34, BRR1 dhan41, Binadhan-4, Biroy and Kalizeera cultivars. Arsenic contamination significantly increased As concentration in grain for all cultivars of rice. Arsenic concentration on rice grains grown in soils without As contamination ranged from 0.06 ppm in BR 11 to 0.13 ppm in BRR1 dhan32 and BRR1 dhan41, while such values ranged from 0.231 ppm in Kalizeera with 10 ppm As to 0.743 ppm in BRR1 dhan40 with applied 20 ppm As to soils. Among the 14 rice cultivars BRR1 dhan32, BR 11 and Najirshail appeared to be more tolerant to As contamination.

Keywords: Rice; yield; arsenic contamination; grain arsenic.

*Corresponding author: Email: sislam_ss@bau.edu.bd;

1. INTRODUCTION

Arsenic (As) is an important geogenic contaminant found ubiquitously in the earth threatening the health of 150 million people worldwide [1-2]. The major sources of As exposure to humans includes drinking water and different food items, especially rice. Exposure of As is associated with various chronic and acute health problems to humans that include skin lesions, cardiovascular diseases, diabetes, cancer, and so on [2]. The environmental pollutant As is bound to Fe hydroxides, oxyhydroxides, and oxides that released into the groundwater through biogeochemical processes and resulted in widespread As contamination in Bengal delta [3-4]. Geogenic As contaminated groundwater are being used for rice cultivation in West Bengal and Bangladesh. This common practice has resulted in As build-up in soil through the years [5-6]. Background As concentration in Bangladesh soils is in the range of 4-8 mg as kg^{-1} . Research studies indicate that, long-term use of As contaminated groundwater for irrigation purposes may result in elevated As concentration in soils and plants, and may lead to crop yield loss and elevated As concentration in cereals, vegetables and other agricultural products [7-9]. Rice can accumulate several-fold higher As than other cereal crops such as wheat and maize [2,6,10]. Every year approximately, 36.60 million tons of rice is produced from 11.42 million hectares of land of Bangladesh [11]. The As accumulation in rice grains varies significantly in different rice varieties [12]. The concentrations of As in different parts of rice varied significantly with the contamination level in soil. However, the identification of a suitable low As accumulating rice variety and its use in breeding programs for the development of a suitable rice variety is now time demanding. Under these circumstances, a successful solution can be to identify rice varieties that accumulate low As in grains. suitable rice variety is now time demanding. Under these circumstances, a successful solution can be to identify rice varieties that accumulate low as in grains.

2. MATERIALS AND METHODS

The experiment was conducted at the net-house of the Department of Soil Science of Bangladesh Agricultural University, Mymensingh, during the Aman rice season (Summer rice) in 2018. Geographically the net-house stands at 24.750 N latitude and 90.500 E longitude at the height of 18 m above the sea level. The rice cultivars

tested were ten high yielding varieties [BR 11, BRRI dhan30, BRRI dhan31, BRRI dhan32, BRRI dhan33, BRRI dhan34, BRRI dhan39, BRRI dhan40, BRRI dhan41, Binadhan-4], and four local varieties [Najirshail, Biroy, Kalizeera, Pajam] were collected from Genetic Resources and Seed Division, BRRI, Gazipur. The soil for the experiment was silt loam in texture, pH 6.7, organic matter 2.0%, total N 0.12%, available P 9.8 ppm, exchangeable K 0.11 me%, available S 10.5 ppm and total As 2.1 ppm. Three levels of as contamination was induced in soil by adding viz. 0, 10 and 20 ppm and tested on 14 rice cultivars (10 modern and 4 local cultivars). The trial was set in a Completely Randomized Design with three replications. A total of 126 plastic pots were used in the experiment. Each plastic pot had the diameter of 43 cm and height 40 cm receiving 10 of kg soil. Thirty-five days old seedlings were transplanted in the pots and three hills were placed in an equal distance and each hill consisted of 3 seedlings. Urea, Triple Super Phosphate (TSP), Muriate of Potash (MoP), Gypsum and Zinc oxide were applied @ 100 mg kg^{-1} N, 25 mg kg^{-1} P, 50 mg kg^{-1} K, 25 mg kg^{-1} S and 5 mg kg^{-1} Zn, respectively. The entire TSP, MoP, Gypsum and zinc were applied during the final pot preparation. Urea was applied in three equal splits; at 7, 30 and 60 days after transplanting. Intercultural operations were employed as and when necessary. Data were recorded on plant height, effective tillers pot^{-1} , panicle length, filled grains panicle⁻¹, 1000-grain weight, grain yield pot^{-1} and straw yield pot^{-1} . The grain As content was determined after digesting the samples with HNO_3 and H_2O_2 at 120°C followed by flow injection hydride generation atomic absorption spectrophotometer with UNICAM model No. 969 with hydride generator assembly using matrix-matched standards. The analysis of variance for various characters of the crop was done following the MSTATC computer program. The mean comparison of the treatments was made by the Duncan's Multiple Range Test (DMRT).

3. RESULTS AND DISCUSSION

3.1 Effective Tillers

The effective tillers pot^{-1} decreased significantly due to as contamination to soil. This reduction was significant for all the varieties. The number of effective tillers pot^{-1} in As control condition was 19.31 and that in 10 and 20 ppm As was 8.10 and 1.57, respectively (Table 1). The high yielding and

Table 1. Effects of added As and varieties on the yield contributing characters, grain and straw yield and grain As concentration of different rice cultivars

Factors	Number of effective tillers pot ⁻¹	Number of filled grains pot ⁻¹	Grain yield (g pot ⁻¹)	Straw yield (g pot ⁻¹)	Grain As (ppm)
As added (ppm)					
0	19.31a	89.54a	42.27a	62.54a	0.10b
10	8.10b	50.12b	11.30b	19.47b	0.45a
20	1.33c	8.75c	2.12b	3.80b	0.64a
S.E. (±)	0.41	3.52	2.73	3.04	0.05
Cultivars					
BR 11	14.22b	48.31b	33.57a	39.74ab	0.31cd
BRR1 dhan30	7.44dc	57.82ab	17.58cd	31.39b	0.26c
BRR1 dhan31	10.45bc	67.33a	24.90b	29.09bcd	0.40bc
BRR1 dhan32	15.01ab	58.71ab	12.00d	34.39b	0.34bcd
BRR1 dhan33	9.00cd	35.95c	19.27cd	23.43d	0.31d
BRR1 dhan34	5.22d	49.40b	8.28d	14.42d	0.22e
BRR1 dhan39	10.00bc	36.00c	12.33d	16.69d	0.48ab
BRR1 dhan40	5.34d	54.86b	20.02b	28.81b	0.49a
BRR1 dhan41	1.44e	24.41d	2.91e	16.64d	0.13e
Binadhan-4	10.00c	48.86b	23.93b	27.21bcd	0.35bcd
Najirshail	19.11a	68.38a	32.55a	45.43a	0.35bcd
Biroy	10.00bc	37.05c	18.55c	31.92b	0.23d
Kalizeera	10.56bc	54.84b	19.56c	28.75b	0.15e
Pajam	7.45dc	50.59b	14.40d	32.55b	0.38bcd
S.E. (±)	1.21	3.78	1.52	2.07	0.05

In a column figures having same letters do not differ significantly at 5% level by DMRT

Table 2. Interaction effects of added As and varieties on the yield contributing characters of different rice cultivars

Cultivars	Number of effective tillers pot ⁻¹			Number of filled grains pot ⁻¹		
	As ₀	As ₁₀	As ₂₀	As ₀	As ₁₀	As ₂₀
BR 11	28.33ab	14.33d-g	0.00m	93.03b-d	51.91j-n	0.00r
BRR1 dhan30	13.66d-h	8.67g-m	0.00m	103.57ab	69.88g-j	0.00r
BRR1 dhan31	20.67b-d	7.00g-m	3.67i-m	94.07b-d	81.85c-h	26.08o-q
BRR1 dhan32	30.37a	12.00e-i	2.67j-m	102.29ab	61.12i-l	12.73qr
BRR1 dhan33	20.67b-d	6.33g-m	0.00m	70.66f-j	37.20m-o	0.00r
BRR1 dhan34	10.33e-j	5.33h-m	0.00m	91.81b-e	56.40i-m	0.00r
BRR1 dhan39	18.00c-e	8.00g-m	4.00i-m	58.48i-l	33.56n-p	15.97p-r
BRR1 dhan40	12.00e-i	3.33i-m	0.68lm	114.31a	37.39m-o	12.89qr
BRR1 dhan41	4.33i-m	0.00m	0.00m	73.24d-i	0.00r	0.00r
Binadhan-4	20.68b-d	9.33f-l	0.00m	84.46b-g	62.13h-k	0.00r
Najirshail	30.33a	17.33c	9.67e-k	91.11b-f	70.52f-g	43.51k-o
Biroy	23.33a-c	6.68i-m	0.00m	84.56b-g	26.58o-q	0.00r
Kalizeera	24.00a-c	7.68g-m	0.00m	93.03a-c	71.50e-j	0.00r
Pajam	13.68b-h	7.33g-m	1.33k-m	98.88a-c	41.57l-o	11.33qr
S.E. (±)		1.501			3.658	

In a column figures having same letters do not differ significantly at 5% level by DMRT

local varieties selected for the experiment showed significant variation in terms of number of effective tillers pot⁻¹ (Table 1). The highest number of effective tillers pot⁻¹ was obtained in Najirshail with the value of 19.11 which was statistically similar to BRR1 dhan32. The interaction of As levels and different high yielding and local varieties selected for the experiment showed remarkable variation in terms of number of effective tillers pot⁻¹ (Table 2). Application of

10 and 20 ppm as significantly reduced the number of effective tillers pot⁻¹. The highest number of effective tillers pot⁻¹ (30.37) was found in BRR1 dhan32 with 0 ppm as application. The cultivar BRR1 dhan41 failed to produce any effective tillers due to the application of 10 and 20 ppm As and the other seven rice varieties BR 11, BRR1 dhan30, BRR1 dhan33, BRR1 dhan34, Binadhan-4, Biroy and Kalizeera failed to produce plants in 20 ppm As treatment. The

corresponding reductions in tiller numbers also reported in Aman rice ranged from 0–22 % [13]. Similarly, a significant effect on plant height and effective tiller numbers has been observed when a different variety of rice is cultivated in soil artificially enriched with arsenic acid [14].

3.2 Filled Grains

The filled grains pot^{-1} was decreased significantly with addition of As to soil. The number of filled grains pot^{-1} recorded in 0 ppm As was 89.54 and that in 10 and 20 ppm As was 50.12 and 8.75, respectively (Table 1). Higher As concentration causes grain sterility and grains panicle^{-1} were also significantly affected by the As treatments [13,15]. All the rice cultivars showed significant variation in terms of number of filled grains pot^{-1} (Table 1). The highest number of filled grains pot^{-1} was obtained in Najirshail with the value of 68.38 which was statistically similar to BRR1 dhan31 and BRR1 dhan30. The interaction of As levels and different high yielding and local varieties selected for the experiment showed remarkable variation in terms of number of filled grains pot^{-1} (Table 2). Application of 10 and 20 ppm As significantly reduced the number of filled grains pot^{-1} . The highest number of filled grains pot^{-1} (114.31) was found in BRR1 dhan40 with 0 ppm As application.

3.3 Grain Yield

Grain yield decreased significantly with levels of As application. Grain yield was considered as one of the most important parameters in variety selection in terms of As tolerance or sensitivity. The grain yield of rice cultivars was statistically affected by As contamination (Table 1). The application of 10 and 20 ppm As in soil produced 11.30 g and 2.12 g grain pot^{-1} , respectively, whereas in control (0 ppm As) the grain was 42.27 g pot^{-1} i.e. yield reduction was almost 73.31% and 94.98 %, respectively over control. Previously, decreased of rough rice yield from about 7 to 2 t/ha in 2006 and 9 to 3 t/ha in 2007 across the soil As gradient has been reported [9]. A number of authors [16,17] also reported yield reduction as a result of higher levels of As concentrations. Generally higher levels of As addition resulted in greater reduction in yield and the application of As to soil resulted in 80% yield reductions in Aman rice (BRR1 dhan33), indicating a strong residual effect of added As on the next crop [13]. In winter and summer rice, above ground biomass of rice genotypes progressively decreased because of soil As

gradient from 0 to 60 mg/kg [18]. The grain yield of rice varied significantly in different cultivars of Aman rice (Table 3). The highest grain yield (33.57 g pot^{-1}) was found in BR 11 which was statistically similar to Najirshail. On the other hand, the lowest grain yield (2.91 g pot^{-1}) was found in BRR1 dhan41. Almost similar result represents by another study in Bangladesh [17]. The yield markedly declined at 20 ppm As treatment and above. The grain yield was reduced to 93.1% for 50 ppm As treatment, whereas its reduction was only 1.8% in 1 ppm As treatment compared to the control. The interaction effect of added As and different rice varieties selected for the experiment significantly affected the grain yield (Table 3). The application of 10 and 20 ppm As reduced the grain yield of all cultivars of rice selected for this experiment. In As control treatment, the highest grain yield (75.66 g pot^{-1}) observed in BR 11. The lowest grain yield (8.73 g pot^{-1}) in As control treatment was found in BRR1 dhan41. The addition of 10 ppm As significantly reduced the grain yield of rice compared to that obtained in As control treatment. The application of 20 ppm As significantly decreased the yield and most of the varieties failed to produce grain yield except BRR1 dhan31, BRR1 dhan32, BRR1 dhan39, BRR1 dhan40 and Najirshail. Thus, it can be said that in respect to grain yield Najirshail is the most tolerant variety to As contamination.

3.4 Straw Yield

Straw yield of rice cultivars was statistically affected by As contamination (Table 1). The yield due to As contamination varied from 62.54 g pot^{-1} in control (0 ppm As) to 19.47 g and 3.80 g pot^{-1} in 10 and 20 ppm As treatment, respectively. Application of 10 and 20 ppm As to soil caused 64.87 % and 93.93 % straw yield reduction of rice, respectively. Similarly straw yield was reduced by 7–64 % in the rice [13]. The straw yield of rice ranged from 14.42 g pot^{-1} in BRR1 dhan34 to 45.43 g pot^{-1} in Najirshail (Table 1). The interaction effect of different rates of As and cultivars of rice was significant in producing the straw yield (Table 3). The application of 10 ppm As decreased the straw of all rice varieties but the degree of decrease in straw yield was quite variable. The application of 20 ppm As significantly decreased the yield and some of the varieties failed to produce straw yield viz. BRR1 dhan33, BRR1 dhan34, BRR1 dhan41 and Binadhan-4. Considerable reduction in straw biomass in the higher As treatment recorded in this experiment primarily as a result of lower

Table 3. Interaction effects of added As and cultivars on grain yield, straw yield and grain As concentration of different rice cultivars

Cultivars	Grain yield pot ⁻¹ (g)			Straw yield pot ⁻¹ (g)			Grain As (ppm)		
	As ₀	As ₁₀	As ₂₀	As ₀	As ₁₀	As ₂₀	As ₀	As ₁₀	As ₂₀
BR 11	75.66a	25.05f-h	0.00k	83.57a	32.79gh	2.87m-o	0.063e	0.559ab	-
BRR1 dhan30	39.40de	13.35g-k	0.00k	66.62b-e	25.99hi	1.57no	0.095e	0.417a-d	-
BRR1 dhan31	59.14bc	10.60g-k	4.95i-k	66.64b-e	14.06i-o	6.57k-o	0.105de	0.458a-c	0.625ab
BRR1 dhan32	17.25ab	17.26g-k	1.48i-k	80.08a-c	19.50h-m	3.58m-o	0.133d-e	0.333b-e	0.550ab
BRR1 dhan33	55.21bc	2.61i-k	0.00k	63.97c-e	6.31k-o	0.00o	0.097e	0.525a-c	-
BRR1 dhan34	17.54g-i	7.29i-k	0.00k	27.52g-i	15.75i-o	0.00o	0.108de	0.333b-e	-
BRR1 dhan39	26.19fg	9.40h-k	1.39i-k	32.76gh	13.49i-o	3.81l-o	0.108de	0.642ab	0.683ab
BRR1 dhan40	47.06de	10.01g-k	2.98i-k	61.54de	21.30h-k	3.59m-o	0.104de	0.625ab	0.743a
BRR1 dhan41	8.73h-k	0.00k	0.00k	42.40fg	7.53j-o	0.00o	0.128de	-	-
Binadhan-4	56.90bc	14.90g-k	0.00k	63.52de	18.12h-n	0.00o	0.116de	0.583ab	-
Najirshail	52.78cd	26.49fg	18.39g-i	81.46ab	34.18gh	20.65h-l	0.082e	0.333b-e	0.620ab
Biroy	49.24c-e	6.42i-k	0.00k	68.29a-e	23.78h-j	3.68m-o	0.075e	0.375a-e	-
Kalizeera	52.06cd	6.63i-k	0.00k	64.94c-e	21.31h-k	0.00o	0.071e	0.231c-e	-
Pajam	34.58ef	8.18i-k	0.43jk	72.25a-d	18.46h-n	6.93k-o	0.078e	0.450a-c	0.600ab
S.E. (±)		2.845			2.864			0.053	

In a column figures having same letters do not differ significantly at 5% level by DMRT

tillers pot⁻¹ and stunted rice plant. The same type of declining As uptake pattern for straw is found for one variety of rice grown in a glasshouse [19]. Exposure to As, causing a reduction in shoot bio-mass and/ or growth, was also reported by many research findings [20-21].

3.5 Grain Arsenic

Arsenic concentration is an important consideration for the selection of As tolerant rice genotypes. Arsenic concentration in grains of rice cultivars was significantly increased due to application of different levels of As in soil (Table 1). The rate of increase was significant in all of the cultivars. Arsenic concentration in grain of control pot (0 ppm As) was 0.10 ppm and that in pot receiving 10 and 20 ppm As was 0.45 and 0.64 ppm, respectively. The As concentrations in different cultivars ranged from 0.13 ppm in BRR1 dhan41 to 0.49 ppm in BRR1 dhan40. There was a significant interaction effect of different rates of As and cultivars of rice on As concentration of rice grains (Table 3). The application of 10 and 20 ppm As increased the grain As concentration of rice. All the varieties selected for the experiment showed higher As concentration in the pots receiving 10 and 20 ppm As in soil compared to As control pot. Among the treatment combinations As₂₀ × BRR1 dhan40 had the highest As concentration (0.743 ppm) in grain and followed by As₂₀ × BRR1 dhan39 and As₂₀ × BRR1 dhan31. The lowest As concentration (0.063 ppm) was found in As₀ × BR 11 treatment combination. Cultivation of rice in soils with elevated concentration has been reported a number of authors [22-23]. The As concentrations in rice grain vary widely depending on the cultivars, As status of soil and irrigation water [8,16]. The grain As concentration increased from values of 0.22–0.32 ppm with the As₀ treatment to 0.59–0.81 ppm with the As₂₀ treatment, but varied inconsistently between the varieties.

4. CONCLUSION

From this experiment, it was found that high level of As contamination to soil significantly affected the yield contributing characters viz. number of effective tillers number of filled grains of rice cultivars. The grain and straw yields of these rice cultivars were also markedly reduced by the artificial introduction of As into soil. A high As uptake in rice grain and rice straw was observed in this experiment. Rice cultivation in the naturally As-contaminated soil of Bangladesh may also reduce the yield of rice grain and straw.

Among the fourteen rice cultivars, BRR1 dhan32, BR 11 and Najirshail appeared to be tolerant to As contamination. Therefore, at high levels of As contamination, the varieties viz. BRR1 dhan32, BR 11 and Najirshail may be considered for cultivation. However, further studies with varying levels of As contamination with different varieties in the field are needed to reach a final conclusion. Therefore, choosing less As-responsive rice genotypes to grow in As affected soil could effectively minimize As bioaccumulation in rice grains, leading to a reduction in health risks caused by As exposure.

ACKNOWLEDGEMENTS

Author would like to express sincere appreciation and gratefulness to the Department of Soil Science, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh for providing necessary supports to govern this research findings.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Majumder S, Banik P. Geographical variation of arsenic distribution in paddy soil, rice and rice-based products: A meta-analytic approach and implications to human health. *J Environ Manage.* 2019; 233:184-99.
2. Shikawa S, Arao T, Makino T. Agronomic strategies for reducing arsenic risk in rice. In: *Arsenic Contamination in Asia* Yamauchi H, Sun G, editors. Singapore: Springer. 2019;181-98.
3. Majumdar A, Bose S. Toxicogenesis and metabolism of arsenic in rice and wheat plants with probable mitigation strategies. In: *Arsenic: Risks of Exposure, Behavior in the Environment and Toxicology* Knežević R, editor. Hauppauge, NY: Nova Science Publishers. 2017;149-66.
4. Raessler M. The arsenic contamination of drinking and groundwaters in Bangladesh: featuring biogeochemical aspects and implications on public health. *Arch Environ Contam Toxicol.* 2018;75(1): 1-7.
5. Panauallah GM, Alam T, Duxbury JM, Loeppert RH, Meisner CA, Lauren JG et al. Arsenic in soil and its effect on the growth

- of rice at a high-arsenic site in Bangladesh. The 18th World Congress of Soil Science. Philadelphia; July 9-15. 2006;2006.
6. Upadhyay MK, Shukla A, Yadav P, Srivastava S. A review of arsenic in crops, vegetables, animals and food products. Food Chem. 2019;276:608-18.
 7. Hossain MB, Jahiruddin M, Loeppert RH, Panaullah GM, Islam MR, Duxbury JM. The effects of iron plaque and phosphorus on yield and arsenic accumulation in rice. Plant Soil. 2009;317(1-2):167-76.
 8. Islam S, Rahman MM, Islam MR, Naidu R. Effect of irrigation and genotypes towards reduction in arsenic load in rice. Sci Total Environ. 2017;609:311-8.
 9. Panaullah GM, Alam T, Hossain MB, Loeppert RH, Lauren JG, Meisner CA et al. Arsenic toxicity to rice (*Oryza sativa* L.) in Bangladesh. Plant Soil. 2009;317(1-2): 31-9.
 10. Williams PN, Raab A, Feldmann J, Meharg AA. High levels of arsenic in South Central, US rice grain: consequences for human dietary exposure. Environ Sci Technol. 2007;41(7):2178-83.
 11. BBS (Bangladesh Bureau of Statistics). Yearbook of agricultural statistics of Bangladesh. Bangladesh Bureau of Statistics. Government of Bangladesh; 2020.
 12. Norton GJ, Duan G, Dasgupta T, Islam MR, Lei M, Zhu Y et al. Meharg. Environmental and genetic control of arsenic accumulation and speciation in rice grain: comparing a range of common cultivars grown in contaminated sites across Bangladesh, China, and India. Environ Sci Technol. 2009;43(21):8381-6.
 13. Khan MA, Islam MR, Panaullah GM, Duxbury JM, Jahiruddin M, Loeppert RH. Accumulation of arsenic in soil and rice under wetland condition in Bangladesh. Plant Soil. 2010;333(1-2):263-74.
 14. Wang F-M, Chen Z-L, Zhang L, Gao Y-L, Sun Y-X. Arsenic uptake and accumulation in rice (*Oryza sativa* L.) at different growth stages following soil incorporation of Roxarsone and arsanilic acid. Plant Soil. 2006;285(1-2):359-67.
 15. Islam S, Islam MR, Jahiruddin M, Islam MM. Genotypic variation for yield, yield attributes and grain arsenic content of boro rice grown in arsenic contaminated soils. Bangladesh J Crop Sci. 2008;19(20): 183-90.
 16. Rahman MA, Hasegawa H, Rahman MM, Rahman MA, Miah MA. Accumulation of arsenic in tissues of rice plant (*Oryza sativa* L.) and its distribution in fractions of rice grain. Chemosphere. 2007;69(6):942-8.
 17. Azad MAK, Islam MN, Alam A, Mahmud H, Islam MA, Rezaul Karim M et al. Arsenic uptake and phytotoxicity of T-aman rice (*Oryza sativa* L.) grown in the As-amended soil of Bangladesh. Environmentalist. 2009;29(4):436-40.
 18. Iqbal M, Rahman GKMM, Panaullah GM, Kabir H, Biswas JC. Influence of soil arsenic levels on biomass production and relationship the concentration of arsenic between Rice straw and grain. AJSSPN. 2019;5(2):1-10.
 19. Azizur Rahman MA, Hasegawa H, Mahfuzur Rahman M, Mazid Miah MA, Tasmin A. Arsenic accumulation in rice (*Oryza sativa* L.): human exposure through food chain. Ecotoxicol Environ Saf. 2008;69(2):317-24.
 20. Huq SM, Shila UK, Joardar JC. Arsenic mitigation strategy for rice using water regime management. Land Contam Reclam. 2006;14(4):805-13.
 21. Islam MR, Islam S, Jahiruddin M. and Islam MA. Effects of irrigation water arsenic in the rice-rice cropping system. Journal of Biological Science. 2004; 4: 542-546.
 22. Williams PN, Islam MR, Adomako EE, Raab A, Hossain SA, Zhu YG et al. Increase in rice grain As for regions of Bangladesh irrigating paddies with elevated arsenic in ground waters. Environ Sci Technol. 2006;40(16): 4903-8.
 23. Habiba KK, Jahiruddin M, Islam MR, Islam S. Phosphorus increases arsenic toxicity in rice. Bangladesh J Seed Sci Technol. 2007;11(1&2):31-6.

© 2022 Islam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91631>