

Spontaneous Rupture of a Huge Subserous Uterine Fibroid Vessel with Massive Hemoperitoneum: A Rare Case of Fibroid

Awopola Ibiebelem Jumbo^a and Esther Ijeoma Nonye-Enyidah^{a*}

^a Department of Obstetrics and Gynaecology, Rivers State University Teaching Hospital, Port Harcourt, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors managed the patient. Author AIJ wrote the first draft of the manuscript while author EINE did the literature research. Both authors read and approved the final manuscript.

Article Information

Editor(s):

- (1) Dr. Arun Singh, Rohilkhand Medical College & Hospital Bareilly, India.
- (2) Dr. Hab. Mariusz Cycon, Medical University of Silesia, Poland.

Reviewers:

- (1) Hemant Deshpande, India.
- (2) Umashankar Km, MVJ Medical College, India.
- (3) N Nagendra Prasad, Dayananda Sagar University, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here: <https://www.sdiarticle5.com/review-history/79314>

Case Report

Received 20 October 2021
Accepted 29 December 2021
Published 04 January 2022

ABSTRACT

Background: Spontaneous rupture is a potentially life-threatening acute complication of uterine fibroid. It is a very rare clinical diagnosis and hardly presents with massive hemoperitoneum. This is a case of spontaneous rupture of uterine fibroid vessels with massive hemoperitoneum, necessitating prompt surgical intervention and massive transfusion.

Case Report: Mrs A. I was a 35-year-old Para 1 who presented to the accident and emergency department of the Rivers State University Teaching Hospital (RSUTH) on the 14th October 2021 with a three-month history of amenorrhoea, and 10 hours history of severe lower abdominal pains associated with dizziness. She was in shock and also had abdominal distension and tenderness. Ultrasonography showed massive hemoperitoneum, uterine fibroids and a left adnexal mass. She had an emergency exploratory laparotomy and the findings were massive hemoperitoneum, a left ovarian cyst and a huge subserous uterine fibroid with an actively bleeding surface vessel. Myomectomy and left ovarian cystectomy were done. She received 5 units of blood perioperatively and had an uneventful post-operative recovery.

Conclusion: Spontaneous rupture of uterine fibroid is a very rare clinical condition. It can lead to significant morbidity and mortality if not recognised and managed promptly. Therefore a high index of suspicion is necessary for women with huge uterine fibroids and hemoperitoneum.

Keywords: Spontaneous rupture; subserous fibroid; hemoperitoneum; laparotomy; myomectomy.

1. INTRODUCTION

Uterine fibroids are the commonest benign pelvic tumour in women [1,2]. They are often asymptomatic and rarely associated with life-threatening complications [2,3]. Acute complications of a uterine fibroid are uncommon and may require immediate surgical treatment [4]. The most dreaded acute complications are acute abdomen due to fibroid torsion or intra-abdominal haemorrhage from rupture of the fibroid [4]. Fibroid rupture is mostly due to trauma [2]. However, Spontaneous rupture of uterine fibroid is very rare and only about 100 cases so far have been reported in the literature [4,5]. The exact mechanism of spontaneous fibroid rupture is poorly understood [5]. The diagnosis is difficult pre-operatively due to the rarity of the condition [6]. The occurrence of massive intra-abdominal haemorrhage due to spontaneous rupture of uterine fibroid is extremely rare [4,7]. This is a case of rupture of a uterine fibroid with massive hemoperitoneum necessitating emergency laparotomy, myomectomy and massive transfusion.

2. CASE REPORT

Mrs A. I was a 35-year-old Para 1 (alive) who presented to the accident and emergency department of the Rivers State University Teaching Hospital on the 14th of October, 2021. She presented with severe lower abdominal pain of 10 hours duration, associated with weakness and dizziness. There was no vaginal discharge, vaginal bleeding or trauma. She was amenorrhagic for three months following the removal of a subdermal implant.

On examination, she was conscious and alert, dyspnoeic and severely pale. The pulse was 126 beats/minute and the blood pressure was 80/50mmHg. Her abdomen was distended, and there was generalised tenderness and guarding. Intraperitoneal fluid was also demonstrated. Her packed cell volume was 18%. An abdominopelvic ultrasound scan that was done before the presentation was suggestive of ruptured ectopic pregnancy with massive hemoperitoneum extending into the Morrison's pouch. The

ultrasound scan also showed a 14cm x 12cm sized uterine fibroid and a complex echogenic mass in the left adnexa. A diagnosis of ruptured ectopic pregnancy with massive hemoperitoneum and a huge uterine fibroid was made. She was resuscitated with two litres of intravenous normal saline, a unit of blood and intranasal oxygen. The diagnosis and management were explained to her and consent was obtained for surgery. She subsequently had an emergency exploratory laparotomy, and the findings were hemoperitoneum of 3.2 Litres, a left ovarian cyst about 5 cm x 5 cm and a 20 weeks size uterus harbouring a fundal subserous fibroid. The fibroid measured 16 cm x12 cm x10 cm and had a surface vessel that was ruptured at two points (about 2 cm apart) and was actively bleeding. The vessel was ligated, then myomectomy and left ovarian cystectomy were done. She was transfused with 2 units of blood intraoperatively. She received antibiotics, analgesics and haematinics postoperatively, as well as another 2 units of blood. She had an uneventful recovery and was discharged on the 5th postoperative day. She was seen two weeks later at the gynaecology clinic and had no complaint. The histology report was consistent with uterine fibroid and endometrioid cyst.

3. DISCUSSION

Uterus fibroid is one of the commonest benign tumours of the female genital tract. It is often asymptomatic, when symptomatic it may present with abnormal uterine bleeding, abdominal swelling, pelvic pressure or subfertility [8].

Acute complications of uterine fibroids are uncommon [4]. Intra-abdominal haemorrhage from rupture of uterine fibroid is a potentially life-threatening complication. It is very rare, and about 100 cases have so far been documented in the literature [4,5]. It usually results from rupture of the surface fibroid veins as in this case, rarely arterial bleeds can occur due to hypertension [6,7].

The exact mechanism that initiates the rupture of the surface vessels is poorly understood [5,6]. Common risk factors include large fibroid size




Fig. 1. Ruptured uterine fibroid with massive hemoperitoneum

(>10cm), trauma, torsion, alcohol consumption, and venous congestion (due to pregnancy, weight-bearing, defaecation, or strenuous exercise) [6]. Mrs A. I had a huge fundal uterine fibroid.

Spontaneous rupture of uterine fibroid is a very rare phenomenon [5,7]. It is often mistaken for other common pathologies and diagnosed incidentally at surgery as was done in this case [6,9]. Diagnosis of ruptured fibroid is difficult preoperatively owing to the rarity of the condition [6,10], it could be suspected from a history of abdominal pain and swelling with radiologic findings of hemoperitoneum and uterine fibroid [6]. Although ultrasonography is insensitive in cases of small hemoperitoneum and non-specific in locating the source of the hemoperitoneum [2,7], it is a very useful imaging modality to aid the preoperative diagnosis of ruptured fibroid and can be performed at the bedside in unstable patients [2,7]. Magnetic resonance imaging and computed tomography scan can locate the source of the haemoperitoneum but may be difficult to perform in emergencies [7,10]. Preoperative diagnosis has been reported using contrast-enhanced computerized tomography [10-13].

Spontaneous fibroid rupture is a rare cause of massive hemoperitoneum [6,7]. Fewer than 30 cases of massive hemoperitoneum due to uterine fibroid has been reported since 1950 [6]. Common causes of massive hemoperitoneum include ruptured ectopic pregnancy, ruptured ovarian cyst/tumours and ruptured viscus [9]. Mrs A. I presented with massive hemoperitoneum and shock which was misdiagnosed for ruptured ectopic pregnancy.

Management involves prompt evaluation, stabilisation and emergency surgical intervention [7,9]. The surgical options of management are ligation of the bleeding vessels and/or Myomectomy or hysterectomy [7,9]. The decision at surgery depends on the patient age and desire for fertility [14]. Mrs A. I had ligation of the bleeding vessel and myomectomy as she was 35year-old and desires fertility.

4. CONCLUSION

The diagnosis of spontaneous rupture of uterine fibroid can often be challenging as the condition is rare. Therefore Gynaecologist should have a high index of suspicion when hemoperitoneum co-exist with huge uterine fibroids in women of

reproductive age. Since laparotomy is necessary for treatment, adequate preparation for myomectomy or hysterectomy should be made as well.

CONSENT AND ETHICAL APPROVAL

As per university standard guidelines, participant consent and ethical approval have been collected and preserved by the authors.

ACKNOWLEDGEMENT

We wish to express our profound gratitude to everybody who participated in the management of this patient.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Baird DD, Dunson DB, Hill MC, Cousins D, Schectman JM. High cumulative incidence of uterine leiomyoma in black and white women: ultrasound evidence. *Am J Obstet Gynecol.* 2003;188(1):100-7.
2. Rokhgireh S, Kashi AM, Kermansaravi M, Tajbakhsh B, Allahqoli L, Alkatout I, et al. Hemoperitoneum due to bleeding from a vein overlying a subserous uterine myoma: A case report. *J Med Case Rep.* 2020;14(1):1-4.
3. Gupta S, Jose J, Manyonda I. Clinical presentation of fibroids. *Best Pract Res Clin Obstet Gynaecol.* 2008;22(4):615-26.
4. Schwartz M, Powell K. Spontaneous Rupture of a Leiomyoma Causing Life-Threatening Intra-Abdominal Hemorrhage. *Case Rep Obstet Gynecol.* 2017;2017:1-3.
5. Lim WH, Cohen SC, Lamaro VP. Intra-abdominal haemorrhage from uterine fibroids: A systematic review of the literature. *BMC Surg.* 2020;20(1):1-7.
6. Swarray-Deen A, Mensah-Brown SA, Coleman J. Rare complication of fibroids in pregnancy: Spontaneous fibroid rupture. *J Obstet Gynaecol Res.* 2017;43(9):1485-8.
7. Lotterman S. Massive hemoperitoneum resulting from spontaneous rupture of uterine leiomyoma. *Am J Emerg Med.* 2008;26(8):974.e1-974.e2.
8. Okolo S. Incidence, aetiology and epidemiology of uterine fibroids. *Best Pract Res Clin Obstet Gynaecol.* 2008;22(4):571-588.
9. Gulati N, Raman S, Srinivasan M, Bakour S. Case Report: Rare gynaecological emergency: massive intraperitoneal haemorrhage from spontaneous rupture of a superficial vessel on a large leiomyoma. *BMJ Case Rep.* 2016;bcr2015212576.
10. Althobaiti FA, Alsaadi KK, Althobaiti AA. A case of hemoperitoneum due to spontaneous bleeding from a uterine leiomyoma. *Am J Case Rep.* 2019;20:167-70.
11. Daimon A, Tanaka T, Kogata Y, Tanaka Y, Fujita D, Ohmichi M. Hemoperitoneum associated with uterine fibroids: A case report. *Medicine (Baltimore).* 2021;100(10):e24024.
12. Tan J, Siraj M. A Rare Case of Submucosal Fibroid Presented With Acute Abdomen: A Case Report. *J Med Cases.* 2020;11(6):145-147.
13. Alharbi SR. Uterine leiomyoma with spontaneous intraleiomyoma haemorrhage, perforation, and hemoperitoneum in a postmenopausal woman: Computed tomography diagnosis. *Avicenna J Med.* 2013;3(3):81-83.
14. Jayanthi D. Hemoperitoneum due to bleeding myoma in pregnancy. *Univ J Surg Surgical Spec.* 2016;2(7):675-84.

© 2022 Jumbo and Nonye-Enyidah; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79314>