



Role of Family Support, Spirituality and Self-efficacy in Type 2 Diabetes Self-care in Greater Accra Region, Ghana

Elizabeth Oduraa Ampofo ^a, Isaac Nyarko Kwakye ^{b*}, Eliza Oduraa Ampofo ^c and Ernestina Aidoo ^d

^a Department of Psychiatry, Korle-Bu Teaching Hospital, Accra, Ghana.

^b Faculty of Education, West End University College, Accra, Ghana.

^c Social Service Department, Police Hospital, Accra, Ghana.

^d Department of Psychiatry, Korle-Bu Teaching Hospital, Accra, Ghana.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JESBS/2022/v35i730436

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/88897>

Original Research Article

Received 20 April 2022

Accepted 26 June 2022

Published 04 July 2022

ABSTRACT

Aims: The purpose of this study was to examine the relationship between family support, spirituality and self-care in Type 2 Diabetes patients in the Greater Accra Region, Ghana. It also aimed at determining whether self-efficacy would moderate the relationship between spirituality, family support and self-care.

Study Design: A cross sectional study was adopted and 270 diabetes patients living in Greater Accra region were conveniently sampled. The data was collected using questionnaire such as FACIT-sp, DSMQ, SDS and S4-Mad. The IBM SPSS version 2020 was used in analysing the data. The Pearson Product Moment Correlation test and Hierarchical Multiple Regression were used respectively to test for the correlation between variables and moderation role of self-efficacy.

Results: Spirituality correlated with self-care ($r = .151, p < .05$). Also, family support correlated with self-care among T2DM patients ($r = .237, p < .01$). Self-efficacy moderated the relationship between family support and self-care ($\Delta R^2 = .031, \beta = .254, p < .05$). However, self-efficacy did not moderate the relationship between spirituality and self-care ($\Delta R^2 = .002, \beta = -.358, p > .05$).

Conclusion: More spirituality and more social support received by patients are associated with greater self-care. Again, self-efficacy moderates the relationship between family support and self-

care but fails to influence the association between spirituality and self-care. Self-efficacy, spirituality and family support are critical in diabetes self-care. It is therefore beneficial that health care providers incorporate psychosocial interventions into health care delivery.

Keywords: Family support; spirituality; Type 2 diabetes; self-Care.

1. INTRODUCTION

“Type 2 Diabetes (T2DM), also known as Non-Insulin Dependent Diabetes Mellitus, occurs as a result of irregular physiological responses to insulin production and uptake, including insulin resistance at the level of the cell membrane” [1]. According to Walsh and Crumbie [2], “the basic problem is that either there is increased peripheral resistance to the action of insulin or the islets of Langerhans gradually diminish their insulin output or there is a combination of decreased insulin secretion and increased insulin resistance”. “The beginning of T2DM is often slow, and patients could live with the condition for a couple of years before diagnoses. T2DM has psychological, social and health implications and it is the most common type of diabetes which accounts for 90% of diabetes cases” [3]. Accordingly, International Diabetes Federation [IDF] [4] added that T2DM accounts for 85% to 95% of diabetes cases in low-income countries.

“It is a disease that occurs among older adults; however, children, adolescents and young adults have increasingly reported the disease and this is often due to lack of physical activity, poor eating habits and nutrition, obesity and growth” [4]. “Past history of gestational diabetes mellitus (GDM) and the exposure of the unborn child to high blood glucose and smoking have been identified as a significant adjustable risk factor. Furthermore, excessive consumption of sweets and beverages high in sugar content is also an associated risk factor” [5,6,7]. Hörnsten [8] posits that “the cause of T2DM is multifactorial with a substantial genetic component but the disease is also associated with environmental and life style changes. Other factors include inadequate intake of fruits, vegetables, whole grains and dietary fibre, and high intake of energy as saturated fat” [4]. T2DM is characterized by slow healing of wounds, tiredness, repeated fungal infections, numbness in limbs, frequent urination and increased thirst.

“T2DM can result in various complications if not given the needed attention. These complications may be medical, psychological as well as social in nature. The medical complications include

hypoglycaemia, ketoacidosis, nephropathy, neuropathy, diabetic foot ulcers, macrovascular comorbidities, dyslipidaemia, hypertension, metabolic syndrome, retinopathy and blindness” [9]. Also, the psychological complications include depression and anxiety [10]. As a result of the distress, patients are unable to take care of themselves and effective diabetes management behaviours become difficult to achieve and even harder to maintain, which suggests that people with diabetes especially T2DM require urgent and needed support to attain proper self-care.

Self-care is defined by Orem [11] as the practice of activities for the maintenance of life, health and well-being, carried out by the individual for his or her own benefit. However, these activities may be imposed by others such as health care professionals or family members. These behaviours could be related to life-style (smoking cessation, preparing healthy food, and coping with stress) or the medical regimen (taking medication as prescribed) so adaptation is often needed to accommodate changing conditions. Anderson et al. [12] argued that “self-care in diabetes is an important factor to keep the disease under control and about 95% of self-care usually carried out by the affected individual or their families consists of self-monitoring of blood glucose (SMBG), nutrition, physical activity, and compliance to medication”.

“Social support is one of the well-documented psychosocial factors influencing physical health outcomes” [13]. “Studies of social support on chronic disease have found social support essential to self-management” [14,15]. Chio [16] found that “sufficient social support was correlated with high diabetes self-care ability. Social support may come from family, peers and friends, professional or community”. Similarly, Kwakye et al. [17] found that support from family, community members and health care provide a productive force which enhances the psychosocial care of diabetes patients. Family and friends social support (FSS) can either facilitate or threaten Diabetes Self-Management [18]. “Family and friends are often a part of the patient’s everyday life; thus, it is projected that they play a significant role in assisting people

with diabetes and may influence the extent to which T2DM patients adhere to diabetes management” [19]. “Most often social support for persons with chronic disease includes emotional encouragement and instrumental help with monitoring blood glucose, taking medications, foot and eye care, following meal plans, and increasing physical activity” [20,15]. “However, behaviours such as nagging, critical comments and over protection from family members have been found to have negative effects on self-management” [21]. “A recent systematic review found that there are considerable gaps and inconsistencies in research about social support and its effects on diabetes management” [22].

“Additionally, spirituality has also been found to help people cope with a wide range of illnesses or variety of stressful situations. These include people dealing with general medical illness such as diabetes” [23]. In 120 studies conducted on spirituality and well-being, 82% of the studies reported positive relationships between spirituality and well-being [24]. On the contrary, another study reported a negative relationship between spirituality and well-being [25]. Spirituality, although identified as relevant [26], has not been explored in detail globally [27]. The story is not different in Ghana as the phenomenon has not been adequately explored. The complications of diabetes are not only medical, so most people rely on their individual resources such as spiritual beliefs, prayers, fasting and spiritual essence to cope with the illness. In Ghana, most people with chronic illness like diabetes often visit one prayer camp or the other for help. Unmet spiritual needs, especially if they involve spiritual struggles, can adversely affect health and may increase mortality independent of mental, physical, or social health [28]. Also, spiritual beliefs affect patients’ medical decisions and can influence compliance with treatments and self-care. Therefore, when these spiritual needs are not understood and incorporated into the care of diabetic patients, a lot cannot be achieved.

“Research shows that one’s own beliefs of efficacy function as an important determinant of motivation, affect, thought and action” [29]. “Since its conception, self-efficacy has been applied to different contexts” [30] and related to better health, better self-development and greater social integration. “Again, one’s beliefs about one’s self can act as a moderating variable in relationships. These beliefs have been considered in other areas within organisational

psychology, showing for example the moderating effects of self-esteem on the results of teamwork’ [31]. However, not much is known about how beliefs of efficacy moderate the relationships between family support, spirituality and diabetes self-care in the Ghanaian society. This has necessitated the call for public concern about the illness as well as government policies and measures to reverse the trend. Again, the few studies in Ghana and other parts of Africa suggests that diabetes patient’s adherence to self-care behaviours is still low despite its significance [32]. Considering the gaps and inconsistencies in literature concerning the relationship between family support, spirituality and self-care in T2DM and given that little is known about how social support and spirituality impacts self-care in T2DM in Ghana, there is the need to conduct research to investigate the relationship. There is also the need to examine specifically family support in diabetes self-care since few studies have been carried out on informal social support compared to formal support.

Thus, the study seeks to examine the following hypothesis:

H₁: Spirituality will positively correlate with self-care among T2DM patients.

H₁: Family support will be positively correlated with self-care.

H₁: Self-efficacy will moderate the relationship between family support and self-care.

H₁: Self-efficacy will moderate the relationship between spirituality and self-care.

2. MATERIALS AND METHODS

2.1 Research Design

A cross-sectional design was adopted and participants were made up of T2DM patients seeking treatment at some hospital facilities in Greater Accra region.

2.2 Recruitment of Participants

A sample of 270 was selected from some hospitals in Greater Accra using the convenient sampling technique. To be selected for the study, participants must be a T2DM for not less than six months, and must be aged 18 years and above. After gaining approval from the hospital, the necessary arrangements were made with the nurses in charge of the various diabetic clinics. Usually, the principal researcher and assistants

were present every morning at the various hospitals on clinic days. Permission was granted for the researcher to explain the purpose of the study to patients, and those who met the inclusion criteria and were willing to participate in the study were given the questionnaire to answer after signing the informed consent form. Participants were guided as to how to respond to the questionnaires without influence to the responses they provided.

2.3 Study Instrument

The data collection instruments included the Functional Assessment of Chronic Illness Therapy-spiritual well-being scale (FACIT-sp) developed by Brady, Peterman, Fitchett, Mo and Cella [33]; Diabetic Self-Management Questionnaire (DSMQ) by Schmitt et al. [62]; Self-efficacy for Diabetes Scale (SDS) also developed by the Stanford Patient Education Research Center [34]; and Social Support Scale for Self-Care in Middle-Aged Patients with T2DM (S4-Mad) developed by Naderimaghani, Niknami, Abolhassani, Hajzadeh, and Montazeri [35].

Questionnaires were pre-tested with 26 diabetic patients. The analysis of the pre-tested questionnaires using SPSS version 20.0 showed an overall Cronbach alpha of $\alpha = .941$ for the whole Social Support Scale for Self-Care in Middle-Aged Patients with T2DM scale. Again, the pre-test indicated a good reliability of $\alpha = .715$ for Self-Management Questionnaire (DSMQ). The self-efficacy scale as well produced a reliability of $\alpha = .81$. Based on comments from participants in the pre-test, some minor changes were made to arrive at the final questionnaire used for the study. Changes to the draft questionnaire comprised rewording some of the questions for clarity and comprehension, and correcting few typographical errors.

2.4 Data Analysis

After data collection, all questionnaires were stored in files and access restricted to the researcher. Data from the questionnaires were double-checked and cleaned before entered onto a computer by the investigator. Data was then entered onto International Business Machines of Statistical Package for Services Solutions

version 20 (IBM SPSS, 2020). Data cleaning was done to check for any error during the data entry. Frequencies and percentages were calculated for all socio-demographic factors. Hypothesis 1 and 2 were tested using Pearson Product Moment Correlation Coefficient. Hypothesis 3 and 4 was analysed using Hierarchical Multiple Regression.

3. RESULTS AND ANALYSIS

3.1 Demography of Participants

The participants were made up of 81.1% females 18.9% males. Eighty three respondents representing 30.7 % were in the age group of 58-67years, 24.4% were in the age range of 48-57 years, while 23% were aged between 68-77 years. Those within the ages of 38-47 were 8.9%, 6.3% (78-87 years), 5.9% (28-37 years) and 0.7% (18-27 years). Most of the respondents were above 50 years and in their late adulthood stage. Concerning the educational level, 81.5% had low level of education and 18.5% respondents had high level of education. Again, 51.5% were married while 4.8%, 14.4% and 29.3% were found to be single, divorced, separated and widowed respectively.

3.2 Preliminary Analysis

The preliminary analysis involves testing for normality, reliability and computing descriptive statistics for the variables studied. The results displayed in Table 1 indicated that the data was normally distributed as the test for normality produced Skewness and Kurtosis figures between -1 and +1. This indicated that the variables were all normal and can be used for parametric analyses [36]. Descriptive statistics of the predictor and criterion variables (means and standard deviations) were computed. Inter-correlations among the variables were also computed using Pearson Product Moment Correlation and the coefficients together with the Cronbach alpha of the scales used. The Cronbach alpha value of the whole scale was .84. Cohen [37] suggests that Cronbach alpha values of 0.7 and above are adequate for use in psychological research. The subscales for this study had Cronbach alpha values ranging from .74 to .89 (Table 1).

Table 1. Descriptive Statistics and Normality of the Study Variables (N=270)

	Min	Max	Mean	SD	Skewness	Kurtosis	Alpha Values
Spirituality	4.00	72.00	32.56	6.89	-.25	.83	.78
Self-Efficacy	20.00	80.00	59.22	11.66	-.60	.24	.85
Family-Support	30.00	111.00	54.27	16.37	.54	-.14	.74
Self-Care	11.00	36.00	21.17	3.49	.28	.12	.89

Table 2. Correlations among Variables (N=270)

	1	2	3	8
1 Spirituality	-			
2 Self-Efficacy	.34**	-		
3 Family-Support	.19*	.24**	-	
4 Self-Care	.15*	.14*	.24**	-

**p<.01, *p<.0

Table 3. Relationship between Spirituality and Self-Care (N=270)

Variable	N	Mean	Std. Dev.	Df	R	P
Spirituality	270	32.56	6.89	268	.151	.007
Self-care	270	21.17	3.49			

Table 4. Relationship between Family support and Self-Care (N=270)

Variable	N	Mean	Std Dev.	Df	R	P
Family S.	270	54.27	16.37	268	.237	.000
Self -care	270	21.17	3.49			

3.2.1 Hypothesis 1: Spirituality will positively correlate with self-care among T2DM patients

This hypothesis was tested using Pearson Moment Correlation Coefficient. As shown in Table 3, spirituality is positively related to self-care ($r = .151, p < .05$). We therefore fail to reject the hypothesis. The finding therefore supports the proposition that spirituality positively correlate with self-care among T2DM patients.

3.2.2 Hypothesis 2: Family support will be positively correlated with self-care

As indicated on Table 4, the relationship between family support and self-care is positive and significant [$r = .237, p < .01$]. We therefore fail to reject the hypothesis. Thus, the second hypothesis which stated that family support would be positively correlated with self-care was supported.

3.3 Testing for the Moderation

To test hypotheses 3 and 4 that seek to find out whether self-efficacy moderate the relationship

between family support and self-care, and the relationship between spirituality and self-care, the procedure proposed by Baron and Kenny [38] for testing moderation effect using hierarchical multiple regression was used. According to Baron and Kenny, [38] a common framework for illustrating moderating effect from both correlational and experimental perspectives is possible using a causal path analysis. The three causal paths as illustrated in Fig. 1 below (a, b, and c) was used. This involves fed into the criterion or dependent variable (DV), self-care: the effect of the IV (family support and spirituality) on the DV (self-care) (path a), the effect of the moderation variable (self-efficacy) on the DV (self-care) (path b), and the interaction or product of these two paths on the DV (path c). The moderator hypothesis is supported if the interaction term (path c) is significant [38]. With respect to the interaction term, the independent and the moderating variables were centred to reduce the effect of multicollinearity [39]. In centring, the mean value of the variable was subtracted from the individual scores of the variables.




Fig. 1. Path diagram of moderation model (Baron & Kenny, 1986)

Table 5. Results of Hierarchical Multiple Regression Analyses for the moderation effect of Self-efficacy on the relationship between family support and self-care (N=270)

Model		B	Std. Error	B	t
Step 1	(Constant)	18.424	.718		25.66**
	Family Support	.051	.013	.237	3.99**
Step 2	(Constant)	17.109	1.162		14.72**
	Family Support	.046	.013	.216	3.54**
	Self-Efficacy	.026	.018	.188	1.43*
Step 3	(Constant)	15.738	3.972		3.96**
	Family Support	.071	.069	.332	1.02*
	Self-Efficacy	.049	.066	.164	.74
	Family support*Self-Efficacy	.041	.112	.254	1.06*

$R^2 = .046$ for step 1, $R^2 = .063$ for step 2, $R^2 = .094$ for step 3, $\Delta R^2 = .056$ for step 1, $\Delta R^2 = .008$ for step 2, $\Delta R^2 = .031$ for step 3, ** $p < .01$ * $p < .05$

3.3.1 Hypothesis 3: Self-Efficacy would moderate the relationship between family support and self-care.

To test this hypothesis 3, the hierarchical regression in which three distinct steps are stipulated was conducted. The main effect of family support was entered first, the main effect of self-efficacy (moderator) was entered second, and the interaction term (family support and self-efficacy) was entered third.^[39] The basic requirement for testing for moderation effect is that there should be a relationship between the predictor variable(s) and the criterion variables [40] and this was met as illustrated in Table 2.

The result of the moderation analyses is shown in Table 5.

The step 1 of Table 5 indicated that family support had a significant influence on self-care ($\beta = .237, p < .01$). The second step also indicated that self-efficacy explained a significant increase in variance of self-care ($\Delta R^2 = .018, \beta = .188, p < .05$). From the third step of the regression

analysis, the interaction term between family support and self-efficacy explained a significant increase in variance in self-care ($\Delta R^2 = .031, \beta = .254, p < .05$). We therefore fail to reject the hypothesis. Thus, the third hypothesis which stated that self-efficacy would moderate the relationship between family support and self-care was supported.

3.3.2 Hypothesis 4: Self-efficacy would moderate the relationship between spirituality and self-care.

To test hypothesis 4, the hierarchical regression in which three distinct steps are stipulated was also conducted. The main effect of spirituality was entered first, the main effect of self-efficacy (moderator) was entered second, and the interaction term (spirituality and self-efficacy) was entered third [39]. The basic requirement for testing for moderation effect that there should be a relationship between the predictor variable(s) and the criterion variables [40] was met as illustrated in Table 2. The result is shown in Table 6.

Table 6. Results of Hierarchical Multiple Regression Analyses for the moderation effect of Self-Efficacy on the relationship between Spirituality and Self-Care (N=270)

Models		B	Std. Error	B	t
Step 1	(Constant)	18.685	1.018		18.35
	Spirituality	.076	.031	.151	2.49*
Step 2	(Constant)	17.469	1.281		13.64
	Spirituality	.059	.032	.117	1.83*
Step 3	Self-Efficacy	.030	.019	.100	1.55
	(Constant)	13.880	4.649		2.98
	Spirituality	.173	.146	.342	1.19
	Self-Efficacy	.092	.080	.308	1.15
	Spirituality*Self-Efficacy	-.196	.244	-.358	-.80

*R*² = .023 for step 1, *R*² = .032 for step 2, *R*² = .034 for step 3, $\Delta R^2 = .023$ for step 1, $\Delta R^2 = .009$ for step 2, $\Delta R^2 = .002$ for step 3, **p* < .05

Assessing the results in Table 6, the step 1 indicated that spirituality had a significant influence on self-care ($\beta = .153, p < .05$). From the second step, self-efficacy did not explain a significant increase in variance of self-care ($\Delta R^2 = .009, \beta = .100, p > .05$). The third step of the regression analysis revealed that the interaction term between spirituality and self-efficacy did not explain a significant increase in variance in self-care ($\Delta R^2 = .002, \beta = -.358, p > .05$). Thus, the fourth hypothesis which stated that self-efficacy would moderate the relationship between spirituality and self-care was not supported.

4. DISCUSSION

Family support and spirituality are psychosocial factors that have been associated with optimum self-care. Every individual at a point in time rely on his/her spiritual essence or family for support. The hypothesis that spirituality will correlate positively with self-care was supported. This means that the more people are spiritual the more they engage in self-care practices. The current finding is confirmed by a study by Polzer and Miles [41] who found that spirituality played a significant role that influenced diabetes self-management among African Americans with diabetes. Similarly, the finding is supported by 98 (82%) studies which found positive relationships between spirituality and well-being [42]. Again, the current study is supported by a study in which positive association was found between spirituality and a healthier diet [43]. The study result is also confirmed by a meta-analysis involving 37 studies. Twenty five of these studies reported significant positive relationship between spirituality and greater exercise or physical activity. This finding however is incongruent with a study by Chio [16]. Chio cited by Abraham [44] in a systematic literature review on factors

contributing to self-care among T2DM patients found that patients who were more spiritual had poor self-care than those loose or non-spiritual patients. Meaning more spiritual T2DM patients do not adhere or engage in self-care practices instead they resort to prayer and other spiritual beliefs. Additionally, the finding is inconsistent with a study that found that more spirituality was associated with poor self-care and poor coping abilities among T2DM patients [45]. This could mean that those who do not have high spirituality could not have a good sense of the meaning of life. Therefore, they are unable to take an active approach in their self-care. The positive relationship between spirituality and self-care is explained by Mazlom, Afkhami-Ardekani and Dadgari [46] to be the identification of God or divine power as a source of power, hope and inspiration for the diabetics in their self-care and management. Again, the significant relationship could be explained by the fact that when diabetics are spiritual they have strong meaning in life. This meaning helps them accept their condition as a trail which must be fought and won. This mind-set helps them to engage in appropriate self-care. This is confirmed by a study in which sense of meaning helped patients cope with their disease, reframed their lives and had an optimistic look on life and a "fighting spirit" against their disease [47]. Similarly, an obvious explanation for the relationship is that when people have strong reasons for living (career, children, and family) they are more likely to take steps or engage in practices (glucose monitoring, diet, and exercise) that would prolong their lives. Given a meaning of spirituality as belief or connectedness with your creator, engaging in little prayer or being prayerful or having meaning in life, we can conclude that prayer, meaning, belief connected to higher being helps the diabetic patient rise above the

physical limitations of the disease. Zinnbauer et al. [48] defines spirituality as either having a relationship with God or a higher power, or having a personal belief system in God or a higher power. This literature-based definition and participants' experience of spirituality seemed similar. Patients had some spiritual essence which could be due to the fact that they are all experiencing personal crisis (illness). This is confirmed by several studies which indicated that people's spirituality or religious faith increases when experiencing personal crisis due to illness. Again, according to Jung [49], individuals in their midlife (40-59) begin to typically turn inward to explore the more aspect of themselves.

The second hypothesis that family support would be positively correlated with self-care was supported. This finding support previous studies where a positive relationship was observed between family support and self-care behaviour in adults with non-insulin dependent diabetes [50]. This is consistent with a study among women with gestational diabetes in which adherence to dietary recommendations was strongly associated with familial social support [51]. Also, the finding is in line with a study by Fisher, Chesla and Staff [52]. In their study, strong family and social support appear to have a positive impact on Glycemic control and or self-management behaviour. This shows that the more diabetics receive support from their family, the more likely they engaged in self-care practices. This finding is in agreement with the Ecological model which presents health as an interaction between the person and their ecosystem such as the family, community and physical environment. There is a reciprocal influence in that people are affected by their ecosystem and likewise people affect their ecosystem [53]. The family as part of the ecosystem is a system where the care, protection and development of individuals are facilitated. This model assumes that the individual efforts at behaviour change will be more likely to succeed within supportive environment such as the family. These environments have collective and cumulative impact on well-being. The Orem self-care theory also supports the current findings. The Orem's self-care theory is based on the assumption that all clients have the capacity to act in order to maintain their health and treat themselves in case of sickness or injury [11]. It is however a complex activity but improves through daily practices and requires help, assistance and guidance from others such as health

professionals and family. Orem in her theory emphasizes that an individual's self-care ability and self-care behaviour is influenced by basic conditioning factors, such as the individual's family system. A functional component of a family system is family support which helps to maintain a healthy lifestyle. Again, Carter [11] emphasizes that basic conditioning factor such as family system factors influence self-care. According to the findings of this present study there is positive correlation among T2DM self-care and family support. Meaning the more support is received or provided; the more self-care activities are performed by diabetics [16].

Additionally, the positive relationship between family support and self-care could be explained by the African-centred worldview, where Africans believe that all elements of the universe (people, animals and inanimate object) are viewed as interconnected. Since they are dependent upon each other, they are, in essence, considered as one so that the human being is never an isolated individual but always the person in the community [55]. The community defines the person. As Mbiti [55] puts it "I am, because we are; and since we are, therefore I am" (p.57). The collective nature of human beings entails collective responsibility for what happens to individuals. "Whatever happens to the individual happens to the whole group and whatever happens to the whole group happens to the individual" (p.57) [55]. This shows that what affect patients affect the entire family so self-care becomes a collective responsibility. Again, the ties of kinship may have created and facilitated the family members' commitment towards the participants and their self-care. If family members are financially sound, they may be eager and readily available to offer support to the patient with diabetes [56]. This in turn affects self-care.

The third prediction that the relationship between family support and self-care will be moderated by self-efficacy was supported. The findings are consistent with a study by Kaşıkçı and Alberto [57]. Their study revealed statistically significant positive relationships between family support and self-care behaviour ($r = 0.302$; $p = 0.01$) and between self-efficacy and self-care behaviour ($r = 0.186$; $p = 0.01$). There was also a statistically significant positive relationship between family support and self-efficacy ($r = 0.412$; $p = 0.01$). Furthermore, the finding is congruent with the Biopsychosocial-spiritual theory [58]. According to the model, an interaction between various factors results in optimal health and not a single

factor. In addition, Potter and Zauszniewski [59] conducted a correlational cross-sectional study using a survey format to examine 47 community-based adults with rheumatoid arthritis. When social, emotional, and physical stressors were examined individually, the individual items did not predict general health perception, but when taken together, a significant effect on health perception was observed. This means that the interaction between family support and self-care is influenced by self-efficacy. Contrary to the current finding, Williams and Bond [60] reported that social support was not a predictor of self-care among 94 diabetics when the effects of self-efficacy were controlled. This finding can be explained by the fact that self-efficacy serves as a bridge that propels an individual to take action. Thus, people's confidence (belief) that they can carry out a task (self-care) coupled with the support from family increases their self-care ability.

The fourth prediction that the relationship between spirituality and self-care will be moderated by self-efficacy was not supported. This means that self-efficacy could not influence the relationship between spirituality and self-care. The lack of relationship is explained by the idea that spirituality alone may help some individuals to "gain a sense of control over their lives and take action. The finding was inconsistent with the Biopsychosocial-spiritual theory [59]. According to this model, illness and health result from the interaction between biological, psychological, spiritual and social factors. This suggests that for an individual to achieve optimal self-care it requires interaction of various factors. Several other factors must play a significant role to enable people achieve adequate self-care in dealing with T2DM.

A strength of the study is that it would help shape the scope of management of diabetes mellitus which is predominantly biomedical by including more psychosocial interventions into diabetes management. It will enable the development of family tailored interventions that would help improve the diabetics' self-care practices and health outcome in general. On the other hand, the findings of the study are limited by the convenience sampling approach which could not permit the generalization beyond the study population. Additionally, the findings are only applicable to Type 2 diabetics and may not fit well with other chronic illness such as cancer. This is mainly because different illnesses have different symptoms and effects, and the impact of

family support and spirituality on self-care may differ from illness to illness.

5. CONCLUSIONS

It is established that T2DM patients who are more spiritual are more likely to care for themselves. It also revealed that the more support diabetics receive from their family, the more they care for themselves. Thus, more availability of social support is more likely to increase self-care among patients. The association between family support and self-care is moderated by self-efficacy. This shows that a person's belief in his capability could also contribute to social support influencing self-care. However, self-efficacy does not influence the relationship between spirituality and self-care. Patient's confidence (belief) that they can care for themselves can actually influence their capabilities to care for themselves. Finally, self-efficacy, spirituality and family support are critical in diabetes self-care. It is therefore beneficial that health care providers incorporate psychosocial interventions into health care delivery.

6. RECOMMENDATIONS

This study has important implications for current policies and programs that are designed to improve self-care among diabetics. Self-care interventions that are solely medical and do not include psychosocial intervention may not achieve optimum management of diabetes. It is therefore recommended that Ghana Health service and Ministry of Health spearhead the drawing of a comprehensive diabetes management programme that would incorporate psychosocial variables. Furthermore, holistic approach should be adopted in the management of the diabetes condition. That is, an approach to health care delivery known as the biopsychosocial involving healthcare professionals with varied expertise such as doctors, nurses, psychiatrists, clinical psychologists and dieticians as recommended by Kwakye [61].

Again, spiritual leaders should be included in the management of diabetes to address the spiritual needs of diabetics since respondents in this study identify themselves with a particular form of religion, and this show the importance respondents attach to religion. This will compliment health professionals' effort. Spiritual leaders, counsellors, social workers and health workers are recommended to organise

workshops that help the patient explore their spiritual essence so they can transcend beyond their condition. The Ghana Diabetes Federation should intensify education on diabetes to create awareness and help patients engage in healthy practices.

CONSENT

Informed consent was obtained from each eligible participant after they had been provided with information sheet containing detailed information about the study.

ETHICAL APPROVAL

Institutional and ethical approval was obtained from the University of Cape Coast Institutional Review Board of the School of Graduate Studies and Research and the selected hospitals before the commencement of the study. This was to ensure that the study was ethically sound and did not violate the rights and privacy of the vulnerable participants.

SOURCE OF FUNDS

The study was funded by the first author with support from the other authors.

ACKNOWLEDGEMENT

A special acknowledgement goes to Prof. Samuel A. Danquah, Dr. Mavis Asare and the research assistants for their enormous support. We are also grateful to management of the hospitals selected for the study and the patients who made time to respond to the questionnaire.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aikins AD, Owusu-Dabo E, Agyemang C. Diabetes in Ghana: a review of research on prevalence, experiences and healthcare. *Chronic Non-Communicable Diseases in Ghana. Multidisciplinary Perspectives.* 2013;41-55.
2. Walsh M, Crumbie A. *Watson's Clinical Nursing and Related Sciences E-Book.* Elsevier Health Sciences; 2007.
3. World Health Organization. *Global report on diabetes.* World Health Organization; 2016.
4. Federation ID. *IDF diabetes atlas 8th edition.* International diabetes federation. 2017:905-11.
5. Imamura F, O'Connor L, Ye Z, Mursu J, Hayashino Y, Bhupathiraju SN, Forouhi NG. Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction. *Bmj.* 2015;351.
6. InterAct consortium d. romaguera-bosch@imperial.ac.uk. Consumption of sweet beverages and type 2 diabetes incidence in European adults: results from EPIC-InterAct. *Diabetologia.* 2013;56:1520-30.
7. Malik VS, Popkin BM, Bray GA, Després JP, Willett WC, Hu FB. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a Meta-analysis. *Diabetes care.* 2010;33(11):2477-83.
8. Hörnsten Å. *Experiences of diabetes care-patients' and nurses' perspectives* (Doctoral dissertation, Omvårdnad, Umeå Universitet).
9. Institute ID. *Type 2 Diabetes: Practical Targets and Treatments.* Melbourne: The International Diabetes Institute; 2005.
10. Lin EH, Von Korff M, WHO WMH survey consortium. Mental disorders among persons with diabetes—results from the World Mental Health Surveys. *Journal of Psychosomatic Research.* 2008;65(6): 571-80.
11. Orem DE, Taylor SG, Renpenning KM. *Nursing: concepts of practice 5th ed.* InSt. Louis (US): Library of Congress 1995.
12. Anderson RM, Funnell MM, Butler PM, Arnold MS, Fitzgerald JT, Feste CC. Patient empowerment: results of a randomized controlled trial. *Diabetes care.* 1995;18(7):943-9.
13. Uchino BN. *Social support and physical health: Understanding the health consequences of relationships.* Yale University Press; 2004.
14. Bai YL, Chiou CP, Chang YY. Self-care behaviour and related factors in older people with Type 2 diabetes. *Journal of clinical nursing.* 2009;18(23):3308-15.
15. Ciechanowski P, Russo J, Katon WJ, Lin EH, Ludman E, Heckbert S, Von Korff M, Williams LH, Young BA. Relationship styles and mortality in patients with diabetes. *Diabetes Care.* 2010;33(3): 539-44.

16. Choi SE. Diet-specific family support and glucose control among Korean immigrants with type 2 diabetes. *The Diabetes Educator*. 2009;35(6):978-85.
17. Kwakye IN, Antwi J, Hormenu T. Lived Experiences of Diabetes Patients in Rural Areas of Ghana: Discovering the Forces that Determine Psychosocial Care; 2021.
18. Paddison CA. Family support and conflict among adults with type 2 diabetes: Development and testing of a new measure. *European Diabetes Nursing*. 2010;7(1):29-33.
19. Rintala TM, Jaatinen P, Paavilainen E, Åstedt-Kurki P. Interrelation between adult persons with diabetes and their family: a systematic review of the literature. *Journal of family nursing*. 2013;19(1):3-28.
20. Anderson D, Christison-Lagay J. Diabetes self-management in a community health center: improving health behaviors and clinical outcomes for underserved patients. *Clinical Diabetes*. 2008;26(1):22-7.
21. Sabone MB. The illness demands of diabetes on couples in Botswana. *Journal of Family Nursing*. 2008;14(3): 363-82.
22. Strom JL, Egede LE. The impact of social support on outcomes in adult patients with type 2 diabetes: a systematic review. *Current diabetes reports*. 2012;12(6):769-81.
23. Fitchett G, Murphy PE, Kim J, Gibbons JL, Cameron JR, Davis JA. Religious struggle: Prevalence, correlates and mental health risks in diabetic, congestive heart failure, and oncology patients. *The International Journal of Psychiatry in Medicine*. 2004;34(2):179-96.
24. Crabtree S, Pelham B. World Gallup Poll, Religion Provides Emotional Boost to World's Poor, *Gallup*; 2009.
25. Dalgarrondo P, Marín-León L, Botega NJ, De Azevedo Barros MB, Bosco de Oliveira H. Religious affiliation and psychiatric morbidity in Brazil: higher rates among evangelicals and spiritists. *International Journal of Social Psychiatry*. 2008;54(6):562-74.
26. Leeman J, Skelly AH, Burns D, Carlson J, Soward A. Tailoring a diabetes self-care intervention for use with older, rural African American women. *The Diabetes Educator*. 2008;34(2):310-7.
27. Deatcher J. Spiritual self-care and the use of prayer. *Diabetes self-management*. 2002;19(6):57-9.
28. Pargament KI, Koenig HG, Tarakeshwar N, Hahn J. Religious struggle as a predictor of mortality among medically ill elderly patients: A 2-year longitudinal study. *Archives of internal Medicine*. 2001;161(15):1881-5.
29. Bandura A. Exercise of personal agency through the self-efficacy mechanism. *Self-efficacy: Thought control of action*. 1992;1:3-7.
30. Bandura A. *Self-efficacy: The exercise of control*. Macmillan; 1997.
31. Brief AP, Aldag RJ. The "self" in work organizations: A conceptual review. *Academic of Management Review*. 1998;6:75-88.
32. Emmanuel OO, Otovwe A. Patterns of adherence to management among patients with type 2 diabetes mellitus in South-South Region of Nigeria. *Journal of Social Health and Diabetes*. 2015;3(02): 115-9.
33. Brady MJ, Peterman AH, Fitchett G, Mo M, Cella D. A case for including spirituality in quality of life measurement in oncology. *Psycho-Oncology: Journal of the Psychological, Social and Behavioral Dimensions of Cancer*. 1999;8(5):417-28.
34. BOOK QC. Stanford Patient Education Research Center. *American Journal of Public Health*. 1990;80:446-52.
35. Naderimagham S, Niknami S, Abolhassani F, Hajizadeh E, Montazeri A. Development and psychometric properties of a new social support scale for self-care in middle-aged patients with type II diabetes (S4-MAD). *BMC Public Health*. 2012;12(1): 1-9.
36. Field A. *Discovering statistics using SPSS*. Berlin: Heidelberg; Sage Publication Limited; 2005.
37. Cohen J. Quantitative methods in psychology: A power primer. In *Psychological Bulletin*; 1992.
38. Baron RM, Kenny DA. The moderator–mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of personality and social psychology*. 1986;51(6):1173.
39. Agerström J, Björklund F, Allwood CM, Aiken LS, West SG. *Multiple Regression: Testing and Interpreting Interactions*. London: Sage. *Journal of Applied Psychology*. 1991;73:305-7.
40. Holmbeck GN. Toward terminological, conceptual, and statistical clarity in the

- study of mediators and moderators: examples from the child-clinical and pediatric psychology literatures. *Journal of consulting and clinical psychology*. 1997;65(4):599.
41. Polzer RL, Miles MS. Spirituality in African Americans with diabetes: Self-management through a relationship with God. *Qualitative Health Research*. 2007;17(2):176-88.
 42. Crabtree S & Pelham B. World Gallup Poll, Religion Provides Emotional Boost to World's Poor, Gallup; 2009.
 43. Underwood SM, Powell RL. Religion and spirituality: influence on health/risk behavior and cancer screening behavior of African Americans. *ABNF Journal*. 2006;17(1).
 44. Abraham M. Self-Care in type 2 Diabetes: A Systematic Literature Review on Factors Contributing to Self-Care among Type 2 Diabetes Mellitus Patients. Unpublished master's dissertation, School of Health and Caring Sciences Linnaeus University; 2011.
 45. Samuel-Hodge CD, Watkins DC, Rowell KL, Hooten EG. Coping styles, well-being, and self-care behaviors among African Americans with type 2 diabetes. *The Diabetes Educator*. 2008;34(3):501-10.
 46. Mazlom N, Afkhami-Ardekani M, Dadgari A. Relationship between the religion orientation and coping with diabetes in patients with type 2 diabetes. *Iranian Journal of Diabetes and Obesity*. 2013;5(1):33-9.
 47. Yi MS, Mrus JM, Wade TJ, Ho ML, Hornung RW, Cotton S, Peterman AH, Puchalski CM, Tsevat J. Religion, spirituality, and depressive symptoms in patients with HIV/AIDS. *Journal of general internal medicine*. 2006;21(S5): S21-7.
 48. Zinnbauer BJ, Pargament KI, Cole B, Rye MS, Butfer EM, Belavich TG, Hipp K, Scott AB, Kadar JL. Religion and spirituality: Unfuzzifying the fuzzy. *Sociology of religion*. 2015:29-34.
 49. Jung CG, von Franz ML, Henderson JL. *Man and his symbols*. New York: Bantam Doubleday Dell Publishing Group; 1964.
 50. Wang CY, Fenske MM. Self-care of adults with non-insulin-dependent diabetes mellitus: influence of family and friends. *The Diabetes Educator*. 1996;22(5): 465-70.
 51. Nicklett EJ, Liang J. Diabetes-related support, regimen adherence, and health decline among older adults. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 2010;65(3):390-9.
 52. Fisher L, Chesla CA, Skaff MM, Gilliss C, Mullan JT, Bartz RJ, Kanter RA, Lutz CP. The family and disease management in Hispanic and European-American patients with type 2 diabetes. *Diabetes Care*. 2000;23(3):267-72.
 53. Glanz K. *Theory at a glance: A guide for health promotion practice*. US Department of Health and Human Services, Public Health Service, National Institutes of Health, National Cancer Institute; 1997.
 54. Carter PA. Self-care agency: the concept and how it is measured. *Journal of Nursing Measurement*. 1998;6(2):195-207.
 55. Mbiti J. *African religions and philosophies* Garden City. NY: Anchor Books; 1970.
 56. Kirk JK, Ebert CN, Gamble GP, Ebert CE. Social support strategies in adult patients with diabetes: a review of strategies in the USA and Europe. *Expert Review of Endocrinology & Metabolism*. 2013;8(4): 379-89.
 57. Kara Kaşıkçı M, Alberto J. Family support, perceived self-efficacy and self-care behaviour of Turkish patients with chronic obstructive pulmonary disease. *Journal of Clinical Nursing*. 2007;16(8): 1468-78.
 58. Kozak L, Boynton L, Bentley J, Bezy E. Introducing spirituality, religion and culture curricula in the psychiatry residency programme. *Medical Humanities*. 2010;36(1):48-51.
 59. Potter ML, Zauszniewski JA. Spirituality, resourcefulness, and arthritis impact on health perception of elders with rheumatoid arthritis. *Journal of Holistic Nursing*. 2000;18(4):311-31.
 60. Williams KE, Bond MJ. The roles of self-efficacy, outcome expectancies and social support in the self-care behaviours of diabetics. *Psychology, health & medicine*. 2002;7(2):127-41.
 61. Kwakye IN. *Lived Experiences of Diabetics Living in Rural Areas in the Eastern Region (Doctoral dissertation, University of Cape Coast)*; 2020.

62. Schmitt A, Gahr A, Hermanns N, Kulzer B, Huber J, Haak T. The Diabetes Self-Management Questionnaire (DSMQ): development and evaluation of an instrument to assess diabetes self-care activities associated with glycaemic control. Health and quality of life outcomes. 2013;11(1):1-4.

© 2022 Ampofo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/88897>