



---

## Application of Mitscherlich-bray Equation to Establish Fertilizer Recommendation for Strawberry under Aduyon Clay Loam

Gerlie S. Jambaro <sup>a</sup>, Louell M. Ozarraga <sup>a</sup> and Edgel O. Escomen <sup>a\*</sup>

<sup>a</sup> Department of Plant Science, College of Agriculture, Mindanao State University Marawi Campus, Lanao del Sur- 9700, Philippines.

### Authors' contributions

This work was carried out in collaboration among all authors. Author GSJ designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Author LMO wrote the results and discussion. Author EOE edited and package the manuscript. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/IJPSS/2022/v34i2031186

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/88680>

Original Research Article

Received 15 April 2022  
Accepted 27 June 2022  
Published 28 June 2022

---

### ABSTRACT

The nutrient requirement of strawberry plants is high making them very responsive to fertilization. This study utilized the Mitscherlich-Bray equation to determine the theoretical maximum yield as the basis of comparison to the actual yield. The economics of fertilizer application was also considered in the determination of the fertilizer recommendation. The theoretical maximum yield of the study is 12.29 t/ha<sup>-1</sup>. Strawberry is found responsive to nitrogen and potassium fertilization. The application of 225-225-360 kg ha<sup>-1</sup> of N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O, respectively could achieve 95% of the theoretical maximum yield, however, it had a much higher amount spent on fertilization per kg of fruit yield compared to the current Hafza's recommendation. The fertilizer recommendation of 150-150-240 kg ha<sup>-1</sup> of N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O, respectively still the best option under Aduyon clay loam for both economic and horticultural benefits.

**Keywords:** Fertilizer recommendation; mitscherlich-Bray equation; theoretical maximum yield; actual yield.

## 1. INTRODUCTION

Strawberry (*Fragaria ananassa* Duch) is a crop with high economic value and documented nutritional benefits, popular in consumer diet, and suitable for sensory evaluation [1]. It grows best in cool areas and is tolerant to most soil types.

In the Philippines, the demand for strawberry fruits is high especially during holiday seasons when strawberry is utilized for dessert and wines [2]. La Trinidad, Benguet is known as the strawberry capital of the Philippines. They devoted 1.54% of the total agricultural land area to strawberry production. There are 18,820 farmers engaged in such production [3]. The average productivity of strawberries is 12 t/ha with prices ranging from PHP 250 to 300/kg which gave the province PHP 3.0 to 3.6 M gross income per hectare. On top of this, the biggest impact of strawberries had brought the municipality into the world of agritourism which registered about 259,000 tourists in 2016 (Ticbaen, 2017).

In some areas in Mindanao, around 30 farmers in Dumangas, T'boli, and South Cotabato were the beneficiary of the government anti-poverty program to start strawberry production in their 1.4 ha of lands [4]. In Davao, the Benwa Farm in Marilog District, Buda, Davao City makes its name known through strawberry picking [5] as well as in Vincent Heights, Brgy. Obrero, Mintal [6]. Moreover, the Taglucop Strawberry Farm in Kitaotao, Bukidnon proves successful in strawberry production which somehow has the same features as that of Lanao del Sur.

The successful venture of strawberry production and processing in Bukidnon as exemplified by the Epol Strawberry Farmers Association offers an enormous possibility of augmenting the lives of low-income farmers in Lanao del Sur. Both Bukidnon and Lanao del Sur are endowed with a cool climate which is an important requirement for strawberry production. Lanao del Sur, particularly, Marawi City is nested on a plateau with an altitude of 833 meters above sea level. The cool temperature of the area ranges between 16.0 to 22.0 °C with an even distribution of rainfall throughout the year.

Fertilizer is supplied to plants to meet the nutrient demand for growth and development. It assists the plants in the growth of vegetative mass and maintains the plant's current activity for fruit development. For strawberries, fertilizer

requirement is high despite their small size [7]. At the high level of nitrogen usually at 500 kg/ha, production of runners or strawberry transplants is viable under Claveria, Misamis Oriental conditions [2] while fruit production requires a high amount of potassium [8].

Fertilizer recommendation needs to be reevaluated because the current rates are already obsolete. Since strawberry is viable to be grown in the Lanao del Sur Province, a fertilizer program should be developed to increase the size and weight of the fruit, attractive to the market, which is considered one of the challenges in strawberry production [9-12]. The use of the Mitscherlich-Bray equation was found helpful in understanding the response of sweet potatoes to fertilizer application [13]. Hence, this study utilized the Mitscherlich-Bray equation on judicious use of fertilizer for strawberry production.

This study aims to determine the theoretical maximum yield as the basis of comparison to the actual yield. It also looks into the economics of fertilizer application under Aduyon clay loam.

## 2. MATERIALS AND METHODS

This study was conducted at the College of Agriculture Experimental Area, Mindanao State University-Marawi Campus, Lanao del Sur (7.9939° N, 124.2581° E) under Aduyon clay loam. Aduyon clay loam falls under *Typic Kandiodult*. It is soil that has undergone extensive leaching and accumulation of clays in the subsoil with sufficient moisture throughout the year. It has a low cation exchange capacity. A typical representative of *Kandiodults* with andesite and basaltic rocks as the parent material. Hence, fertilization and liming are highly required.

Variety *Florida* was used in this study because of its aroma and sweetness even though it is medium-sized. Due to expensive planting materials, mother strawberry plants were allowed to reproduce for six months inside the Plant Science Nursery. Fertilizer recommendations were 150- 150- 240 kg/ha N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O, respectively [14]. Treatments were established using this recommendation with no application, 50, 100, 150, 200, and 250% levels. It was laid out using Randomized Complete Block Design with four replications. Fertilizer was applied in four split applications from transplanting until fruit set development with 2 weeks intervals. The soil

contained 0.11% Total N, 1.06 ppm available P and 104.72 ppm exchangeable K [15].

Damaged, old and diseased leaves were removed from the plants with the use of pair of scissors. Removal of the runner was done after 60 days to promote flowering. Cypermethrin was used to control insect pests following the manufacturer’s recommendation.

The yield data collected were subjected to the Mitscherlich-Bray equation as given below (Afzal et al 2014).

$$\log (A-y) = \log A-c1b-cx$$

The theoretical yield was obtained by plotting log y (actual yield) to 1/x (amount of fertilizer applied).

Constants c1 and c were calculated separately following the equations:

$$c_1 = \frac{\log A - \log(A-y_0)}{b}$$

$$c = \frac{(\log A - c_1 b) - \log(A-yx)}{x}$$

Where A=% theoretical maximum yield, y=actual yield obtained in q ha<sup>-1</sup>, b=native soil test value in kg ha<sup>-1</sup>, x=fertilizer nutrient applied in kg ha<sup>-1</sup>,

c<sub>1</sub> and c=constants i.e., the efficiency of soil and fertilizer nutrients respectively; yx is the yield obtained from the fertilized plots., y<sub>0</sub> is the yield obtained from control plots.

### 3. RESULTS AND DISCUSSION

The theoretical yield of strawberries was determined using the actual yield in terms of the Mitscherlich-Bray equation by plotting the log y and 1/x (Fig. 1). Strawberry plants can yield as much as 12.29 t ha<sup>-1</sup>. This theoretical maximum yield was comparable to the Municipality of Benguet with an average yield of 12 t/ha (Ticbaen, 2017). The c<sub>1</sub> and c values were 0.0061 and 0.00464 for N, 0.03179 and 0.00471 for P<sub>2</sub>O<sub>5</sub> and 0.000377 and 0.002903 for K<sub>2</sub>O, respectively. The ratio of c<sub>1</sub>/c indicates that a higher ratio means a lower response of strawberry plants while a lower ratio means a greater response to fertilizer application [13]. A lower ratio was obtained from the fertilization of N and K<sub>2</sub>O primarily because the interactive effects of these mineral nutrients are necessary as the former improved yield while the latter improved fruit quality [16]. The application of N and P alone results in a lower yield than with a complete NPK application [17]. The importance of K is vital since it participated in many metabolic activities such as fruit quality improvement and stress tolerance.

**Table 1. Strawberry yield and the efficiency coefficient of soil and fertilizer NPK**

Treatments N applied kg ha <sup>-1</sup>	Actual yield (t ha <sup>-1</sup> )	Calculated Log y	1/x	c1	c	c1/c
				0.0016		
0	2.04					
75	9.25	0.9663	0.01333		0.007046	
150	11.40	1.0568	0.00667		0.007067	
225	11.79	1.0715	0.00444		0.00583	0.344454
300	9.82	0.9923	0.00333		0.002062	
375	8.70	0.9397	0.00267		0.001216	
Mean					(0.00464)	
Theoretical yield	12.29					
Treatments P <sub>2</sub> O <sub>5</sub> applied kg ha <sup>-1</sup>	Actual yield (t ha <sup>-1</sup> )	Calculated log y	1/x	c1	c	c1/c
				0.03179		
0	2.04					
75	9.25	0.9663	0.01333		0.0071985	
150	11.40	1.0568	0.00667		0.0071436	
225	11.79	1.0715	0.00444		0.0058812	6.743697
300	9.82	0.9923	0.00333		0.0021001	
375	8.70	0.9397	0.00267		0.0012465	
Mean					(0.00471)	
Theoretical yield	12.29					

Treatments K <sub>2</sub> O applied kg ha <sup>-1</sup>	Actual yield (t ha <sup>-1</sup> )	Calculated log y	1/x	c1	c	c1/c
0	2.04			0.000377		
120	9.25	0.9663	0.9663		0.004404	
240	11.40	1.0568	1.0568		0.004417	
360	11.79	1.0715	1.0715		0.003644	0.129947
480	9.82	0.9923	0.9923		0.001289	
600	8.70	0.9397	0.9397		0.00076	
Mean					(0.002903)	
Theoretical yield	12.29					




Fig. 1. Theoretical maximum yield (A) of strawberry from the plot of log y vs 1/x as affected by levels of NPK

Since this experiment was conducted in the field setting, plants received more stress brought by environmental factors such as too much sunlight, air and soil temperature and rainfall [18]. Nitrogen is necessary during vegetative growth towards flower development. Also, the difference in the amount of fertilizer among treatments used was wide, thus responses were highly defined. The response of phosphorus in this experiment is low maybe because of the strongly acidic soil pH (pH 5.5). Phosphorus availability is between 6.0-7.5 [19], hence liming is necessary for this soil.

Fertilizer recommendation of 225-225-360 kg ha<sup>-1</sup> of N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O, respectively could achieve 95% of the theoretical maximum yield based on the actual yield. The current fertilizer recommendation of Hafza (2008) which is 150-150-240 kg ha<sup>-1</sup> of N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O, respectively could achieve 93% of the theoretical maximum yield. Half of Hafsa's fertilizer recommendation could achieve 75% while twice the amount could attain 80%. This means 225-225-360 kg ha<sup>-1</sup> of N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O, respectively already reached the peak of fruit production following fertilizer application. A further increase in the fertilizer application could lead to a decline in yield. In terms of the % calculated yield, it can achieve up to 98% of the strawberry yield among N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O fertilizer applications.

Strawberry is highly responsive to fertilizer application [17]. Nutrient management is a key factor to ensure higher yields and fruit quality of strawberry plants (Trejo-Trelles and Gomez-Merino, 2014). With this, fertilization should be a

critical consideration in the production management of strawberries to avoid undersupply which resulted in low and unprofitable yield, or oversupply which will increase production cost and later contribute to environmental problems [13].

The result of the experiment can also be interpreted using the Baule unit. Baule unit is defined as the amount of nutrient added which results in moving one way half closer to maximum yield [20]. This study explains that at very low rating (Table 2), one Baule unit is equal to 47.45 kg ha<sup>-1</sup> for N and 87.69 kg ha<sup>-1</sup> for K, both 50% target yield. There was no recommendation drawn from P<sub>2</sub>O<sub>5</sub> because of the wide range of treatments used. The common range for phosphorus is between 10 kg/ha for very low to 50 kg/ha for high [21]. The experiment ranges from 75-325 kg/ha which already surpassed the high P<sub>2</sub>O<sub>5</sub> rating.

Table 3 shows the amount of fertilizer computed based on the Mitscherlich-Bray Equation depending on the target yield set. The N recommendation for 50, 60, 70, 80 and 90% target yield of the highest actual yield obtained from the experiment were 47.45, 66.52, 90.50, 122.84, and 172.62 kg/ha<sup>-1</sup>, respectively for very low and 19.82, 38.94, 62.92, 95.25 to 145.03 kg/ha<sup>-1</sup>, respectively for low. No fertilizer recommendation is needed for soil with medium soil nitrogen content if the target yield is between 50-60% and high N content between 50-80%. Medium N soil requires 21.54, 53.87 and 103.65 kg/ha-1 of N application for a yield target between 70, 80 and 90% target yield, respectively.




Fig. 2. Percent actual yield and calculated yield to that of the theoretical yield

**Table 2. General Interpretation of Total Nitrogen, Extractable Phosphorus and Exchangeable Potassium (PCARRD 2007)**

Rating	Nitrogen (%)	Phosphorus (mg/kg or ppm)	Potassium	
			(mg/kg or ppm)	(meq/100 g)
Very high	>1.0		>460	>1.2
High	0.6 - 1.0	>40	230-460	0.6-1.2
Medium	0.3 - 0.6	20-40	115-230	0.3-0.6
Low	0.1 - 0.3	10-20	40-115	0.1-0.3
Very Low	<0.1	<10	<40	<0.1

**Table 3. Fertilizer Recommendation for very low, low, medium, and high soil content for N and K<sub>2</sub>O**

% From the Highest Actual Yield for N	Kg ha <sup>-1</sup>			
	Very Low	Low	Medium	High
50	47.45	19.82		
60	66.52	38.94		
70	90.50	62.92	21.54	
80	122.84	95.25	53.87	
90	172.62	145.03	103.65	48.48

% From the Highest Actual Yield for K <sub>2</sub> O	Kg ha <sup>-1</sup>			
	Very Low	Low	Medium	High
50	87.69	67.41	36.99	
60	118.21	97.93	67.51	6.67
70	156.58	136.30	105.88	45.04
80	208.31	188.03	157.61	96.76
90	287.96	267.68	237.26	176.42

**Table 4. Amount spent on fertilizer (Peso) / yield production) of strawberry**

Treatments	The total amount of fertilizer materials spent*	Yield (t ha <sup>-1</sup> )	Amount spent on fertilizer (Peso) / yield production)
0.5	18,304.78	9.25	8.97
1.0	36,609.55	11.40	3.95
1.5	54,914.33	11.79	4.81
2.0	73,219.10	9.82	6.21
2.5	91,523.88	8.70	9.31

\*Prevailing market prices of the fertilizer materials as of December 2021: Diammonium phosphate- PhP 2,149.59; potassium chloride- PhP 1,621.49 and ammonium sulfate PhP 1,111.12 (PSA, 2022)

For K, the target yield for 50, 60, 70, 80 and 90% of the highest actual yield requires 87.69, 118.21, 156.58, 208.31 and 287.96 kg/ha<sup>-1</sup>, respectively for very low; 67.41, 97.93, 136.30, 188.03 and 267.68 kg/ha<sup>-1</sup>, respectively for low; 36.99, 67.51, 105.88, 157.61 and 237.26, respectively for medium K<sub>2</sub>O soil. There is no recommendation for 50% target yield if the soil had high K<sub>2</sub>O content. At 60% the soil K<sub>2</sub>O requirement is 6.67 kg/ha, 70% is 45.04 kg/ha, 80% is 96.76 kg/ha and 90% is 176.42 kg/ha.

The economics of judicious fertilizer application is important because fertilizer materials command a very high price in the market. In this study, diammonium phosphate (18-46-0),

potassium chloride (0-0-60), and ammonium sulfate (21-0-0) were used. Although the application of 225-225-360 kg ha<sup>-1</sup> of N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O, respectively showed a higher percentage of the maximum attainable yield, the amount spent for every kg of strawberry yield was higher compared to the recommendation set by Hafza [14] which was PhP 3.95 per kg of fruit yield. Half of Hafza's recommendation had a higher amount spent while increasing it further to 200 and 250% of the recommendation added production cost without economic benefits at all. Application of half of the fertilizer recommendation to strawberry plants had the lowest total spent on fertilizer materials. However, the amount spent per kg yield is higher

because of the lower yield. The data suggest that in determining the fertilizer rates, the yield should not be the sole basis for the recommendation. Considering the economic benefits derived from fertilizer application as well as the environmental impact is necessary for sound fertility management practices.

#### 4. CONCLUSION

The Mitscherlich-Bray Equation can be a basis for determining the amount of fertilizer to be applied along with the economic benefits of the recommendation. Strawberry is highly responsive to fertilizer application, particularly in N and K<sub>2</sub>O nutrients. Two fertilizer options were found very responsive as they reached 95 and 93% based on theoretical maximum yield. Considering economic benefits, the application of 150-150-240 kg ha<sup>-1</sup> of N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O, respectively still the best option. The equation is very useful in understanding the response of fertilizer to crops and thus it should be used in soil fertility trials.

#### 5. RECOMMENDATION

This study recommends separately conducting the study for N, P<sub>2</sub>O<sub>5</sub>, and K<sub>2</sub>O set-up. Also, the range of P should be lowered, and the ranges should be narrower. Since this study is a preliminary test of fertilizer recommendation, a verification trial must be conducted.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Reganold JP, Andrews PK, Reeve JR, Carpenter-Boogs L, Schadt CW. Fruit and Soil Quality of Organic and Conventional Strawberry Agroecosystems. PLOS ONE. 2010;5(10). DOI:10.1371/annotation/1eefda4-77af-4f48-98c32c5696ca9e7a. Available:https://doi.org/10.1371/annotation/1eefda4-77af-4f48-98c32c5696ca9e7a
2. Osip CA, Madriaga CS, Pimentel RL. Study on the propagation and production management of strawberry. Philippine Journal of Crop Science; 2008. Available:https://agris.fao.org/agris-search/search.do?recordID=PH2009000079
3. Ticbaen FD. Strawberry Industry in Benguet Province La Trinidad Experience. SGRA SEMINAR 21 “Sustaining the Growth and Gains of Research Development and Extension” at Ben Palispis Hall , Provincial Capitol; 2007.
4. PCCT. T'boli folks plan to start growing strawberries in South Cotabato; 2017. Available:https://medium.com/@rpconditionalcashtransfer/tboli-folks-plan-to-start-growing-strawberries-in-south-cotabato-24b30171c96f
5. Primer. Look: Benwa Farm in Davao City; 2017. Available:https://primer.com.ph/travel/2017/12/27/look-bemwa-farm-in-davao-city/
6. Sun Star. Strawberry urban garden in Davao City; 2019. Available:https://www.sunstar.com.ph/article/1820997/davao/business/strawberry-urban-garden-in-davao-city
7. Morgan L. Hydroponic Strawberry production, A technical guide to the hydroponic production of Strawberries. Suntec (NZ) Ltd, Tokomaru New Zealand. pp118. (PDF) Effect of Chemical, Organic and Bio Fertilization on Growth and Yield of Strawberry Plant; 2006. Available:https://www.researchgate.net/publication/328917238\_Effect\_of\_Chemical\_Organicand\_Bio\_Fertilization\_on\_Growth\_and\_Yield\_of\_Strawberry\_Plant [accessed Mar 09
8. Bibi AC, Oosterhuis DM, Gonias ED.. Exogenous application of putrescine ameliorates the effects of high temperature in strawberry's flowers and fruit development. J. Agron. Crop Sci. 2010; 196:205–211. Available:doi.org/10.1111/j.1439-037x.2009.00414.x
9. Almorado RC. Adaptability of Strawberry (*Fragaria ananassa* Duch) in Lanao Del Sur, Philippines. International Journal of Humanities and Social Sciences p-ISSN: 1694-2620 e-ISSN: 1694-2639. 2019;11(3):25-32, ©IJHSS Available:https://doi.org/10.26803/ijhss.11.3.4
10. Gokmen Yilmaz F, Korkmaz A, Isik Y, Saglik NC, Akcay Kulluk D, Kaya Y, Torun AR, Dursun N, Arisoy RZ, Partigoc F, Kaya Y. Determination of the Nitrate Sufficiency Limit Value for Nitrogenous Fertilization Use in Wheat Grown on a Calcareous Soils Through Application of Mitscherlich-Bray Equation. Communications in Soil Science and Plant Analysis. 2021;52(19): 2207-16.

11. PSA. Updates on Fertilizer Prices; 2022. Available:<https://psa.gov.ph/content/updates-fertilizer-prices-0>
12. Srivastava S, Subba Rao A, Alivelu K, Singh KN, Raju NS, Rathore A. Evaluation of crop responses to applied fertilizer phosphorus and derivation of optimum recommendations using the Mitscherlich–Bray equation. *Communications in soil science and plant analysis*. 2006;37(05-06):847-58.
13. Asio LG, dela Cruz NE. Application of Mitscherlich-Bray Equation to formulate fertilizer recommendation for Sweet Potato in Leyte, Philippines. *Annals of Tropical Research*; 2020. DOI:10.32945/atr4223.2020
14. Hafsa. Crop Guide: Strawberry Fertilizer Recommendation; 2008. Available:<https://www.haifa-group.com/crop-guide/vegetables/strawberry-fertilizer/crop-guide-strawberry-fertilizer-recommendations>
15. Ouano AC. Soil Quality Assessment in Mindanao State University. Unpublished Master's Thesis; 2018.
16. Preciado-Rangel P, Troyo-Diéguez E, Valdez-Aguilar LA, Garcia-Hernandez JL, Luna-Ortega JG. Interactive effects of the Potassium and Nitrogen Relationship on Yield and Quality of Strawberry Grown Under Soilless Conditions. *Plants*. 2020;9: 441. DOI:10.3390/plants 9040441
17. Medeiros RF, Pereira WE, Rodrigues RM, Nascimento R, Suassuna JF, Dantas TAG. Growth and yield of strawberry plants fertilized with nitrogen and phosphorus. *Rev. Bras. Eng. Agríc. Ambient*. 2015;9:865–870.
18. Agüero JJ, Salazar SM, Kirschbaum DS, Jerez EF. Factors affecting fruit quality in strawberries grown in a subtropical environment. *Int. J. Fruit Sci*. 2015;15:223–234.
19. USDA. Undated. Soil Phosphorus: Soil health Guide for Educators. Available:[https://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcs142p2\\_051878.pdf](https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_051878.pdf)
20. Ranganathan V. Baule Units in Augmenting Analysis of Nutrient Interactions. *Indian J Plant Soil*. 2019;6(1):37-39
21. The Philippines Recommends for Soil Diagnosis and Amelioration (PCARRD). Los Banos, Laguna; 2007.

© 2022 Jambaro et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/88680>