



The Effect of Different Concentrations of GA₃ and NAA on Growth and Flowering of Asiatic Lily (*Lilium × asiatica*) cv. Litouwen under Naturally Ventilated Polyhouse Conditions of Prayagraj

Kedar Mahadev Gheware^{a+++*}, Krishna Kaushik^{b+++*}, Rohit Gangwar^{b+++}, Mohd Wamiq^{c+++} and Urfi Fatmi^{d#}

^a *Division of Floriculture and Landscaping, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu, India.*

^b *Department of Floriculture and Landscape Architecture, Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut (U.P.), India.*

^c *Department of Vegetable Science, Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut (U.P.), India.*

^d *Department of Horticulture, SHUATS, Prayagraj, U.P., India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i121557

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95298>

Original Research Article

Received: 20/10/2022

Accepted: 27/12/2022

Published: 29/12/2022

⁺⁺ Ph.D. Student;

[#] Assistant Professor;

^{*}Corresponding author: E-mail: kedar.gheware.1997@gmail.com, koushikkishnaram@gmail.com;

ABSTRACT

This investigation was conducted under a naturally ventilated polyhouse, in the Department of Horticulture, SHUATS, Prayagraj, Uttar Pradesh. In December 2020 with records taken from the month of February to month of March 2021. There were nine treatments comprising two growth regulators Gibberellic acid (GA₃) (at 200, 250, 300, and 350 ppm), and Naphthalic Acetic Acid (NAA) (at 50,100,150, and 200 ppm) along with control. The whole experiment was laid out as a Randomized Block Design (RBD) with three replications. The results revealed that among the *Lilium* cv. LITOUWEN, the foliar application of 250 ppm GA₃ significantly increased the number of leaves per plant (66) with enhanced days to bud initiation (25.7 days). The number of bud per plant was 2.8 and stem length was 43.4cm respectively., GA₃ at a concentration of 350 ppm enhanced vase life (7.9 days). and NAA at a concentration of 100 ppm enhanced the number of bulblets (3.3).

Keywords: Asiatic lily; polyhouse; GA₃; NAA.

1. INTRODUCTION

"*Lilium* sp. is cultivated worldwide and is one of the most important generators of cut flowers, pot plants, and garden plants. The Asiatic hybrids are derived from interspecific crosses among at least 12 species of the Sinomartagon section" [1]. Lilies (*Lilium* species) are significant ornamental bulbous plants belonging to the family Liliaceae. The genus *Lilium* comprises nearly 100 species and more than 9,400 cultivars [2], an essential geophyte with showy flowers, appealing colour, sturdy growth, and robust flowers. which are divided into seven sections. Lily is the common English name for flowering plants of the *Lilium* genus and they are extensively grown in polyhouse as a cut flower in the global flower trade due to a wider choice of growing periods, adorn of colors, and everlasting quality. *Lilium* with showy captivating flowers of different color patterns. It is one of the important genera of flower bulbs produced worldwide. *Lilium* is a Latin name derived from the Greek word "leiron" used by the Theophrastus for the Madonna lily (*Lilium candidum*). In Christianity, the lily flower symbolizes the virgin mother in the semantic world, the symbol of mother wood. It can be grown both as a cut flower and a pot plant. Lily ranks fourth among the top ten cut flowers of the world, next to rose, chrysanthemum, and tulip [3]. The global production of lily bulbs occurs in ten countries [4], with the Netherlands having the maximum production area (4280 hectares, 77%), followed by France (401 hectares, 0.8%), Chile (205 hectares, 0.4%), Japan (189 hectares, 0.3%) and New Zealand (110 hectares, 0.2%). In India, lilies are famous among farmers of Haryana, Himachal Pradesh, Jammu Kashmir, Uttarakhand, and Tamil Nadu because of their high profitability within a short growing period of

four months. "The cultivars of the genus *Lilium* are highly appreciated by horticulturists for their outstanding range of color, fragrance and adaptability to several environmental conditions. *Lilium* can be used for informal planting in grassland or among orchard trees, along crocuses, bluebells and tulips to create flower meadows.

Large and attractive flowers with the capacity to rehydrate after long distance transportation, have made *Lilium* gain popularity fast in our country. The cultivars are highly appreciated for their outstanding range of colours, fragrances, and adaptability to several environmental conditions" [5]. "However, nearly all the cut flowers of *Lilium* available in the florists' shop are being acquired from Bengaluru, Pune and hilly areas of the country like Himachal Pradesh, Uttarakhand and North Eastern regions. However, Asiatic lily varieties/hybrids grown by amateurs in the state of Uttar Pradesh are coming up well and blooming in a wide range of colors, sizes, and shapes. Although agro-climatic conditions of the State are suitable for this flower crop, its commercial cultivation has not attracted the flower growers due to a lack of awareness about the crop, quality planting material and knowledge about its production technology [6].

Large and attractive flowers with the capacity to rehydrate after long-distance transportation, have made *Lilium* gain popularity fast in our country". [5]. "However, nearly all the cut flowers of *Lilium* available in the florists' shop are being acquired from Bengaluru, Pune and hilly areas of the country like Himachal Pradesh, Uttarakhand and North Eastern regions. However, Asiatic lily varieties/hybrids grown by amateurs in the state of Uttar Pradesh are coming up well and blooming in a wide range of colors, sizes and shapes. It is a species of great economic

importance in the production and commercialization of cut flower in the international market" [7].

Research work on the use of traditional plant growth promoters like gibberellins, improving bulb and corm multiplication rate as well as bulb and corm enlargement was carried out in different parts of the country but artificial application of plant growth regulators can also enhance, hasten, or delay the flowering time in some plant species. According to some other reports, gibberellins induced early flowering and prolonged flower life. Growth and yield were enhanced by application of Gibberellic Acid (GA₃) [8,9], and Naphthalic Acetic Acid (NAA) [10].

2. MATERIALS AND METHODS

The present investigation was conducted at Horticulture Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj from December 2020 to March 2021.

3. EXPERIMENTAL DESIGN AND TREATMENTS

An experiment was performed by using randomized block design (RBD) with three replications. Also, in this experiment plant was sprayed with Gibberellic Acid (GA₃) (at 200, 250, 300, 350 ppm), and Naphthalic Acetic Acid (NAA) (at 50, 100, 150, and 200 ppm) with control after planting of *Lilium* bulbs. The observations were recorded with respect to 15 days after planting to know the response of Asiatic lily to different regulators at different concentrations.

4. RESULTS AND DISCUSSION

No. of leaves per plant:- In no. of leaves per plant treatment differed significantly with respect. It was found that the maximum no. of leaves per plant (66) was reached in the T₂ GA₃ at 250 ppm. Gibberellic Acid (GA₃) influencing the increased vegetative growth by increasing cell division and cell elongation may be the reason behind the production of more number of leaves per plant. Similar findings were also reported by Pal and Das [11] where GA₃ application increased the number of leaves per plant in Asiatic lily which in consonance with the present findings.'

Days to bud initiation (days):- The results presented in Table 1 revealed that the growth regulator's treatments had significantly for days

to bud initiation. It was studied among the different treatments in T₂ GA₃ at 250 ppm that took lesser days (25.7 days), which were found maximum days (29.3 days) for bud initiation in T₀ Control. The result revealed that foliar application of 250 ppm GA₃ significantly advanced days to bud initiation in the Asiatic lily cultivar Litouwen. The reason behind this effect may be due to the stimulation and enhancement of vegetative growth. The present findings agree with Singh et al. [12].

Number of buds per plant:- The effect of different growth regulators showed significant influence on the no. of buds per plant. It was spotted that no. of bud per plant was more (2.8) in T₂ GA₃ at 250 ppm. Which had fewer bud per plant with NAA 200 ppm T₈. In this study, the number of flower buds per plant significantly increased with levels of GA₃ at 250 ppm. However, these parameters were recorded maximum in GA₃ at 250 ppm treatment, by the findings of Khan and Tiwari [13] in dahlia.

Stem length (cm):-The data on stem length exhibited significant differences among the treatments. The longest stem length was observed in T₂ GA₃ at 250 ppm (43.4 cm) and the Shortest stem length was seen in T₇ NAA at 150 ppm (15.7 cm). The results are in agreement with the report of Padmalatha et al. [14] recorded the effect of GA₃ on the growth and flowering of two gladiolus cultivars Darshan and Dhiraj and noted maximum spike length with GA₃.

The vase life of flower:- As mentioned in Table 1 plants treated with GA₃ at 350 ppm T₄ had longer vase life (7.9 days) whereas, the shorter vase life (2.8 days) was recorded in T₆ (NAA at 100 ppm) among all the treatments. The findings are in agreement with similarly, Kumar and Gupta [15] recorded significantly higher vase life with pre-soaking and foliar spray of GA₃ on gladiolus cv. Jessica. And Jadhav et al. [16] reported that pre-harvest foliar spray of GA₃ on gladiolus cv. American Beauty was significant in enhancing the vase's life.

Number of bulblets :-As mentioned in Table 1 among the different growth regulator treatments, the number of bulblets increased significantly with T₆ (NAA at 100 ppm, 3.2) whereas, fewer bulblets were reported in T₀ (Control, 0.9).The findings are in agreement with those of NAA accelerates the growth of the plant, stem cell and manufacturing root cells. Similarly, when NAA was used by Simmonds and Cumming [17], in *Lilium longiflorum* cultivars.

Table 1. The influence of GA₃ and NAA on No. of leaves per plant, Days to bud initiation (days), No. of bud per plant, Stem length (cm), Vase life, No. of bulblets

Treatment No.	Treatment	No. leaves per plant	Days to bud initiation (days)	No. of bud per Plant	Stem length (cm)	Vase Life	No. of bulblets
T ₀	Control	59	29.3	2.5	37.7	6.9	0.9
T ₁	GA ₃ at 200 ppm	63	27.7	2.7	40.5	7	1.8
T ₂	GA ₃ at 250 ppm	66	25.7	2.8	43.4	7.4	2.3
T ₃	GA ₃ at 300 ppm	63	27	2.1	41.2	6.7	1.7
T ₄	GA ₃ at 350 ppm	61	27.7	2.0	38.6	7.9	2.7
T ₅	NAA at 50 ppm	54	28	2.1	30.2	4.7	3.0
T ₆	NAA at 100ppm	47	26	1.9	16.5	2.8	3.3
T ₇	NAA at 150ppm	51	28	1.6	15.7	3	3.1
T ₈	NAA at 200ppm	46	27.7	1.1	16.3	3.4	2.6
S.Ed. (±)		2.74	0.75	0.45	3.06	0.67	0.66
C.D. at 5%		5.86	1.62	0.96	6.50	1.42	1.41




Fig. 1. No. of leaves per plant




Fig. 2. Days to bud initiation




Fig. 3. Number of bud per plant




Fig. 4. Stem length (cm)




Fig. 5. Vase life




Fig. 6. Number of bulblets

5. CONCLUSION

It is determined from the current study that among the different growth regulators and treatment GA₃ at 250 ppm gave significant results for No. of leaves per plant, days to bud initiation, no. of bud per plant, and stem length, while GA₃ at 350ppm gave better vase life of a flower in the Asiatic lily cultivar Litouwen. And NAA at 100 ppm gives significant results in no. of bulblets. Hence, GA₃ at 250 ppm can be recommended for plant growth and flowering of Asiatic lily cv. Litouwen. And NAA at 100 ppm for bulblets formation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Leslie AC. The international lily register (including supplements). The Royal Horticulture Society, London; 1982-2005. Available: <http://www.lilyregister.com/>
2. Fatmi U, Singh D, Bharti S. Growth and flowering of Asiatic lily cv. Pollyanna as influenced by different growing environments. Plant Archives. 2018;18(1): 760-762.
3. Chaudhary N, Sindhu SS, Kumar R, Saha TN, Raju DVS, Arora A, Sharma RR. Effect of growing media composition on growth, flowering, and bulb production of LA hybrid (Red Alert) and Oriental (Avocado) group of Liliium under protected condition; 2018.
4. Buschman JCM. Globalisation-flower-flower bulbs-bulb flowers. In IX International Symposium on Flower Bulbs. 2004;673:27-33.
5. Bahr LR, Compton ME. Competence for *In vitro* bulb regeneration among eight Liliium genotypes. Hort Science. 2004;39(1):127-129.
6. Comber H. A new classification of the genus Liliium. Lily Year Book, Royal Horticultural Society, London. 1949;13: 86-105.
7. Jimenez S, Plaza BM, Segura ML, Contreras JI and Lao TM. Peat substrate reuse in Liliium "Haveltia" crop. Communications in Soil Science and Plant Analysis. 2012;43:243-250.
8. Umrao VK, Singh RP, Singh AR. Effect of gibberellic acid and growing media on vegetative and floral attributes. Indian Journal of Horticulture. 2007;64: 73-76.
9. Rana P, Kumar J, Kumar M. Response of GA₃, plant spacing and planting depth on growth, flowering and corm production. Journal of Ornamental Horticulture. 2005;8 (1):41-44.
10. Kumar PS, Bhagawati R, Kumar R, Ronya T. Effect of plant growth regulators on vegetative growth, flowering and corm production of gladiolus. Journal of Ornamental Horticulture. 2008;11(4):265-270.
11. Pal AK, Das SN. Effect of growth regulators on growth and flowering of Liliium longiflorum. Orissa Journal of Horticulture. 1990;18(1-2): 18-21.
12. Singh J, Nigam R, Singh R, Kumar A, Kumar A. Effect of gibberellic acid and cycocel on growth, flowering and yield of chrysanthemum (*Dendranthema grandiflora ramat*) cv. birbalsahni. Journal of Pharmacognosy And Phytochemistry. 2018;1:2753-2758.

13. Khan FU, Tiwari GN. Effect of growth regulators on growth and flowering of dahlia (*Dahlia variabilis* L.). Indian Journal of Horticulture. 2003;60(2):192-194.
14. Padmalatha T, Reddy GS, Chandrasekhar R, Shankar AS, Chaturvedi A. Effect of foliar sprays of bio regulators on growth and flowering in gladiolus. Indian Journal of Agricultural Research. 2013;47(3): 192-199.
15. Kumar S, Gupta AK. Postharvest life of *Gladiolus grandiflorus* L. cv. Jessica as influenced by pre-harvest application of gibberellic acid and kinetin. Journal of Postharvest Technology. 2014;2(3):169-176.
16. Jadhav PB, Mangave BD, Singh A, Dekhane SS, Patel DJ. Effect of plant growth regulators and natural growth substances on growth and flowering of gladiolus cv. American Beauty. International Journal of Current Research. 2015;7:17674-17676.
17. Simmonds JW, Cumming BG. Propagation of *Lilium* hybrids dependence of bulblet production on time of scale removal and growth substances. Scientia Horticulturae. 1976;5:77-83.

© 2022 Gheware et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95298>