



Effect of Macro and Micronutrients on Nutrient Uptake of Groundnut (*Arachis hypogaea* L.) in Coastal Sandy Soils

K. Ramanathan ^{a^o*}, P. Venkata Subbaiah ^{b[#]},
P. Mohana Rao ^{a[†]} and K. Lakshman ^{c[‡]}

^a Department of Soil Science and Agricultural Chemistry, Agricultural College, Bapatla, Andhra Pradesh, India.

^b Saline Water Scheme, Agricultural College Farm, Bapatla, Andhra Pradesh, India.

^c Krishi Vigyan Kendra, Banavasi, Kurnool, Andhra Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i242693

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/90371>

Original Research Article

Received: 13/06/2022
Accepted: 17/08/2022
Published: 30/12/2022

ABSTRACT

Aim: This experiment was conducted to evaluate the effect of macro and micronutrients on nutrient uptake of groundnut in coastal sandy soils.

Materials and Methods: A field experiment was conducted at Agricultural College Farm, Bapatla during *rabi*, 2021-22 to assess the response of macro and micronutrients on nutrient uptake of groundnut in coastal sandy soil.

Results: The results of the experiment clearly indicated that the nutrient uptake of macro, secondary and micronutrients was significantly influenced by the application of macro and micronutrients. The treatment T₅ (125% RDF + Soil application of ZnSO₄ @ 50 kg ha⁻¹ and Borax

^oPG Scholar;

[#]Senior Scientist;

[†]Assistant Professor;

[‡]Scientist;

*Corresponding author: E-mail: ramanathankalaivanan@gmail.com;

@ 10 kg ha⁻¹) recorded significantly higher uptake of N, P, K, S, Fe, Zn, Cu, Mn and B at peg penetration stage. At pod development stage except iron, the highest values of N, P, K, S, Zn, Cu, Mn and B uptake were recorded in the treatment T₅ and it was on par with the treatment T₇. The treatment T₇ which received 125% RDF + Foliar application of ZnSO₄ @ 2 g L⁻¹, FeSO₄ @ 5 g L⁻¹ and Borax @ 1.5 g L⁻¹ at 45 and 65 DAS registered the highest uptake of N, P, K, S, Zn, Cu, Mn and B but on par with T₅ at harvest stage.

Conclusion: For iron uptake at pod development and harvest stage, the treatment T₇ obtained significantly higher values and it was significantly superior over all other treatments.

Keywords: Groundnut; micronutrients; uptake; zinc; boron; iron.

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is one of the important oilseed crops and occupies an important position in the Indian agricultural economy. It is capable of fixing atmospheric nitrogen by the bacteria *Rhizobium* present in the nodules and hence it requires less N containing fertilizers. It contains high oil and protein content hence it is called as the king of oil seed crops. India ranks first in groundnut area with an area of 4.89 million hectares and second in production with 10.10 million tonnes. It is a rich source of edible oil and high quality protein and hence it is valued for both oil and confectionary purposes. Groundnut is mainly grown for its seed but all parts of the plant are utilized [1].

Groundnut is mostly grown in light textured soils especially sandy loam and sandy soils mainly because it has underground pod bearing habit. But these soils have poor nutrient status especially micronutrients, zinc (Zn) and boron (B) due to leaching, low nutrient retention capacity and low organic matter status [2]. Restricted availability of these nutrients in sandy soil greatly impairs the yield of groundnut. Judicious use of fertilizer is an important management practice to increase groundnut production. Balanced use of fertilizers assumes vital important in sustainable agriculture. In these soils, the nutrient uptake will also be low, if unfertilized. Improvement in uptake of macro and micronutrients by the application of micronutrients has been reported by several workers. Keeping all these points in view, this experiment was conducted to evaluate the effect of macro and micronutrients on nutrient uptake of groundnut in coastal sandy soils.

2. MATERIALS AND METHODS

The experiment was carried out during *rabi*, 2021-22 at Agricultural College Farm, Bapatla in RBD with seven treatments replicated thrice. The experimental soil was sandy in texture, neutral in

reaction (6.65), non-saline (0.09 dS m⁻¹), low in organic carbon (0.09 g kg⁻¹), available nitrogen (135 kg ha⁻¹), medium in available phosphorus (39.5 kg ha⁻¹) and low in available potassium (118 kg ha⁻¹), calcium (320 mg kg⁻¹) and magnesium (42 mg kg⁻¹) and sufficient in sulphur (20 mg kg⁻¹), manganese (3.59 mg kg⁻¹), copper (0.69 mg kg⁻¹) and deficient in boron (0.29 mg kg⁻¹), iron (3.95 mg kg⁻¹) and zinc (0.29 mg kg⁻¹). The treatments are T₁- Control, T₂- 100% RDF, T₃- 125% RDF, T₄- 100% RDF + Soil application of ZnSO₄ @ 50 kg ha⁻¹ and Borax @ 10 kg ha⁻¹, T₅- 125% RDF + Soil application of ZnSO₄ @ 50 kg ha⁻¹ and Borax @ 10 kg ha⁻¹, T₆- 100% RDF + Foliar application of ZnSO₄ @ 2 g L⁻¹, FeSO₄ @ 5 g L⁻¹ and Borax @ 1.5 g L⁻¹ at 45 and 65 DAS, T₇- 125% RDF + Foliar application of ZnSO₄ @ 2 g L⁻¹, FeSO₄ @ 5 g L⁻¹ and Borax @ 1.5 g L⁻¹ at 45 and 65 DAS. A common dose of 30 kg nitrogen ha⁻¹, was applied through urea in two equal split doses, half as basal, and a half at 30 DAS by considering the plot size. A common dose of phosphorus @ 40 kg ha⁻¹ in the form of single super phosphate, and potassium @ 50 kg ha⁻¹ in the form of muriate of potash were applied as basal before sowing. ZnSO₄ and borax were applied at the rate of 50 kg ha⁻¹ and 10 kg ha⁻¹ respectively, to the plots as per the treatments as basal and foliar application of ZnSO₄, FeSO₄ and borax were applied at the rate of 2 g L⁻¹, 5 g L⁻¹ and 1.5 g L⁻¹ at 45 DAS and 65 DAS to the respective plots as per the treatments.

The groundnut variety TAG-24 was planted in the second week of November with a spacing of 30 x 10 cm. The crop was raised with all the standard packages of practices as they required. Plant samples were collected at peg penetration, pod development and harvest stages and analyzed using standard procedures in the laboratory. Five representative plant samples were collected, shade dried and kept in oven at 70^o C for 24 to 48 hours till the constant weight and then it was averaged to get data in g/ plant, then calculated on hectare basis. Plant uptakes were worked out by using nutrient content and dry matter

accumulation. The data were analyzed statistically by following the analysis of variance (ANOVA) technique as suggested by Panse and Sukhatme (1978) for RBD.

Macronutrient uptake (kg ha^{-1}) = (Nutrient concentration (%) \times Dry matter yield (kg ha^{-1}))/100

Micronutrient uptake (g ha^{-1}) = (Nutrient concentration (mg kg^{-1}) \times Dry matter yield (kg ha^{-1})) / 1000

3. RESULTS AND DISCUSSION

3.1 Macronutrients Uptake

Application of macro and micronutrient had a significant influence on macronutrients uptake by the groundnut at all the stages of the crop (Table 1). The highest N (52.61 kg ha^{-1}), P (8.59 kg ha^{-1}) and K (41.5 kg ha^{-1}) uptakes were recorded by the treatments T₅ (125% RDF + Soil application of ZnSO₄ @ 50 kg ha^{-1} and Borax @ 10 kg ha^{-1}) at peg penetration stage. Whereas at pod development stage, the treatment T₅ recorded the highest uptake of N (61.45 kg ha^{-1}), P (11.47 kg ha^{-1}) and K (46.3 kg ha^{-1}) but it was on par with T₇. In haulm and kernel at harvest stage, the highest N (30.95 and 92.95 kg ha^{-1}), P (4.22 and 8.94 kg ha^{-1}) and K (33.1 and 22.8 kg ha^{-1}) uptakes respectively, were recorded by the treatment T₇ which received 125% RDF + Foliar application of Zn, Fe and B at 45 and 65 DAS) and it was on par with T₅. The lowest uptake of all macronutrients (N, P and K) was recorded in the control (T₁) at all the stages. Yakadri and Satyanarayana [3] reported that there is a close relationship between nutrient uptake and dry matter production in groundnut. The increased uptake of nitrogen was mainly due to the fact that the micronutrients like zinc and boron are involved in nitrogen fixation and translocation into plant parts, which might have led to higher dry matter production. The higher uptake might be due to the solubilization of native phosphorus in addition to applied fertilizers which ultimately resulted in better root growth and increased physiological activity of roots to absorb more phosphorus (Sudarasan and Ramaswami [4]; Sumangala [5]). Increased K uptake might be due to better plant growth leading to higher uptake of nutrients and further on the stimulatory effect of B and Zn in absorption of potassium. Such findings were in accordance in Elayaraja and Singaravel [6], Abd EL-Kader and Mona [7] and Abhigna [8].

3.2 Secondary and Micronutrients Uptake

Perusal of data revealed that there was a significant influence of secondary and micronutrients uptake at all the stages by the macro and micronutrient application (Tables 2 and 3). Maximum sulphur uptake (7.07 kg ha^{-1}) was recorded with the treatment T₅ at peg penetration stage. At pod development stage, the treatment T₅ recorded maximum sulphur uptake (10.09 kg ha^{-1}) and it was on par with T₇. Whereas at harvest stage, the treatment T₇ obtained the highest uptake in both haulm (6.06 kg ha^{-1}) and kernel (5.69 kg ha^{-1}) which was on par with T₅ (Table 2). The higher sulphur uptake might also be due to stimulatory effect of zinc and boron uptake.

Highest iron (393 g ha^{-1}) uptake was recorded with the treatment T₅ (125% RDF + Soil application of ZnSO₄ @ 50 kg ha^{-1} and Borax @ 10 kg ha^{-1}) at peg penetration stage. At pod development and haulm and kernel at harvest stages, highest iron uptake of 505 , 284 and 224 g ha^{-1} were recorded with the treatment T₇ (125% RDF + Foliar application of ZnSO₄ @ 2 g L^{-1} , FeSO₄ @ 5 g L^{-1} and Borax @ 1.5 g L^{-1} at 45 and 65 DAS) and it was superior over all other treatments.

At peg penetration stage, maximum Zn (119.3 g ha^{-1}), Cu (27.00 g ha^{-1}), Mn (116.0 g ha^{-1}) and B (90.9 g ha^{-1}) uptake was recorded with the treatment T₅. The treatment T₅ recorded significantly higher uptake of Zn (147.0 g ha^{-1}), Cu (36.27 g ha^{-1}), Mn (135.9 g ha^{-1}) and B (142.4 g ha^{-1}) at pod development stage and it was on par with T₇. At harvest stage, the treatment T₇ obtained the maximum uptake of Zn (84.8 and 57.9 g ha^{-1}), Cu (24.48 and 19.34 g ha^{-1}), Mn (91.2 and 57.8 g ha^{-1}) and B (95.9 and 70.6 g ha^{-1}) in both haulm and kernel respectively and this was statistically on par with T₅. The lowest uptake of all micronutrients at all the stages was recorded by the treatment T₁ (Control).

Both soil and foliar application of micronutrients led to an increase in the concentrations of macro and micronutrients in seeds due to the vital physiological roles in the plants which promotes the uptake of plant nutrients. Increase in micronutrients uptake might also be due to higher dry matter production coupled with increased content. Also, a positive interaction between iron, boron and zinc was also reported [9]. The results are in accordance with Gowthami and Ananda [10], Aboyeji et al. [11], Elayaraja and Senthilvalavan [12] and Kamalakannan and Elayaraja [13].

Table 1. Effect of macro and micronutrients on N, P and K uptake by groundnut (PP-Peg penetration stage, PD- Pod development Stage)

Treatments	Nitrogen uptake (kg ha ⁻¹)				Phosphorus uptake (kg ha ⁻¹)				Potassium uptake (kg ha ⁻¹)			
	PP	PD	Harvest		PP	PD	Harvest		PP	PD	Harvest	
			Haulm	Kernel			Haulm	Kernel			Haulm	Kernel
T ₁ : Control	23.70	27.26	12.12	35.69	4.00	5.41	1.70	3.51	19.2	21.2	14.0	8.8
T ₂ : 100% RDF	32.06	36.95	17.06	53.41	5.38	7.24	2.44	5.32	25.5	28.4	19.3	12.9
T ₃ : 125% RDF	42.33	48.48	23.52	72.63	7.12	9.36	3.28	7.15	33.1	36.8	25.5	17.9
T ₄ : 100% RDF + Soil application of ZnSO ₄ @ 50kg ha ⁻¹ and Borax @ 10 kg ha ⁻¹	41.42	47.90	22.59	71.78	7.06	9.21	3.30	7.00	33.0	36.5	25.4	17.7
T ₅ : 125% RDF + Soil application of ZnSO ₄ @ 50 kg ha ⁻¹ and Borax @ 10 kg ha ⁻¹	52.61	61.45	30.45	92.13	8.59	11.47	4.08	8.83	41.5	46.3	32.9	22.7
T ₆ : 100% RDF + Foliar application of ZnSO ₄ @ 2 g L ⁻¹ , FeSO ₄ @ 5 g L ⁻¹ and Borax @ 1.5 g L ⁻¹ at 45 and 65 DAS	32.89	47.41	22.88	72.05	5.52	9.18	3.24	7.08	26.0	36.2	25.4	17.8
T ₇ : 125% RDF + Foliar application of ZnSO ₄ @ 2 g L ⁻¹ , FeSO ₄ @ 5 g L ⁻¹ and Borax @ 1.5 g L ⁻¹ at 45 and 65 DAS	42.33	60.52	30.95	92.95	7.03	11.46	4.22	8.94	33.0	46.0	33.1	22.8
SEm (±)	1.71	2.72	1.11	3.74	0.24	0.39	0.14	0.38	1.86	2.20	0.99	0.81
CD (p=0.05)	5.27	8.38	3.43	11.53	0.74	1.22	0.42	1.17	5.73	6.77	3.06	2.50
CV (%)	7.76	9.99	8.46	9.25	6.52	7.55	7.45	9.63	10.67	10.59	6.85	8.16

Table 2. Effect of macro and micronutrients on S, Fe and Zn uptake by groundnut (PP-Peg Penetration Stage, PD-Pod Development Stage)

Treatments	Sulphur uptake (kg ha ⁻¹)				Iron uptake (g ha ⁻¹)				Zinc uptake (g ha ⁻¹)			
	PP	PD	Harvest		PP	PD	Harvest		PP	PD	Harvest	
			Haulm	Kernel			Haulm	Kernel			Haulm	Kernel
T ₁ : Control	3.64	4.99	2.73	2.43	210	241	115	86	58.5	66.7	35.7	22.3
T ₂ : 100% RDF	4.50	6.23	3.56	3.41	255	282	139	114	71.4	82.8	45.4	30.7
T ₃ : 125% RDF	5.34	7.38	4.38	4.16	303	325	163	136	83.4	97.6	54.5	37.1
T ₄ : 100% RDF + Soil application of ZnSO ₄ @ 50kg ha ⁻¹ and Borax @ 10 kg ha ⁻¹	6.15	8.56	5.15	4.86	347	369	201	157	107.0	130.2	72.3	48.6
T ₅ : 125% RDF + Soil application of ZnSO ₄ @ 50 kg ha ⁻¹ and Borax @ 10 kg ha ⁻¹	7.07	10.09	6.04	5.64	393	411	227	178	119.3	147.0	83.8	56.7
T ₆ : 100% RDF + Foliar application of ZnSO ₄ @ 2 g L ⁻¹ , FeSO ₄ @ 5 g L ⁻¹ and Borax @ 1.5 g L ⁻¹ at 45 and 65 DAS	4.59	8.47	5.00	4.90	258	451	249	197	71.9	129.9	72.5	49.3
T ₇ : 125% RDF + Foliar application of ZnSO ₄ @ 2 g L ⁻¹ , FeSO ₄ @ 5 g L ⁻¹ and Borax @ 1.5 g L ⁻¹ at 45 and 65 DAS	5.44	9.76	6.06	5.69	303	505	284	224	83.3	145.3	84.8	57.9
SEm (±)	0.25	0.37	0.26	0.19	13.84	12.98	7.12	6.23	3.82	4.39	2.88	1.90
CD (p=0.05)	0.76	1.13	0.79	0.60	42.65	39.99	21.93	19.19	11.77	13.54	8.87	5.85
CV (%)	8.09	7.98	9.41	7.59	8.11	6.09	6.26	6.92	7.79	6.66	7.78	7.60

Table 3. Effect of macro and micronutrients on Cu, Mn and B uptake by groundnut (PP-Peg Penetration Stage, PD-Pod development Stage)

Treatments	Copper uptake (g ha ⁻¹)				Manganese uptake (g ha ⁻¹)				Boron uptake (g ha ⁻¹)			
	PP	PD	Harvest		PP	PD	Harvest		PP	PD	Harvest	
			Haulm	Kernel			Haulm	Kernel			Haulm	Kernel
T ₁ : Control	15.44	18.67	11.09	8.48	60.7	68.3	40.8	24.4	34.5	54.1	30.6	21.6
T ₂ : 100% RDF	18.19	22.54	14.05	11.43	73.7	83.4	51.1	35.4	42.3	68.9	39.8	31.5
T ₃ : 125% RDF	21.18	26.58	16.69	13.81	87.0	97.6	61.3	43.0	51.0	83.5	49.1	39.1
T ₄ : 100% RDF + Soil application of ZnSO ₄ @ 50kg ha ⁻¹ and Borax @ 10 kg ha ⁻¹	24.30	31.49	20.76	16.29	103.5	117.9	76.8	49.9	81.4	126.8	78.9	57.8
T ₅ : 125% RDF + Soil application of ZnSO ₄ @ 50 kg ha ⁻¹ and Borax @ 10 kg ha ⁻¹	27.00	36.27	24.00	19.19	116.0	135.9	88.3	57.3	90.9	142.4	94.2	70.3
T ₆ : 100% RDF + Foliar application of ZnSO ₄ @ 2 g L ⁻¹ , FeSO ₄ @ 5 g L ⁻¹ and Borax @ 1.5 g L ⁻¹ at 45 and 65 DAS	18.33	31.04	20.96	16.72	74.2	116.7	76.2	50.1	43.2	126.4	80.6	58.7
T ₇ : 125% RDF + Foliar application of ZnSO ₄ @ 2 g L ⁻¹ , FeSO ₄ @ 5 g L ⁻¹ and Borax @ 1.5 g L ⁻¹ at 45 and 65 DAS	21.35	35.89	24.48	19.34	87.2	134.0	91.2	57.8	51.3	141.4	95.9	70.6
SEm (±)	0.85	1.11	0.81	0.77	3.87	4.51	3.27	1.91	2.29	4.70	2.62	2.33
CD (p=0.05)	2.63	3.41	2.50	2.36	11.94	13.88	10.08	5.89	7.06	14.50	8.07	7.17
CV (%)	7.08	6.62	7.44	8.84	7.80	7.25	8.17	7.29	7.04	7.67	6.77	8.08

4. CONCLUSION

From the analysis of experimental data, it could be concluded that the foliar application of zinc, iron and boron along with 125% RDF improved the uptake of all the nutrients in coastal sandy soils [14].

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh BB, Musa A, Ajeigbe HA, Tarawali SA. Effect of feeding crop residues of different cereals and legumes on weight gain of Yankassa rams. *Int J Livest Prod.* 2012;2:17-23.
2. Elayaraja D, Singaravel R. Zinc and boron application on groundnut yield and nutrient uptake in coastal sandy soils. *Int Res J Chem.* 2016;12:17-23.
3. Yakadri M, Satyanarayana V. Dry matter production and uptake of nitrogen, phosphorus and potassium in rainfed groundnut. *Indian J Agron.* 1995;40(2):325-7.
4. Sudarasan S, Ramaswami PP. Micronutrient nutrition in groundnut black gram cropping system. *Fert News.* 1993;38(2):51-7.
5. Sumangala BJ. Response of groundnut (*Arachis hypogaea* L.) to conjunctive use of micronutrients and bio-inoculants at graded levels of fertilizers under dryland conditions [Ph.D. thesis]. Bengaluru, Karnataka: University of Agricultural Sciences; 2003.
6. Elayaraja D, Singaravel R. Zinc and boron application on groundnut yield and nutrient uptake in coastal sandy soils. *An Asian J Soil Sci.* 2012;7(1):50-3.
7. EL-Kader A, Mona G. Effect of sulfur application and foliar spraying with zinc and boron on yield, yield components and seed quality of peanut (*Arachis hypogaea* L.). *Res J Agric Biol Sci.* 2013;9(4):127-35.
8. Abhigna D. Effect of soil and foliar application of micronutrients on growth and productivity of groundnut in sandy soils. M.Sc. (Ag.) [thesis]. Bapatla: ANGRAU. Andhra Pradesh; 2021.
9. Haneena KM, Subbaiah PV, Rao ChS, Srinivasulu K. Effect of boron on nutritional quality of groundnut grown in coastal sandy soils. *Int J Plant Soil Sci.* 2021;33(19):189-97.
10. Gowthami VSS, Ananda N. Effect of zinc and iron ferti-fortification on growth, pod yield, and zinc uptake of groundnut (*Arachis hypogaea* L.) genotypes. *Int J Environ Agric Biotechnol.* 2017;10(5):575-80.
11. Aboyeji C, Dunsin O, Adekiya AO, Chinedum C, Suleiman KO, Okunlola FO et al. Zinc sulphate and boron-based foliar fertilizer effect on growth, yield, minerals, and heavy metal composition of groundnut (*Arachis hypogaea* L.) grown on an alfisol. *Int J Agron.* 2019;2019:1-7.
12. Elayaraja D, Senthilvalavan P. Soil properties, enzymatic activity, yield and nutrient uptake of groundnut as influenced by nutrient management practices in coastal sandy soil. *Annals Plant Soil Res.* 2019;21(1):87-92.
13. Kamalakannan P, Elayaraja D. Effect of organic and inorganic sources of nutrients in micronutrients uptake and availability on groundnut in sandy clay loam soil. *Plant Arch.* 2020;20(1):3721-6.
14. Panse VG, Sukhatme. *Statistical methods for agricultural workers.* 3rd ed. Indian Council of Agricultural Research Publication. New Delhi. 1978;361.

© 2022 Ramanathan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/90371>