



Assessment of Growth Pattern and Intervention's Effects of Oilseeds Mission on Groundnut Production in Andhra Pradesh, India

Sushmitha Burigi ^{a*}, K. N. Selvaraj ^b, R. Senthil Kumar ^{a, b},
S. Senthilnathan ^b, S. Moghana Lavanya ^b
and U. Arulanandu ^a

^a Anbil Dharmalingam Agricultural College and Research Institute, Trichirapalli, India.

^b Tamil Nadu Agricultural University, Coimbatore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2023/v45i72136

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/99705>

Original Research Article

Received: 01/03/2023

Accepted: 03/05/2023

Published: 09/05/2023

ABSTRACT

High instability in crop area and yield resulted in production shortfall in Andhra Pradesh. As groundnut is raised mostly as rainfed kharif crop, yield during this season is half of the yield attained during rabi/summer season. State's average productivity is also lower than the national average necessitating the importance of improving the oilseeds production in the state since the state contributing 8 % to India's groundnut production. Oilseeds mission was implemented in all the districts of the state to scale up the production through various technological interventions. Therefore, understanding the pattern of production before and after the intervention is critically

*Corresponding author: E-mail: chummi1993@gmail.com;

important for future directions and strategies required to scale up the groundnut production in the state. The present paper aims at assessing the district-wise growth performance of groundnut and examining the sustained effects of oilseeds mission. Using the time series data on area, production, and productivity (1997-98 and 2019-20), growth analysis and interrupted time series segmented regression analyses were employed. Groundnut area tended to decline in all the districts of the state. The rate of decline was higher in Cuddapah (7.47%) followed by Kurnool (3.82%), Chittoor (3.79%) and Anantapur (2.13%) and the state recorded a negative growth rate of 3.49% per annum. Production growth was found low in Anantapur district (1.39%) though the yield registered a 3.63% growth per annum. Similarly, production in Chittoor district declined by 2.1% per annum though yield grew by 4% per annum due to higher fall in area. The Cuddapah also witnessed a negative growth in production and production decline was around 2% per annum. However, productivity in the Cuddapah district grew at faster rate compare to other major districts and the productivity has grown at the rate 6% per annum. Groundnut production in the state declined by 0.4% per annum due to decline in both productivity and area by 3% and 3.4% respectively. It is alarming to note that the maximum decline in production was 12.38% in the State mainly due to the maximum decline in production in major groundnut producing districts namely Anantapur (76%) followed by Kurnool (32.32%), Cuddapah (22.19%) and Chittoor (13.40%). The model results of interrupted time series show that though there is no significant intervention effect, there is sign of productivity improvement after the intervention. The coefficient of sustained effect for yield turned to positive from negative though not significant (immediate effect) for the major growing districts, indicating the long-term effect of the oilseeds mission. Constraints relating to adoption of improved technology, non-suitability of improved technology, and lack of technical guidance were identified by studies affecting groundnut production in the state. Therefore, addressing these issues through policy and technological interventions are crucial for sustaining the productivity and increasing the production in the major districts.

Keywords: Growth pattern; groundnut production; oilseeds; rainfed kharif crop.

1. INTRODUCTION

Andhra Pradesh is second major producer of groundnut in India sharing 8% of the total country's production. In Andhra Pradesh groundnut is grown in Anantapur, Kurnool, Chittoor, Cuddapah, Warangal, Nalgonda, Srikakulam, Visakhapatnam and Mahaboobnagar districts. Groundnut is raised mostly as rainfed kharif crop; nearly 90% of acreage and production comes from kharif crop (June-October) [1] and less than 7% of the area under groundnut is irrigated [2]. As major area is under rainfed, groundnut productivity is erratic and crop size also shrinks. Apart from erratic rainfall, yield potential of groundnut has not been touched till date due to biotic, abiotic, technological, institutional, and socio-economic constraints [3]. Farmers in the major rainfed growing districts of the state are shifting from groundnut to other remunerative crops particularly towards oil palm cultivation due to the incentives provided by the state government (www.horticulture.ap.nic.in). Like announcement of remunerative price, which resulted in decreasing share of groundnut in total oilseeds production. Farmers have also been cultivating redgram and other commercial crops along with groundnut particularly during the

kharif season resulting in lesser farming of groundnuts. Increase in cost of production due to rising input prices, lack of availability of good quality inputs, shortage of subsidised seeds [4], though the government is supplying the seed on subsidy for varietal and to enhance the seed replacement rate of groundnut [5], and insufficient extension services affected groundnut production in the state.

State's average productivity of groundnut continues to be lower than the national average. Oilseeds mission was implemented in all the districts of the state to scale up the production through various technological interventions. Though the implementation of technology mission on oilseeds paved the way for increase in oil seeds productions in the State by increasing the productivity through technological intervention, variability in area under groundnut leads to variability in productivity resulting in inconsistent supply. Instability in crop area and yield of groundnut resulted in production shortfall in most of the districts. Hence, it is vital to understand the performance of groundnut in major districts in Andhra Pradesh in order to identify the determinants of performance and potentials for future expansion of area under

groundnut through the implementation of strategies for scaling -up the productivity of groundnut-based production systems. The present paper aims at assessing the growth performance of groundnut in various districts of Andhra Pradesh and examining the sustained effects of oilseeds mission on groundnut production and identifying the potentials for area increase and productivity improvement.

2. MATERIALS AND METHODS

(i) Growth Performance of Groundnut

Time series data on area, production, and productivity of groundnut for all the 13 districts of Andhra Pradesh for the period between 1997-98 and 2019-20 were sourced from various online publications (aps.dac.gov.in). Compound growth rate was estimated for overall period using the formula $[(\text{Ending period (2019-20)}/\text{Initial value (1997-98)})^{1/23}-1]*100]$ and for maximum decline and increase in area/ production and productivity in the state and districts in terms of compound growth rate was estimated using the formula $[(\text{Min value}/\text{Max value})^{1/\text{No. of years}}-1]*100]$. The growth rates were estimated in excel worksheet and the districts were classified based on compound growth rates.

(ii) Intervention's Effect: Tracking a Long-term Period before and after a Point of Intervention - Quasi Experimental Time Series Analysis

The government of India implemented technology mission on oilseed and oil palm (TMOP) during the period 1985-2003 (7th to 9th plan), integrated scheme on oilseed, pulses, oil palm and maize (ISPOM) during 2004-2013 (10th to 11th plan) and national mission on oilseed and oil palm (NMOOP) during the period 2014-2017 (12th plan) in order to increase oilseed production since a substantial portion of edible oil requirement is met through imports. In order to capture the impact, the above set programmes and classify the districts based on the potential the following interrupted time series analysis is employed.

$$y = \alpha + \beta_1 T + \beta_2 X + \beta_3 XT + \epsilon.$$

Where;

T refers to time period, X indicates the study phase and XT represents time after interruption. β_1 coefficient indicate pre-trend, β_2 shows the

post level change, β_3 is the post trend change and $(\beta_1 + \beta_3)$ is the post-trend.

The interrupted time series enable to investigate no impact, immediate impact, sustained or long-term impact and both immediate and sustained impacts of oilseed mission on area, production, and productivity. In time series it is important to understand the counterfactual and the actuals. The analysis involves tracking a long-term period before and after a point of intervention to assess the intervention's effects. The time coefficient β_1 indicates the trends in area, production, and productivity of groundnut in before intervention. The positive and significant coefficients indicate that area, production, and productivity increase over the period or vice versa. The β_2 normally indicates the immediate effect of oilseed mission and the immediate effect is expected to be positive and significance. Time since intervention impact co-efficient β_3 indicates that trend has changed after the intervention i.e., the sustained effect of intervention.

3. RESULTS AND DISCUSSION

(i) Growth Performance of Groundnut in Various Districts of Andhra

Cultivation of groundnut in Andhra Pradesh is completely dependent on rains as this crop mostly grown in rainfed conditions. Area under the groundnut in all the districts of Andhra Pradesh are declining at faster rate and the rate of decline is higher in the major groundnut growing districts namely Anantapur (2.13%), Chittoor (3.79%), Cuddapah (7.47%) and Kurnool (3.82 %). The area under groundnut in the State also registered a negative rate of growth and the area declined at the rate of 3.49 % between 1997-98 and 2019-20 and maximum decline was 6 % per annum, which is alarming (Table 1). Anantapur district contributes on an average 57 % of the state's area and maximum share in area was 66 %. Since, the maximum decline in this district was 5.12 %, the production growth is found low (1.39%) despite the productivity increased by 3.63 % per annum.

Similarly, the rate of production in Chittoor district declined at 2.1% overall, though the productivity increased at the rate of 4 % and this is mainly due to a decline in area by almost 4 % per annum. The Cuddapah, one of the major groundnuts producing districts in the state, also witnessed a negative growth rate in production

and production decline was around 2% per annum. However, productivity in the Cuddapah district grew at faster rate compare to other major districts and the productivity has grown at 6% per annum (Table 2).

The rate of growth in production in Kurnool district, in which groundnut occupies 14% area in the state, also witnessed very meager rate of growth in production (3.37%) due to an alarming rate of decline in area by 9.2% per annum and the overall rate of decline growth was estimated at 4% per annum. The rate of increasing in productivity was 4.34% per annum in this district and this rate of increase in productivity could be

not able to push up the production due to alarming rate of decline in area. As part of the Andhra Pradesh Primary Sector Mission (APPSM), which is also known as 'Rythu Kosam', farmers in the district were advised through text messages on the best time of sowing and harvest depending on weather conditions, soil and other indicators. Farmers who sowed in the first week of June got meagre yields due to a long dry spell in August, while registered farmers who sowed in the last week of June and first week of July and followed advisories got better yields and are out of loss. Though there is a distinct example of technological success as the farmers in Andhra Pradesh's Kurnool district




Fig. 1. Trend pattern of yield of ground nut in Andhra Pradesh and period of intervention

Table 1. Growth of area of peanuts in various districts of Andhra Pradesh during the period between 1998 and 2020

District	Area – Average Share (%)	Area – share – Max (%)	Area – Share – Min (%)	CGR (%) – Area (ha)	Max decline/increase in area-CGR (%)
Anantapur	56.58	66.38	44.79	-2.13	-5.12
Chittoor	13.51	17.63	9.12	-3.79	-4.13
East Godavari	0.04	0.14	0.02	-10.81	-11.64
Guntur	0.40	0.60	0.26	-1.73	-6.76
Cuddapah	8.09	13.30	2.64	-7.47	-14.82
Krishna	0.45	1.03	0.16	-9.68	-9.67
Kurnool	14.06	18.24	8.92	-3.82	-9.20
Prakasam	0.66	1.26	0.26	-7.65	-10.27
SPSR Nellore	1.05	1.64	0.66	-3.60	-3.94
Srikakulam	1.92	3.02	0.67	-8.97	-9.71
Visakhapatnam	0.57	1.43	0.22	-9.61	-10.57
Vizianagaram	2.31	4.68	0.23	-15.38	-16.90
West Godavari	0.36	0.21	0.57	-3.40	-13.90
Andhra Pradesh	100.00	-	-	-3.49	-6.00

Source: Authors' estimation using the data sourced from aps.dac.gov.in.

Table 2. Growth of production and yield of peanuts in various districts of Andhra Pradesh during the period between 1998 and 2020

District	Production Share (Average %)	Production share (Max %)	Production Share (Min %)	CGR (%) - Production (tonnes)	Max decline/increase -Production CGR (%)	CGR (%) - Yield (t/ha)
Anantapur	36.16	49.22	11.86	1.39	-76.54	3.63
Chittoor	18.98	26.47	13.42	-0.21	-13.40	3.74
East Godavari	0.13	0.36	0.05	-8.45	-8.81	2.65
Guntur	1.50	4.28	0.40	2.38	-6.08	4.17
Cuddapah	8.40	14.77	0.87	-1.77	-22.19	6.17
Krishna	1.39	3.10	0.54	-5.84	-8.98	4.25
Kurnool	18.24	26.84	8.11	0.37	-32.32	4.34
Prakasam	1.87	3.65	0.80	-2.89	-5.26	5.15
SPSR Nellore	4.10	8.29	1.03	-1.38	-28.44	2.31
Srikakulam	3.15	6.21	0.87	-7.28	-7.93	1.83
Visakhapatnam	1.01	2.93	0.31	-7.49	-11.72	2.34
Vizianagaram	3.47	8.97	0.29	-13.12	-15.13	2.65
West Godavari	1.25	2.47	0.37	-1.88	-10.56	1.57
Andhra Pradesh	100.00	-	-	-0.40	-12.38	1.56

Source: Authors' estimation using the data sourced from aps.dac.gov.in.

increased their per hectare groundnut yield by 30% owing to the use of cloud technology and business intelligence, in many of the districts the productivity continues to decline. The lower spread of suitable high yielding varieties/hybrids, moisture stress, high costs of production; untimely availability of inputs; low and fluctuating prices were found to be some of the key production constraints confronted at the farm level for the cultivation of groundnut [6]. Apart from few technological successes, the above issues confronting groundnut production should also be addressed for achieving higher production in the major groundnut producing districts.

(ii) Gaps and Potentials for Scaling up the Production

The estimated growth rates clearly indicate that in the recent past the area under groundnut cultivation is declining as a farmer shifting to remunerative crops like tobacco, cotton, maize including oil palm as market for these crops are stable and the prices are higher and risk-free. The studies also found that market prices were below the minimum support price for the most of the days in the state and the highest average difference in the state was 14%. Groundnut production in the state declined by 0.4% per annum due to decline in both productivity and area by 3% and 3.4% respectively. It is alarming to know that the maximum decline in production

was 12.38% in the State mainly due to the maximum decline in production in major groundnut producing districts namely Anantapur (76%) followed by Kurnool (32.32%), Cuddapah (22.19%) and Chittoor (13.40%). Production of the groundnut in the state declined by 0.4% per annum due to decline in both productivity and area by 3% and 3.4% respectively. It is alarming to note that the maximum decline in production was 12.38% due to maximum decline in production in major districts namely Anantapur (76%) followed by Kurnool (32.32%), Cuddapah (22.19%) and Chittoor (13.40%). Due to instability in crop area, the yield of groundnut is falling down as a result production either fell drastically or increase meagerly in most of the districts. Though the rate of decline in growth in terms of area was higher in Cuddapah district, which accounts for 8% of the total area in the state, compare to other major districts, the rate of growth in productivity of this district is very high which is more than 6%. The district constituted almost 15% of the state production, its production trended to decline mainly due to area. Since high productivity potential exists in this district, in order to increase the production of groundnut in the state, the area under groundnut in this district must be sustained without any further decline through technological and policy interventions (Tables 3 and 4).

Study showed that of the total farmers surveyed, 66 % adopted K-6 variety, 18 % adopted K-9 and

23 % adopted Dharani Variety [7,8]. The western mandals of Chittoor district in Andhra Pradesh are prone to frequent drought marked by deficit and late onset of rainfall and prolonged dry spells impacting productivity of groundnut. The study highlighted with demonstrations that yield, gross returns, net returns and BC Ratio for Dharani' groundnut variety were higher than Kadiri-6 [9]. The K-6 variety notified nationally in 2005, was renotified in 2015 till 2020, and its good taste and kernel look was commanding a good price, but overzealous farmers and [10]. Under various existing production constraints, increasing the adoption of suitable and high yield potential varieties signifies the timely distribution seeds.

Maximum productivity was recorded was 1.26 t/ha during 2007-08 in the Anantapur district and at present the productivity was 0.84 t/ha. The yield gap between the achieved productivity and the present productivity is 0.42 t/ha. The yield gap clearly reveals that if the maximum achieved productivity is the potential productivity of the district, then if this productivity could have been sustained over the period through technological and policy interventions, the production could have improved even at fall in area. Similarly, in case of Chittoor district the highest productivity

was 2.03 t/ha and which was achieved during 2017-18 thereafter the yield declined to 1.88 t/ha at present losing almost 1.5 t/ha as a result production declined despite overall increase in productivity growth. Cuddapah district achieved 2.42 t/ha at present which is comparatively higher than the other major producing districts. Hence, given production potential in this district, sustaining area through rainfed technological dissemination is critical for higher level of production in the state as this district contributes more than 8 % of the production in the State (Table 5). By interventions there could be a reduction in the cost of production paving the way for farmers to allocate more area under groundnut thereby sizeable fall in area in major districts can be reduced. The earlier study estimated that per hectare groundnut net income was found to be Rs.28, 971, whereas the competing crop maize was Rs.37, 710 per hectare. Per hectare cost of cultivation (C2) increased by 40% in the period II (2012-13 to 2016-17) over period I (2007-08 to 2011-12) and the operational costs increased by 52% compared to fixed costs which increased by 18%. [11,12]. Cost reduction by achieving the higher productivity through various interventions with assured market is paramount important for increasing the production in the state.

Table 3. Classification of districts based on the rate of growth of area

Rate of growth (%)	No of districts	Name of the district
-1 to - 2	1	Guntur
-2 to -4	5	Anantapur, Chittoor, Kurnool, SPSR Nellore and West Godavari
-6 to -8	2	Cuddapah and Prakasam
-8 to -10	3	Krishna, Srikakulam and Visakhapatnam
> -10	2	East Godavari and Vizianagaram
Total	13	-

Source: Authors' estimation using the data ranking classification from Table-1

Table 4. Classification of districts based on the rate of growth of production

Rate of growth	No of districts	Name of the district
< 1 to - 2	4	Chittoor, Cuddapah, SPSR Nellore, and West Godavari
-2 to -4	1	Prakasam
-4 to -6	1	Krishna
-6 to -8	2	Srikakulam and Visakhapatnam
- 8 to -10	1	East Godavari
>-10	1	Vizianagaram
< 1 to 2	2	Anantapur, Kurnool
2 to 4	1	Guntur
Total	13	-

Source: Authors' estimation using the data ranking classification from Table-2

Table 5. Classification of districts based on the rate of growth of yield

Rate of growth (%)	No of districts	Name of the district
1-2	2	Srikakulam and West Godavari
2-4	6	Anantapur, Chittoor, East Godavari, SPSR Nellore, Visakhapatnam and Vizianagaram
4-6	4	Guntur, Krishna, Kurnool and Prakasam
>6	1	Cuddapah
Total	13	

Source: Authors' estimation using the data ranking classification from Table-2

(iii) Intervention's Effects on Groundnut Production - Oilseeds Mission

India is one of the major edible oil consuming and producing countries in the world. National Mission on Oilseeds and Oil Palm (NMOOP) was implemented during the 12th Five Year Plan to expand the oil palm areas and increase the production of edible oils. Considering the growing domestic demand for edible oils, the staggering deficiency, and the cost to the exchequer on account of imports, the urgency of scaling up the oilseeds production does not need over-emphasis. It has now been planned to achieve a production of 45.64 Million tonnes (mts) from nine (9) annual oilseed crops by 2022-23, expecting an additional production of about 15.58 mt over and above the 30.06 Million tonnes production (Quinquennium Ending 2016-17). The projected demand for groundnut will be 20 million tonnes 2050. A growth rate of 4% to 5% in productivity is required to meet the demand both domestically and internationally. The strategies for enhancing the productivity of oilseed-based production system are prepared for the annual oilseeds and for oil palm in the country with low-cost technologies with high impact on productivity resulting in higher income. Technologies with high impact that involve reasonable investment with high return on investment (ROI), with emphasis on eco-friendliness, high input use efficiency, and strategies with emphasis on quality improvement and value addition leveraging technologies with a bearing on the employment through skill development.

Groundnut holds the first place in India presently growing in an area of 50.89 lakh hectares (2020-21) with annual production of 101.46 lakh tonnes (2020-21). The major groundnut producing states are Gujarat, Rajasthan, Tamil Nadu, Andhra Pradesh, Karnataka, Madhya Pradesh, West Bengal and Telangana contributing more than '00 thousand tonnes each. Among the various states Tamil Nadu stands first in terms of

productivity with an average of 2,980 kg per hectare followed by Gujarat (2001 kg/ha), Rajasthan (2064 kg/ha), Andhra Pradesh (957 kg/ha), Karnataka (802 kg/ha), Madhya Pradesh (1571 kg/ha), West Bengal (2363 (kg/ha), Telangana (2148 (kg/ha). MMI on Oilseeds under NMOOP was implemented in all 13 districts of Andhra Pradesh since the state average yield of oilseeds is less than the national average.

The State need to make effort for improving productivity of oilseeds particularly in case of groundnut which is less than national average. Under MM-I, Annual Action Plan (AAP) for 2016-17 has been approved with a total outlay of Rs. 2918.33 lakh (central share Rs 1751. lakh & state share Rs. 1167.33 lakh) which is being implemented by the State Department of Agriculture. The Components under MM-I on Oilseeds are classified broadly in three categories namely Seed Components, Production inputs, Transfer of Technology, Farm Machineries and irrigation tool, Mission Management expenses and Flexi funds. Increasing the certified seed availability by maintaining seed chain of recently released varieties, integrated crop management by increasing area under IPM / INM / Micro irrigation and demonstrating latest production / protection technologies at farmers' field through FLD / demonstrations, promotion of mechanization through distribution of improved farm implements / equipment and capacity building of farmers / extension workers through inter / intra trainings including FFS are major initiatives.

The interrupted time series model results for major groundnut growing districts are presented in the Tables 6- 9. In Andhra Pradesh, groundnut production and area are highest in Anantapur district. However, due to vagaries of rainfall, about 55-60 percent of groundnut cultivation acreage remained uncultivated in the district and groundnut seed sowing was confined to only 45 percent of the total area. Those farmers, who ventured to cultivate, are those with guaranteed

water sources and having the ability to withstand financial vagaries. Farmers and believed since there were no guarantees for marketing and remunerative price are fluctuating and volatile the farming operations have become expensive and investment burden has increased considerably. The model results of interrupted time series show that (Table 6) though there is no significant intervention effect, there is sign of productivity improvement after the intervention. Productivity increased from 0.45 t/ha to 0.84 t/ha after the intervention. The coefficient of sustained effect for production turned out to positive from negative though not significant (immediate effect), indicating the effect of mission. It is evident from the study [13] that, the growth

performance of groundnut in Ananthapuramu district declined over the years, particularly in Period III, which implies that, despite lot of efforts from researchers and government to encourage groundnut production at macro level, the contribution of groundnut at district level showed a declining trend. This scenario could be attributed to low farm level yields with higher farm level inefficiencies, gaps in production technology, geographical location of the district under rain shadow region, declining scenario in the length of growing period, shift towards competing crops, poor post-harvest support, changing climate, lower yields, low extension contact at field level, low market prices and high abiotic and abiotic stresses.

Table 6. Impact of Oilseeds Mission: Interrupted Time Series Model Results for Anantapur District

	Area (ha)	Production (tonnes)	Yield (t/ha)
β Coefficient			
Before Intervention (β_1)	-2282.05 (-0.44)	-22573.40* (-1.48)	-0.03* (-1.47)
Immediate Effect (β_2)	-163269.95*(-1.63)	-39656.50 (-0.13)	0.05 (0.12)
Sustained Effect (β_3)	-23976.06 (-1.02)	40132.66 (0.58)	0.09 (1.09)
Constant (β_0)	790634.25***(14.47)	639458.90***(3.97)	0.80***(3.97)

(Figures in parentheses are t ratios). * P < 0.10; *** P < 0.01
 Source: Authors' estimation using the data report sourced from aps.dac.gov.in.




Fig. 2. Interrupted Times Series – Model Results for Yield of Groundnut in Anantapur District

Table 7. Impact of Oilseeds Mission: Interrupted Time Series Model Results for Chittoor District

	Area (ha)	Production (tonnes)	Yield (t/ha)
	β Coefficient		
Before Intervention (β_1)	-5172.55***(-3.07)	-4747.98(-1.39)	0.004(0.19)
Immediate Effect (β_2)	7889.91(0.24)	-34138.29(-0.51)	-0.36(-1.03)
Sustained Effect (β_3)	-88.13(-0.01)	20718.99(1.34)	0.18**(2.21)
Constant (β_0)	223961.60***(12.59)	220545.76***(6.11)	0.97***(5.06)

(Figures in parentheses are t ratios). **P < 0.05; *** P < 0.01
 Source: Authors' estimation using the data report sourced from aps.dac.gov.in.

Table 8. Impact of oilseeds mission: interrupted time series model results for Cuddapah District

	Area (ha)	Production (tonnes)	Yield (t/ha)
	β Coefficient		
Before Intervention (β_1)	-5872.48**(-2.64)	-2774.70(-0.75)	0.02(0.91)
Immediate Effect (β_2)	-38074.04(-0.88)	-27197.70(-0.38)	-0.18(-0.37)
Sustained Effect (β_3)	1833.34(0.18)	1433.58(0.08)	0.15(1.30)
Constant (β_0)	190268.58*** (8.12)	130948.30***(3.35)	0.57** (2.17)

(Figures in parentheses are t ratios). * P < 0.05; *** P < 0.01
 Source: Authors' estimation using the data report sourced from aps.dac.gov.in.

Table 9. Impact of oilseeds mission: Interrupted time series model results for Kurnool District

	Area (ha)	Production (tonnes)	Yield (t/ha)
	β Coefficient		
Before Intervention (β_1)	-5518.27*** (-3.32)	-5796.45* (-1.69)	-0.01 (-0.42)
Immediate Effect (β_2)	-51118.60*(-1.59)	-50244.90(-0.75)	-0.02(-0.07)
Sustained Effect (β_3)	3790.70(0.50)	13162.88(0.84)	0.10(1.41)
Constant (β_0)	254203.32***(14.49)	241317.00***(6.66)	0.97***(5.78)

(Figures in parentheses are t ratios). * P < 0.10; *** P < 0.01
 Source: Authors' estimation using the data report sourced from aps.dac.gov.in.

The estimated coefficient of immediate effect for yield is -0.36 in the case of Chittoor district (Table 7) and turned to significant and positive (0.18) implying the sustained effect of the intervention. After intervention productivity of groundnut in this district increased from 0.85 t/ha to 2.03 t/ha with variability. Similarly, it is also found that there are no immediate and sustained effects of oilseed mission in Cuddapah district. Though there is no immediate effect of oilseed mission, the long-term effects of the co-efficient was positive and estimated at 0.15 indicating that as the time passes after the intervention the productivity of groundnut increases to 0.15 t/ha (Table 8). As a result, the productivity of groundnut in the Cuddapah district reached to 2.42 t/ha, which was achieved in the year 2019-20. Predominant crop is rainfed groundnut, which accounts for more than 50% of the annual cropped area in the district. The studies conducted in Cuddapah districts identified major technical constraints relating to adoption of

improved technology, non-suitability of improved technology, and lack of technical guidance [14]. Not much impact of intervention both in during short and longer periods was observed in the Kurnool district (Table 9). Addressing these issues through policy and technological interventions are crucial for sustaining the productivity and increasing the production of groundnut [15].

4. CONCLUSION AND IMPLICATIONS

Area under the groundnut in all the districts of Andhra Pradesh are declining at faster rate and the rate of decline is higher in the major groundnut growing districts namely Anantapur (2.13%), Chittoor (3.79%), Cuddapah (7.47%) and Kurnool (3.82 %). The area under groundnut in the State also registered a negative rate of growth and the area declined at the rate of 3.49 % between 1997-98 and 2019-20 and maximum decline was 6 per cent, which is alarming.

Anantapur district contributes on an average 57 % of the state area and maximum share was 66 %. Since, the maximum decline is in this district was 5.12 %, the production growth was found low (1.39%) despite the productivity increased by 3.63 % per annum. Similarly, the rate of production in Chittoor district declined at 2.1% overall, though the productivity increased at the rate of 4 % annum and this is mainly due to a decline in area by almost 4 % per annum. The Cuddapah, one of the major groundnuts producing districts in the state, also witnessed a negative growth rate in production and the production decline was around 2% per annum. However, productivity in the Cuddapah district grew at faster rate compare to other major districts and the productivity has grown at 6% per annum. Groundnut production in the state declined by 0.4% per annum due to decline in both productivity and area by 3% and 3.4% respectively. It is alarming to know that the maximum decline in production was 12.38% in the State mainly due to the maximum decline in production in major groundnut producing districts namely Anantapur (76%) followed by Kurnool (32.32%), Cuddapah (22.19%) and Chittoor (13.40%).

Hence, given production potentials in the major groundnut producing districts, sustaining area through rainfed technological dissemination is critical for higher level of production in the state as these districts contributes more than 80 % of the production in the State. By interventions there could be a reduction in the cost of production paving the way for farmers to allocate more area under groundnut thereby sizeable fall in area in major districts can be reduced. The yield gap clearly reveals that if the maximum achieved productivity is the potential productivity of the districts, then this productivity could have been sustained over the period through technological and policy interventions, the production could have improved even at fall in area.

Those farmers, who ventured to cultivate, are those with guaranteed water sources and having the ability to withstand financial vagaries. Farmers believed that since there are no guarantees for marketing and remunerative prices are fluctuating and volatile, the farming operations have become expensive and investment burden has increased considerably. The model results of interrupted time series show that though there is no significant immediate intervention effect, there is sign of productivity

improvement in the long run after the intervention. The coefficient of sustained effect for production turned to positive from negative though not significant (immediate effect), signifying the effect of oilseeds mission in the major districts. The studies identified major technical constraints relating to adoption of improved technology, non-suitability of improved technology, and lack of technical guidance. Addressing these issues through policy and technological interventions are crucial for sustaining the productivity and increasing the production.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Acharya NG. Ranga Agricultural University. Crop Outlook Reports of Andhra Pradesh; 2021.
2. Reddy G. Raghunadha, M. Chandrasekhar Reddy, K. Gurava reddy, S. K. Sneha and B. Meher Gita. An economic analysis of oilseed sector in Andhra Pradesh. J. Res. Angrau. 2022;50(2):118-133.
3. Nayak A, Lokesha H, Gracy CP. Growth and Instability Analysis of Groundnut Production in India and Karnataka. Economic Affairs. 2021;66(1):61-69.
4. Krishna N Priya, Padmodaya B, Srinivasulu DV, Shilpakala V. Production Constraints in Groundnut Crop in Kadapa District of Andhra Pradesh. J Krishi Vigyan. 2021;10 (1):218-222.
5. Available: www.apseeds.ap.gov.in
6. Teja I Krishna, Ramana Rao SV, Praveen Kumar P, Lora Anusha P. Production Constraints faced by the farmers in groundnut, sesame and sunflower cultivation in Andhra Pradesh. The Pharma Innovation Journal. 2022;11(11):2239-2242.
7. Chowdary K. Raghavendra, Vemaraju, M. Tejaswini and M. Reddi Kumar. A study on the extent of adoption of ANGRAU Technologies in groundnut crop by farmers of Rayalaseema zone of Andhra Pradesh. Asian Journal of Agricultural Extension, Economics & Sociology. 2021;39(5): 29-33.
8. Reddy A. Malliswara, B. Sahadeva Reddy, R. Veeraraghavaiah, G. Ravindrachary and

- K.A. Gopinath. Evaluation of climate resilient varieties of groundnut for scarce rainfall zone of Andhra Pradesh. Extended Summaries: 5th International Agronomy Congress; 2021, India.
9. Sreenivasulu S, Divya V, Sudhakar PS, Ramu Kumar T, Prasad JV, Prasad YG, Prasad JVNS. Performance of Drought Tolerant Groundnut Variety in Chittoor District of Andhra Pradesh. J Krishi Vigyan. 2021;10 (1):196-199.
 10. The Hindu, May, 3; 2022.
 11. Sunandini GP, Irugu Shakuntala Devi. Economic analysis of groundnut production in Andhra Pradesh. International Journal of Economic Plants. 2020; 7(4):176-179.
 12. Naidu Challa Bhargavi, Sanjay Kumar and A.K .Rai (2019). An economic analysis of production of groundnut (Arachis Hypogaea) in Anantapur district of Andhra Pradesh. *International Journal of Innovative Science and Research Technology*.4 (5). 482-487
 13. Teja I Krishna Teja, Ramana Rao SV, Bhavani Devi I, Prasad SV, Ravindra Reddy B, Paladugu Praveen Kumar. Growth performance of groundnut in Ananthapuramu district of Andhra Pradesh – A Temporal Analysis. Asian Journal of Agricultural Extension, Economics & Sociology. 2021;39(7): 81-87.
 14. Veeraiah A, Shilpakala V, Ramalakshmi Devi S, Ankaiah Kumar K. Constraint Analysis of Groundnut Cultivation in YSR District of Andhra Pradesh, India. International Journal of Current Microbiology and Applied Sciences. 2019;8(7):1488 -1493.
 15. Available:www.apagrisnet.gov.in/2017/Agri%20Action%20Plan%20201718%20(English)/19%20NMOOP.pdf

© 2023 Burigi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/99705>