



Impact of Mulch and Irrigation Schedules on Yield and Quality of Spring Sunflower in Tarai Region of Northern India

Rakesh Dawar ^{a*}, Mahendra Singh Pal ^b,
Akshay Kumar Yogi ^a, Deepak Kumar Meena ^a
and Sunil Kumar ^a

^a Division of Agronomy, ICAR - Indian Agricultural Research Institute, New Delhi - 110 012, India.

^b Department of Agronomy, College of Agriculture, G. B. Pant University of Agriculture and Technology, Pantnagar- 263145 (U.S.Nagar, Uttarakhand), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i92218

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100987>

Original Research Article

Received: 10/04/2023

Accepted: 12/06/2023

Published: 28/06/2023

ABSTRACT

The field experiment was carried out at Forage Block, Instructional Dairy Farm, Nagla, G. B. Pant University of Agriculture & Technology, Pantnagar (Uttarakhand) in spring 2019 to, study the effect of mulches and irrigation schedules on productivity and water use efficiency of spring sunflower (*Helianthus annuus* L.) in Mollisols of Uttarakhand (India). The experiment consisted of 03 levels of mulches i.e., control (no mulch), straw mulch @ 6t/ha and polythene mulch (black) in the main plot and 04 levels of irrigation schedules i.e., critical stages (button, flowering, and seed filling stage), 0.8, 1.0 and 1.2 IW/CPE ratios in subplots, was laid out in split-plot design with three replications. Under the application of polythene mulch, significant seed output was observed that was

*Corresponding author: E-mail: dawarrakesh601@gmail.com;

statistically comparable to straw mulch. Under polythene mulch, seed yield was found to be 4.6% and 18.5% greater than under straw mulch and control, respectively. The polythene mulch produced significantly higher stover yield that was non-significant with straw mulch Polythene mulch produced 8.10% and 10.4 % higher stover than straw mulch and control, respectively. Sunflower seed yield differed significantly among irrigation schedules. Irrigation applied at 1.2 IW/CPE produced significantly the highest seed yield that was 5.4, 11.1, and 14.8% greater than irrigation scheduled at 1.0, 0.8 IW/CPE, and critical stages, respectively. Among the irrigation schedules, stover yield was recorded 5.8, 8.0, and 10.8% higher under the 1.2 IW/CPE ratio than 1.0, 0.8 IW/CPE ratio, and critical stages, respectively. Overall, it is highlighted that polythene mulch gives significantly higher productivity and quality parameters. 1.2 IW/CPE ratios give significantly the highest seed yield, higher protein content and oil content.

Keywords: Heading; irrigation schedules; mulch; protein; Tarai.

1. INTRODUCTION

Sunflower oilseed production has been steadily increasing worldwide, driven by the rising demand for vegetable oils and the utilization of its oil in various sectors. In terms of sowing areas worldwide, oil crops are second in importance only to cereals. "Sunflower (*Helianthus annuus* L.) is a day-neutral, short duration, drought, and salinity tolerant oilseed crop belonging to the aster family and is native to southern parts of the USA and Mexico. Sunflower ranks fourth next to soybean, groundnut, and rapeseed within the total production of oilseeds of the planet. It is cultivated in a vicinity of 27.9million hectares with annual production and productivity of 38.9 million tonnes and 1700 kg per hectare, respectively within the world within the year" (USDA, 2022) "Now, the crop has been well accepted by the farming community due to its desirable attributes like short duration, photoperiod insensitivity, adaptability to a good range of soil and atmospheric condition, drought tolerance, lower seed rate, higher seed multiplication ratio and good quality edible oil. The cultivation of sunflower is basically confined to southern parts of the country comprising the states of Karnataka, Maharashtra, Tamil Nadu, and state. These four states contribute about 90 percent of total acreage and 78 percent of total production. Out of these four states Karnataka is the leading sunflower-producing state contributing nearly 52 percent of the whole area and 40 percent of the entire production within the country. However, recently sunflower has moved to northern parts of the country where the productivity is extremely high. In India, sunflower is grown over a region of 0.22 M ha with production and productivity of 0.23 Million tonne and 1023 kg per hectare, respectively during the year 2020-21" (Directorate of Economics & Statistics, 2021). "It is considered to be an exhaustive crop. While the realm under the sunflower crop is increasing, the

productivity is not on par with the cultivable area. Among the several reasons because of low productivity, inadequate and imbalanced nutrition of essential nutrients is taken into account as a serious one. The availability of water is decreasing day-by-day and therefore the share of water to agriculture is decreasing very sharply over the previous couple of decades" (Kadasiddappa et al., 2015).

"Among the assorted approaches used for scheduling irrigation, a modified meteorological approach supported the ratio of fixed amount of irrigation water (IW) and cumulative pan evaporation (CPE) has been found suitable for the guidance of irrigation scheduling to different crops" (Parihar et al., 1976). "Among different conservation practices, mulching is an important technique which is known for its beneficial impacts on the soil-plant systems. Several advantages of mulching on soil and crops are prevention of soil erosion and runoff loss, reduction in the evaporation loss of water, conservation of residual soil moisture, maintenance of soil temperature and improvement in the soil physical, biological and chemical properties over time" [1]. According to Domber et al. (2009), sunflower hybrids reacted admirably to irrigation and mulch control. Tunio et al. (2007) reported "significant differences in seed germination; seed per head, seed index and seed yield due to different types mulch and irrigation frequencies. Therefore, an experiment was carried out to investigate the impact of different mulches and irrigation scheduling on growth and yield characteristics in northern region".

2. MATERIALS AND METHODS

The field experiment was conducted in spring 2019 at the Forage Block, Instructional Dairy Farm, Nagla, G. B. Pant University of Agriculture

& Technology, Pantnagar, Uttarakhand, India. The purpose was to investigate the impact of mulches and irrigation schedules on the productivity and water use efficiency of spring sunflower (*Helianthus annuus* L.) in the Mollisols soil of Uttarakhand. The experimental site had silty clay loam soil with specific characteristics such as pH 7.13, electrical conductivity (EC) 0.20 ds/m, organic carbon content 0.68%, bulk density 1.57 g/cc, and available nutrients including nitrogen, phosphorus, and potassium at levels of 280.2, 25.2, and 215.2 kg/ha, respectively. The experiment included three levels of mulches: control (no mulch), straw mulch at a rate of 6 tons/ha, black polythene mulch and Four levels of irrigation schedules viz., critical stages (button, flowering and seed filling), 0.8, 1.0, and 1.2 IW/CPE ratios. The experiment was laid out in split plot design with three replications. The previous crop grown at the site was dhaincha (*Sesbania rostrata*). The sunflower hybrid variety 'DRSH-1' was sown manually and were treated with Carbendazim @3g/kg seed. The plants were spaced in lines 60 cm apart with a seed-to-seed distance of 30 cm. The plant-to-plant distance was adjusted after 10 days of sowing. The recommended chemical fertilizer dose of 120:60:40 kg N:P:K₂O /ha was applied, with half of the nitrogen and the full dose of phosphorus and potassium applied at sowing. The remaining nitrogen was top-dressed after 25 days of sowing at the first irrigation. Mulching was performed with oat straw at 6 tons/ha or black polythene sheet after the first common irrigation at 25 days of sowing, depending on the treatment. Based on the IW/CPE ratio and the crop's crucial stages, irrigation was planned. The cumulative pan evaporation (CPE) results were collected from a USWB Class-1 open pan evaporimeter situated at the university's crop weather observatory. The number of irrigations applied at different IW/CPE ratios 0.8, 1.0, and 1.2 was 6,4, and 3, respectively. The crop was harvested when the lower portion of the heads turned yellowish-brown, bracts dries and the seeds became dark brown, and the thalamus turned yellow. Observations on growth and yield attributes were recorded from five randomly selected plants in each plot. The seed yield was calculated based on the net plot area for each treatment and presented in kg/ha. Protein content was determined using the Kjeldahl method.

CGR and RGR values were calculated for 30-60 DAS and 60 DAS-harvest by the formula (Watson et al., 1952).

$$\text{Crop growth rate (g/m}^2\text{/day)} = W_2 - W_1/t_2 - t_1$$

$$\text{Relative growth rate (mg/g/day)} = \frac{\text{Loge}W_2 - \text{Loge}W_1}{t_2 - t_1}$$

Where, W1 and W2 are the dry weights (mg) of plants at time t1 and t2, respectively. The statistical analysis of the data for various growth, yield, and quality parameters was conducted using a split-plot design and the standard approach of Analysis of Variance (ANOVA) as described by Gomez and Gomez (1984).

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

3.1.1 Effect of mulch

“The application of mulch had a significant effect on 50 % flowering with maximum and minimum days taken under polythene mulch and control, respectively, however, polythene mulch and straw mulch were found non-significant to each other. This might be due to an inadequate moisture regime under control. A significantly higher crop growth rate was recorded at 30-60 DAS under polythene mulch than straw mulch and control but it was significantly equal to straw mulch. Significantly lowest crop growth rate was recorded under control. 60 DAS to harvest, significantly highest crop growth rate was observed under polythene mulch followed by straw mulch and significantly lowest crop growth rate was recorded under control. 30-60 DAS, significantly higher relative growth rate was recorded under polythene mulch followed by straw mulch and control but both were non-significant with each other. Significantly lowest relative growth rate was observed under control. At 60 DAS-harvest, the relative growth rate was not affected significantly by mulches however the highest RGR was recorded under polythene mulch followed by straw mulch and under the lowest in the control treatment. This might be attributed mulching reducing water loss by evapotranspiration avoid salt accumulation, conserving moisture in soil, promoting crop growth and increasing crop water use efficiency” [1].

3.1.2 Effect of irrigation schedules

Effect of irrigation schedules the days taken to 50% flowering were affected greatly by irrigation schedules. The irrigation schedule at 1.2 IW/CPE took maximum while other irrigation schedules i.e. critical stages, 0.8 and 1.0 IW/CPE ratio took minimum days to 50 % flowering; however, all irrigation schedules had a non-significant effect

on 50 % flowering. Early flowering was recorded due to a higher frequency of irrigation leading to the early formation of reproductive parts. The crop growth rate was influenced significantly by irrigation schedules. At 30-60 DAS, a significantly higher crop growth rate was recorded when irrigation was scheduled at 1.2 IW/CPE than 0.8, 1.0 IW/CPE, and critical stages, but it was statistically at par with 1.0 IW/CPE. Significantly the lowest crop growth rate was observed when irrigation was applied at critical stages. A similar trend was observed at 60 DAS to harvest where the highest crop growth rate value was recorded at irrigation scheduled at 1.2 IW/CPE and the lowest crop growth rate was observed when irrigation was applied at critical stages. The increasing irrigation frequency increased crop growth rate due to the availability of adequate soil moisture that supported higher cell divisions and expansion of cells [2]. Irrigation schedules had a significant effect on RGR at both 30-60 DAS and 60 DAS harvest. Significantly the highest relative growth rate was found at the 1.2 IW/CPE ratio which remained non-significant with the 1.0 IW/CPE ratio. The lowest RGR was recorded when irrigation was applied at critical stages, though it was non-significant with 0.8 and 1.0 IW/CPE ratios. Similarly, at 60 DAS – harvest, significantly highest RGR was measured at 1.2 IW/CPE irrigation schedules that were statistically at par with 0.8 and 1.0 IW/CPE ratio. The higher value was attributed to more number of leaves as well as LAI.

3.2 Yield Attributes and Seed Yield

3.2.1 Effect of mulch

The mulch application had a significant effect on sunflower seed yield. Under the application of polythene mulch, which was statistically comparable to straw mulch, a significantly higher seed yield was noted, and the lowest yield was discovered under control conditions. In comparison to straw mulch and the control, the sunflower seed output was 4.6% and 18.5% higher under polythene mulch, respectively. A 13.3% higher seed output than the control was achieved with the addition of straw mulch. The higher sunflower yield was due to greater head diameter, seed weight, 100 seed weight, and seed weight per head. Similar findings were also recorded by Arora et al., [3].

The polythene mulch produced significantly the highest stover yield that was non-significant with straw mulch. Significantly the lowest stover yield was observed under control. Polythene mulch

produced 8.10% and 10.4 % higher stover than straw mulch and control, respectively. The higher stover yield was due to more plant height, the number of leaves, and dry matter accumulation. Din et al., (2013) and Rajput et al., (2014) also supported the above results. The mulch had a significant effect on the harvest index of sunflower and a significantly higher value of harvest index was recorded under polythene mulch that was statistically at par with straw mulch. Significantly the lowest harvest index was recorded under control. A higher harvest index recorded under polythene mulch might be due to a higher fraction of seed yield compared to stover yield [4,5].

3.2.2 Effect of irrigation schedules

Sunflower seed yield differed significantly among irrigation schedules. Irrigation applied at 1.2 IW/CPE produced significantly the highest seed yield that was 5.4, 11.1, and 14.8% greater than irrigation scheduled at 1.0, 0.8 IW/CPE, and critical stages, respectively. The seed yield didn't vary significantly between irrigation applied at critical stages and 0.8 IW/CPE ratio, however, irrigation scheduled at 1.0 IW/CPE ratio also gave a significantly higher seed yield than 0.8 IW/CPE ratio and critical stages. Such differences in the seed yield were attributed to favorable water regimes in soil for better mobilization of nutrients and also enhanced source capacity and sink strength which in turn influenced yield attributing characters. Gurumurthy et al. (2008) also recorded a higher yield of a sunflower when irrigation was applied at 1.2 IW/CPE mainly due to higher values of yield contributing attributes. The interaction effect between mulch and irrigation schedule was found non-significant. Among the irrigation schedules, significantly the highest stover yield was found when irrigation was scheduled at 1.2 IW/CPE followed by irrigation applied at 0.8 and 1.0 IW/CPE ratio. Significantly the lowest stover yield was produced at irrigation applied at critical stages. The stover yield was recorded as 5.8, 8.0, and 10.8% higher under the 1.2 IW/CPE ratio than 1.0, 0.8 IW/CPE ratio, and critical stages, respectively. The higher stover yield was attributed to higher values of growth and yield attributes. These results were also supported by Mahal et al. [6] the irrigation schedules did not have any significant effect on the harvest index however higher value of harvest index was recorded when irrigation was applied at 1.2 IW/CPE ratio followed by 1.0 and 0.8 IW/CPE ratio. The lowest harvest index was recorded at irrigation applied at critical stages.

Table1. Effect of mulch and irrigation schedules on CGR, RGR, and 50% flowering of sunflower

Treatments	Crop growth rate (g/m ² /day)		Relative growth rate (g/g/day)		50 % flowering (Days)
	30-60 DAS	60 DAS- at harvest	30-60 DAS	60 DAS- at harvest	
A) Mulch					
Control	10.11	5.32	0.077	0.015	63
Straw mulch	10.45	5.75	0.079	0.018	64
Polythene mulch	10.50	5.90	0.083	0.020	65
SEm ±	0.03	0.02	0.010	0.023	0.16
CD (P=0.05)	0.10	0.08	0.030	NS	01
B) Irrigation schedules					
Critical stages	9.39	5.10	0.075	0.016	64
0.8	10.00	5.43	0.076	0.018	64
1.0	10.88	5.65	0.077	0.018	65
1.2	10.95	5.80	0.079	0.019	64
SEm±	0.05	0.05	0.001	0.001	0.24
CD (P=0.05)	0.20	0.18	0.002	0.003	NS

Table 2. Effect of mulch and irrigation schedules on yield attributes and quality parameter

Treatments	Stover yield (kg/ha)	Seed yield (kg/ha)	Harvest index	Oil Percent %	Oil yield (kg/ha)	Protein (%)	Protein yield (kg/ha)
Control	7470.10	2594.54	25.50	37.50	885.00	17.79	461.71
Straw mulch	7483.64	2938.34	26.94	37.91	1015.05	17.87	525.43
Polythene mulch	8356.82	3074.30	27.56	38.81	1083.78	18.16	558.78
SEm ±	82.03	75.50	0.34	0.16	24.16	0.03	13.05
CD (P=0.05)	330.72	235.62	1.34	0.65	97.5	0.12	52.63
Critical stages	7367.00	2687.50	26.58	36.95	903.02	17.71	476.30
0.8	7674.52	2776.00	26.74	37.35	943.42	17.92	497.80
1.0	7698.52	2927.76	26.54	37.85	1004.15	18.02	527.88
1.2	8286.25	3085.00	26.81	40.16	1127.80	18.11	559.16
SEm±	59.42	39.54	0.37	0.22	16.42	0.02	7.25
CD (P=0.05)	177.94	118.50	NS	0.66	49.17	0.07	21.71

3.3 Quality Parameter

3.3.1 Effect of mulch

The significantly highest oil content was recorded under polythene mulch but values of oil content were statistically at par grown under straw mulch and control. Higher oil content values were caused due to higher 100 seed weight favored by a better plant grown and development. Sheikh et al. (2002) also reported higher oil content under mulch application. Significantly higher oil yield was recorded under polythene mulch that was statistically similar to straw mulch. The oil yield was recorded 6.0 to 18.4% higher than straw mulch and control, respectively. Higher oil yield was the result of high oil content and seed yield. The above result was supported by Yenpreddiwar et al. [7]. The polythene mulch

recorded significantly highest protein content followed by straw mulch and control. However, there was no significant difference between control and straw mulch. The higher protein content was attributed to higher nitrogen content in the seed. Similar findings were reported by Jain et al. [8]. Significantly highest protein yield was found under polythene mulch that remained non-significant with straw mulch and the lowest protein yield was found under control. The higher protein yield was the result of higher protein content and seed yield. Hingonia et al. [9] also reported a similar agreement.

3.3.2 Effect of irrigation schedules

The oil content increased with increasing IW/CPE ratio and the highest value was recorded when irrigation was applied at 1.2 IW/CPE ratio

but the oil content was found significantly similar at both 0.8 and 1.0 IW/CPE ratios. Significantly lowest oil content was extracted at irrigation scheduled at critical stages. The higher values of oil content might be due to better soil moisture caused higher availability and uptake of nutrients which in turn produced bold seed and greater seed test weight and it was supported by Hittinahalli et al. [10]. The interaction effect was found non-significant. The oil yield was increased with increasing irrigation frequencies with the significantly highest value at 1.2 IW/CPE ratio followed by 1.0 IW/CPE ratio. Significantly lowest oil yield was found when irrigation was applied at critical stages that remained statistically at par with a 0.8 IW/CPE ratio. The oil yield was recorded 10.9, 16.3, and 19.9% higher under the 1.2 IW/CPE ratio than 1.0 and 0.8 IW/CPE ratio as well as critical stages, respectively. This was ascribed due to better formation and development of seeds associated with higher oil content and seed yield. Similar results were also reported by Hittinahalli et al., [10]. The interaction effect was found non-significant. The irrigation schedules had a significant effect on protein yield and it was increased with increasing irrigation frequencies. The protein content increased with increasing irrigation frequencies and a significantly higher value was found at the 1.2 IW/CPE ratio followed by a 1.0 IW/CPE ratio that had a significantly higher value than irrigation schedules at critical stages and 0.8 IW/CPE ratio. The higher value was attributed to high N content under higher irrigation frequency favored by more N uptake. The interaction effect between mulch and irrigation schedules was recorded as non-significant. Hussain et al. (2001) also reported the same findings. Significantly highest protein yield was recorded under 1.2 IW/CPE ratio and irrigation scheduled at critical stages gave significantly lowest value that was non-significant with irrigation scheduled at 0.8 IW/CPE ratio. Higher protein yield was the result of higher protein content and seed yield. The same results were recorded by Hussain et al. (2001).

4. CONCLUSION

The results indicated that growth, stover yield and seed yields of sunflower were highest under polythene mulching and also irrigation scheduled at 1.2 IW/CPE ratio. On other hand, the scheduling irrigation at 1.2 IW/CPE ratios gave significantly higher in quality parameter. It is, therefore, concluded that sunflower may be grown with application of straw mulch @ 6 ton/ha

with irrigation Scheduled at 1.2 IW/CPE ratios for higher productivity, as well as protein and oil content during spring season in Uttarakhand and may also be replicated in similar agro-ecological zones of India.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar T, Haque MA, Md Saiful I. Effect of polythene mulch on growth and yield of sunflower (*Helianthus annuus*). Archives of Crop Science. 2018;2(1):38-46.
2. Ramamoorthy K, Arthanari PM, Subbian P. Effect of irrigation scheduling and method of irrigation on productivity and water economy in hybrid sunflower (*Helianthus annuus* L.). Helia. 2009;32(50):115-122.
3. Li J, Qu Z, Chen J, Yang B, Huang Y. Effect of planting density on the growth and yield of sunflower under mulched drip irrigation. Water. 2019;11():752.
4. Kadasiddappa MM, Rao VP, Reddy KY, Ramulu V, Devi MU, Reddy SN. Effect of irrigation (drip/surface) on sunflower growth, seed and oil yield, nutrient uptake and water use efficiency-A review. Agricultural reviews. 2017;38(2):152-8.
5. Arora VK, Singh AS, Thind SS. Irrigation, tillage and mulching effects on soybean yield and water productivity in relation to soil texture. Agriculture Water Management. 2011;98(1):563-68.
6. Mahal SS, Brar JS, Brar AS, Vashist KK. Water productivity of spring sunflower (*Helianthus annuus* L.) under varying planting methods and irrigation scheduling. Crop Research. 2012;44(1&2):82-84.
7. Yenpreddiwar MD, Nikam RR, Dange RB, Gaidhane SN. Water use studies in linseed as influenced by different irrigation levels and moisture conservation practices. Journal of Soils and Crops. 2007;17(1): 176-177.
8. Jain NK, Meena HN, Bhaduri D. Improvement in productivity, water use efficiency, and soil nutrient dynamics of summer peanut (*Arachis hypogaea* L.) through use of polythene mulch, hydrogel, and nutrient management. Communications in Soil Science and Plant Analysis 2017;48(5):549-564.

9. Hingonia K, Singh RK, Meena NR, Verma HP, Meena RP. Effect of mulch and irrigation levels on yield and quality of barley (*Hordeum vulgare* L.). Journal of Pure and Applied Microbiology. 2016; 10(4):2925-2930.
10. Hittinahalli NM, Malligawad LH, Giriraj K. Productivity of hybrid sunflower as influenced by irrigation, nitrogen and sulphur fertilization during the summer season. Proceedings Sixteenth International Sunflower Conference, Fargo, ND USA. 2007:363-369.

© 2023 Dawar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/100987>