



# Impact of Tillage, Irrigation Regimes and Nitrogen Levels on Soil Moisture Dynamics, Growth and Productivity of Canola (*Brassica napus*)

Gagandeep Kaur <sup>a</sup>, Satinder Singh Brar <sup>a\*</sup>, C. B. Singh <sup>a</sup>,  
Madhu Dhingra <sup>a</sup> and K. B. Singh <sup>a</sup>

<sup>a</sup> Department of Soil Science, Punjab Agricultural University, Ludhiana-141004, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i173214

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102670>

Original Research Article

Received: 01/05/2023  
Accepted: 01/07/2023  
Published: 07/07/2023

## ABSTRACT

A field experiment was conducted at Punjab Agricultural University, Ludhiana, during *rabi* 2018-19 to examine the effects of tillage, irrigation and nitrogen rates on the productivity of canola (*Brassica napus*). Combinations of two tillage systems (Deep tillage- DT, and conventional tillage- CT), with three irrigation regimes viz; no irrigation ( $I_0$ ), one irrigation ( $I_1$ ) and two irrigations ( $I_2$ ) in main plots and four nitrogen (N) rates viz; 0 ( $N_0$ ), 50 ( $N_{50}$ ), 75 ( $N_{75}$ ) and 100 ( $N_{100}$ ) kg ha<sup>-1</sup> in sub-plots with three replications. Lower soil moisture content was recorded under deep tillage. Root density in upper 60 cm soil depth was higher in  $I_2$  followed by  $I_1$  and  $I_0$  irrigation regimes whereas below 60 cm, it was higher under  $I_0$ . Higher root density was recorded under DT and  $N_{100}$  plots. Irrigation and N application significantly improved plant height, relative leaf water content and SPAD value at different stages. Yield attributes and yield were highest under  $I_2$ , deep tillage and  $N_{100}$  treatment. Oil

\*Corresponding author: E-mail: brarpau2010@gmail.com;

content also improved with successive increments of N rate. It may be concluded that, for higher productivity, canola can be grown under deep tillage with sufficient irrigation (I<sub>2</sub>) and N fertilization (N<sub>100</sub>).

**Keywords:** Canola; soil moisture dynamics; N rates; root length density; yield attributes.

## 1. INTRODUCTION

Rice-wheat (R-W) cropping system in the north-western region of India, although ensuring national food security, has indeed resulted in soil degradation and over-use of underground water resources. In addition, conventional practices for crop management in the R-W system involve a high cost of production and inefficient input utilization [1]. Diversifying R-W systems with oilseed-based systems and alternative crop-saved irrigation water and soil management practices may improve the productivity of the system, environmental quality and sustaining soil health. Rapeseed and mustard (*Brassica* species; Family *Brassicaceae*), being the second largest edible oilseed crop after groundnut, accounts for 31.4% of the Indian edible oilseed pool [2,3]. India has a 59.77 lakh ha area under rapeseed mustard with a production of 8430 thousand tonnes [4]. The share of oilseeds is 14.1 per cent out of the total cropped area in India; rapeseed-mustard accounts for 3 per cent of it [5]. However, high levels of erucic acid and glucosinolates in Indian cultivars have reduced their preference in the international market. Canola is an internationally accepted nomenclature for *Brassica* varieties with less than 2% erucic acid in the oil and glucosinolates less than 30 micromoles per gram defatted meal [6]. Canola oil is used for human consumption and seed meal as a rich source of protein for livestock including poultry [7]. It is a magnificent feedstock for biofuel production. Canola-quality oilseed rape (*Brassica napus* L.) varieties have been fairly successful in India.

The yield of oilseed rape is greatly influenced by various agronomic practices such as tillage, irrigation and fertilization [8,9,10]. Tillage has been an important aspect of technological advancement in the evolution of agriculture, especially in food production. Among the main factors of crop production, this is an important factor contributing to approximately 20% of the increase in crop yield [11]. Mechanical modifications of soil profiles, commonly referred to as deep tillage, could reduce high subsoil

strength, promoting deeper rooting and thus, the plant availability of subsoil resources [12]. Irrigation promotes the growth and yield attributes of mustard by supplementing the water requirement of the crop. Better results both in terms of biometric components and seed yield can be achieved by the application of optimum irrigation. Non-availability of sufficient irrigation water as per the requirements of mustard crops causes moisture stress at critical stages of growth and development. Irrigation provided at the most critical growth stages (at the flower initiation stage and silique development stage) produced the maximum growth and yield attributes [13]. Rapeseed-mustard group of crops have a relatively higher demand for nitrogen than many other crops [14] and the nitrogen fertilizer rate influences canola seed yield and quality [15]. Excess nitrogen results in greater succulency in plants and thus may lead to a higher incidence of insect pests and diseases. Hence, an optimum dose of N is of utmost importance to maintain high N use and water use efficiency.

Therefore, the current study was conducted to evaluate the effect of tillage, irrigation and N rates on the growth, yield and oil content of canola along with soil moisture distribution during the growing season.

## 2. MATERIALS AND METHOD

### 2.1 Site Description

A multi-factor study was conducted during the *rabi* season of 2018-19 at the Punjab Agricultural University, Ludhiana situated at 30° 54' N latitude and 75° 48' E longitude at a height of 247 m above mean sea level. Important physical and chemical properties of the soil are outlined in Table 1. Total rainfall during the growing season was 171.6 mm. Pan-evaporation (545 mm) was below the long-term average (685.9 mm). During the growing season, the mean maximum air temperature ranged from 18.5-35.1°C to the normal 18.2-34.4°C; whereas the mean minimum temperature ranged from 5.5-19.5°C compared to the normal 5.6-17.1°C, respectively.

**Table 1. Physico-chemical properties of experimental site**

Depth (cm)	Soil separates (%)			Textural class	Bulk Density (Mg m <sup>-3</sup> )	Water holding capacity (% v/v)	pH (1:2 soil:water suspension)	EC (dS m <sup>-1</sup> )	Organic carbon (%)	Available N (kg ha <sup>-1</sup> )	Available P (kg ha <sup>-1</sup> )	Available K (kg ha <sup>-1</sup> )
	Sand	Silt	Clay									
<b>0-15</b>	79.3	11.7	7.6	Loamy Sand	1.38	42.7	8.25	0.19	0.30	80.3	31.8	164.9
<b>15-30</b>	80.6	12.3	6.3	Loamy Sand	1.42	43.5	8.50	0.18	0.22	77.1	25.4	133.3
<b>30-60</b>	81.9	9.5	8.5	Loamy Sand	1.47	40.4	8.46	0.14	0.15	60.3	19.3	111.4
<b>60-90</b>	83.5	7.7	8.8	Loamy Sand	1.53	41.7	8.44	0.13	0.09	52.8	17.9	103.3
<b>90-120</b>	84.3	8.4	7.3	Loamy Sand	1.55	41.3	8.54	0.12	0.05	55.5	15.7	102.8
<b>120-150</b>	82.8	8.7	8.5	Loamy Sand	1.57	41.4	8.56	0.11	0.01	42.1	7.7	101.3
<b>150-180</b>	81.5	9.8	8.7	Loamy Sand	1.59	41.2	8.57	0.13	0.01	40.1	5.1	103.2

## 2.2 Treatments

Combinations of irrigation regimes and tillage systems as main plots and nitrogen rates as subplots were evaluated in a factorial split-plot design with three replications. Tillage included conventional tillage (CT) - two discs, two cultivators followed by planking operation and deep tillage (DT) - sub-soiling/chiselling ploughed up to 45 cm deep and 50 cm apart followed by CT. Irrigation regimes comprised of no post-sowing irrigation ( $I_0$ ), one irrigation ( $I_1$ ); at 4 weeks after sowing (WAS) and two irrigations ( $I_2$ ); one at 4 WAS and the second in December end or January start. Nitrogen rates were 0 ( $N_0$ ), 50 ( $N_{50}$ ), 75 ( $N_{75}$ ) and 100 kg ha<sup>-1</sup> ( $N_{100}$ ). The gross plot size was 3.9 x 3.3 m<sup>2</sup> and the net plot size was 3.6 x 3.0 m<sup>2</sup>.

## 2.3 Crop Management

The field plots with DT treatment were deeply tilled (sub-soiled) with a tractor-drawn chiseler in the first week of October after harvesting the preceding maize crop and then the entire field was ploughed twice with a disc harrow. After heavy pre-sowing irrigation (10 cm), the field was prepared by giving two cultivations with a tractor-drawn cultivator followed by planking at proper moisture conditions to obtain a fine seed bed. Canola crop (GSC 7) was sown @ 3.75 kg ha<sup>-1</sup> at a row spacing of 45 cm on October 18, 2018. The whole amount of phosphorus (30 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> as single super phosphate) and potassium (15 kg K<sub>2</sub>O ha<sup>-1</sup> as muriate of potash) was applied at sowing. In plots with  $I_0$  irrigation regime, the full dose of nitrogen fertilizer (as urea) was applied at the time of sowing as per treatment while in plots with other irrigation regimes, 50 per cent of N as per treatment was applied. Before the first irrigation, the remaining dose of N as per treatment was applied. No nitrogen was applied in  $N_0$  plots. In  $I_1$  and  $I_2$  irrigation regimes, the first irrigation was applied on November 16 and in  $I_2$  regime, the second irrigation was applied on December 27, 2018. The measured known amount (70 mm) of irrigation water was applied using the Parshall flume. Harvesting of the crop was done manually in the first week of April.

## 2.4 Measurements

Soil cores for determining root growth were sampled at 50 per cent flowering with 0.15 m depth increments down to 1.80 m soil depth with 0.05 m diameter auger centred at 0.075 m away

from the plant base [16]. Roots from each sample were washed in a net cloth and cleaned. Root length was measured with CI-203 Area Meter and root length density (root length per unit volume of soil) was calculated. The moisture content of 0-15, 15-30, 30-60, 60-90, 90-120, 120-150 and 150-180 cm profile layers was determined at different stages of crop growth by thermo gravimetric method with three replications of each treatment. Three plants were randomly selected from each plot and plant height was measured at 64, 91, 132 and 154 DAS and maturity of the crop from the base to the tip of the plant. Chlorophyll content in leaves of intact plants was measured with Minolta – SPAD 502 Chlorophyll Meter at 49, 64, 103 and 132 DAS. For each observation, a second or third fully opened leaf from the apex was selected from ten plants for each treatment by taking the precaution that the midrib should not come under the sample area/sensor of the instrument. The mean value of 10 readings was reported as SPAD value. Relative leaf water content (RLWC) was measured at 70, 104 and 152 DAS from the fully developed 3<sup>rd</sup> leaf from the top. Among nitrogen treatments, two extremes (0 and 100 kg N ha<sup>-1</sup>) were taken for measurements. Immediately after cutting, leaves were sealed within plastic bags and quickly transferred to the laboratory. Five circular disks of 2 cm diameter were cut from each leaf and fresh weight (FW) was recorded. Turgid weight (TW) was obtained after soaking disks in distilled water in Petri plates for 24 hours at room temperature (about 20°C) and under low light conditions in the laboratory. After soaking, disks were quickly and carefully blotted dry with tissue paper in preparation for determining turgid weight. The dry weight (DW) of the disk was obtained after oven drying at 70°C till the weight becomes constant. Relative leaf water content was calculated by the formula:

$$RLWC (\%) = \frac{(FW-DW)}{(TW-DW)} \times 100$$

The number of primary and secondary branches per plant and siliquae per plant in each treatment was counted at physiological maturity from three randomly selected plants and computed as mean. The total number of seeds in them was counted from twenty-five siliquae collected randomly in each treatment at maturity just before harvesting. The mean number of seeds per siliqua was calculated. After threshing the crop, a representative sample of seeds from each treatment was collected from bulk produce of the whole plot. One thousand seeds were

counted and weighed to give a thousand seed weight. Grain and stover yield was calculated from harvested net plot area. The oil content in seed (expressed in percentage) was determined with MQC benchtop Nuclear Magnetic Resonance (NMR) Analyser (Oxford Instruments, UK) by using a non-destructive method of oil estimation as suggested by Alexander et al. [17].

## 2.5 Statistical Analysis

Treatment effects on various parameters were tested for their statistical significance using ANOVA for a factorial split-plot design. Regression analysis was done to relate the seed yield of canola with nitrogen application rate and yield attributing characters viz; primary branches, secondary branches, siliquae per plant, seeds per siliqua and 1000-seed weight.

## 3. RESULTS AND DISCUSSION

### 3.1 Soil Moisture Distribution


The distribution of soil moisture in different layers of the root zone reflects water extraction patterns by roots. In general, soil moisture content increased with an increase in soil depth due to lesser evaporation from sub-surface. A major contribution for evaporation and greater root activities were responsible for the notably lower moisture content in the upper (0-1.2 m) soil layers [18]. At 39 DAS, higher soil moisture content was observed under  $I_1$  irrigation regime than  $I_0$  by 3.5 to 1.7 per cent of soil moisture content throughout the 180 cm of soil profile (Fig. 1). Higher moisture content in  $I_1$  irrigation regime was attributed to the application of irrigation water in  $I_1$  before sampling. At 70 DAS, the difference in soil moisture content between  $I_1$  and  $I_0$  reduced throughout the soil profile with higher values obtained under  $I_1$  irrigation regime than  $I_0$ . At 84 DAS,  $I_2$  irrigation regime recorded 8.5, 8.9, 10, 10.1, 10.6, 10.8 and 11.1 per cent soil moisture content against 4.4, 5, 8.1, 9.1, 9.7, 10.1 and 10.7 per cent obtained in  $I_0$  in 0-15, 15-30, 30-60, 60-90, 90-120, 120-150 and 150-180 cm of soil depth, respectively. Higher moisture content in  $I_2$  irrigation regime than  $I_1$  and  $I_0$  was observed due to the application of irrigation water in  $I_2$  before sampling. The differences were lower among soil moisture content in upper soil layers under  $I_0$ ,  $I_1$  and  $I_2$  irrigation regimes at 153 DAS due to rainfall however, higher differences were recorded at lower depths. The results were similar to the findings of Mandal et al. [19] and Zeleke et al. [20].

Higher soil moisture content was recorded under conventional tillage as compared to deep tillage at all growth stages but most of the difference was observed in lower layers (Fig. 2). Denser rooting and increasing dry matter with deep tillage and higher irrigation frequency caused greater extraction of profile stored water. As a consequence, there was less residual water at the end of the growing season in deep-tilled plots [21]. There was no difference between soil moisture content under CT and DT at harvest due to rainfall near harvest.

In general, higher soil moisture content was recorded under  $N_0$  as compared to  $N_{100}$  at all growth stages (Fig. 3). At 39 DAS, higher values of soil moisture content were obtained under  $N_0$  than  $N_{100}$  with a maximum difference in 0-60 cm of the soil profile. Soil moisture content recorded under  $N_0$  was 4.3, 4.9, 7.6, 9.5, 10, 10.4 and 11.1 per cent against 4.8, 5.4, 7.2, 8.9, 9.3, 9.5 and 10.2 per cent under  $N_{100}$  in 0-15, 15-30, 30-60, 60-90, 90-120, 120-150 and 150-180 cm of soil depth, respectively. Soil moisture content recorded in 0-30 cm of the soil profile at 84 DAS had no difference between  $N_0$  and  $N_{100}$  but below 30 cm of soil depth difference was observed to be higher under  $N_0$  than  $N_{100}$  and thereafter, it reduced with an increase in depth. A similar trend was observed at 153 DAS and harvest. Kumar et al. [22] observed that 80 kg N ha<sup>-1</sup> recorded significantly lower soil moisture content than 40 kg N ha<sup>-1</sup> and control in mustard. Increase in nitrogen application rate provided sufficient nitrogen required for plant development enhancing the root growth and above-ground biomass. This caused the plant to extract more water from the soil profile and hence, resulting in lower moisture content in the soil. The findings were in correspondence with the study of Taylor et al. [23] and Beard et al. [24].

### 3.2 Root Length Density

Among irrigation regimes,  $I_2$  recorded the highest RLD followed by  $I_1$  and  $I_0$  in upper layers of soil up to 60 cm (Fig. 4). The mean increase in RLD under  $I_2$  irrigation regime was 0.34, 0.24 and 0.23 mm cm<sup>-3</sup> over  $I_1$  and 0.82, 0.48 and 0.44 mm cm<sup>-3</sup> over  $I_0$  in 0-15, 15-30 and 30-60 cm of soil depth, respectively. In 60-90 and 90-120 cm of soil layers,  $I_0$  irrigation regime recorded maximum RLD of 0.52 and 0.29 mm cm<sup>-3</sup> which was higher by 10 and 5.1 per cent than  $I_1$  and 16.3 and 9.4 per cent than  $I_2$ . Below 120 cm soil depth, RLD increased with an increase in the number of irrigations but the difference was



**Fig. 1. Effect of irrigation on depth-wise volumetric soil moisture content during growing season of canola**

smaller. Increased RLD with an increase in irrigation frequency conform with the findings of Arora et al. [21]. Root length density was higher in DT as compared to CT with maximum effect

observed in 60-90 cm of soil depth (Fig. 4). Averaged across nitrogen treatments, DT resulted in RLD of 1.5, 1.3, 0.7, 0.4, 0.2 and 0.1 mm cm<sup>-3</sup> in 0-15, 15-30, 30-60, 60-90, 90-120

and 120-150 cm of soil depth under unirrigated plots that was 9.6, 5.1, 18, 40, 47.3 and 15.3 per cent higher than CT, respectively. In  $I_1$  irrigation regime, the increase in RLD under DT plots over

CT was 4.8, 5, 16.9, 44.7, 45.6 and 15.9 per cent in 0-15, 15-30, 30-60, 60-90, 90-120 and 120-150 cm of soil depth, respectively. The corresponding increase in  $I_2$  irrigation regime





Fig. 2. Effect of tillage on depth-wise volumetric soil moisture content during growing season of canola



**Fig. 3. Effect of nitrogen application on depth-wise volumetric soil moisture content during growing season of canola**

was 5.1, 8.3, 13.6, 48.3, 49.4 and 24.3 per cent under DT over CT. Arora et al. [21] reported that below 60 cm depth, DT plots had higher rooting density than CT plots in mustard. Similar results were obtained by Kaur and Arora [25] in maize crop. With an increase in nitrogen dose, RLD increased being highest under N<sub>100</sub> followed by N<sub>75</sub>, N<sub>50</sub> and N<sub>0</sub>. The effect was maximum in upper soil layers and reduced with an increase in

soil depth. Little difference was observed in deeper layers of soil. Under I<sub>0</sub> irrigation regime, N<sub>100</sub> recorded 1.6, 1.4, 0.8, 0.6, 0.3 and 0.1 mm cm<sup>-3</sup> of RLD in 0-15, 15-30, 30-60, 60-90, 90-120 cm and 120-150 cm of soil depth against 1.5, 1.2, 0.7, 0.5 and 0.3 mm cm<sup>-3</sup> under N<sub>0</sub> in 0-15, 15-30, 30-60, 60-90 and 90-120 cm of soil depth, respectively. In I<sub>1</sub> and I<sub>2</sub> irrigation regimes also, N<sub>100</sub> resulted in the highest RLD as compared to



N<sub>75</sub>, N<sub>50</sub> and N<sub>0</sub>. Dreccer et al. [26] reported higher root length density with higher nitrogen dose in oilseed rape. The results were similar to the findings of Liu et al. [27] and Kaur and Arora

[25]. Better root growth may be explained that nitrogen stimulates early root development and their growth [28].




Fig. 4. Effect of (a) tillage and (b) nitrogen rates on depth-wise root length density (RLD, mm cm<sup>-3</sup>) of canola under different irrigation regimes

### 3.3 Plant Height

Irrigation regimes and nitrogen rates significantly influenced the plant height recorded at various growth stages (Table 2). A significant increase in plant height was observed with two irrigations at all growth stages over one irrigation and no post-sowing irrigation. Maximum mean plant height recorded was 102.9 cm under  $I_2$  which was significantly higher than  $I_1$  (95.3 cm) and  $I_0$  (80.1 cm) at 91 DAS and corresponding values of about 155, 135 and 111 cm at 132 DAS, 167, 149 and 144 cm at 154 DAS and 188, 170 and 155 cm at maturity, respectively were obtained. Increased plant height with the application of irrigation water would have been due to the availability of higher soil moisture resulting thereby higher leaf water potentials, stomatal conductance, absorption and translocation of nutrients which ultimately reflected in the healthy growth of the crop. Saud et al. [29] recorded significantly higher plant height with one irrigation (at the flowering stage) than no post-sowing irrigation in Indian mustard. Maximum mean plant height of 182 cm was obtained with  $N_{100}$  followed by  $N_{75}$  (179 cm),  $N_{50}$  (175 cm) and  $N_0$  (166 cm) at maturity. A similar trend was also observed at other growth stages. Pattam et al. [30] and Khan et al. [31] also recorded an increase in plant height with an increase in nitrogen dose.

### 3.4 Leaf Chlorophyll Content (SPAD Value)

The SPAD value increased up to 103 DAS and decreased thereafter to 132 DAS (Table 2). With an increase in the number of irrigations, SPAD value increased significantly at all growth stages with a mean maximum value of 51.4 obtained with two irrigations followed by 49.3 and 46.7 with one irrigation and no irrigation, respectively at 103 DAS (Table 2). The improvement in chlorophyll content of leaves under various irrigation levels was probably due to better plant growth and N uptake. It is a fact that periodical water stress leads to many anatomical changes like a decrease in the size of cell and inter-cellular space, thicker cell wall and decrease in chlorophyll content and its stability but sufficient moisture condition reverses these anatomical changes by enhancing the turgidity of cells and uptake of nutrients which in turn affects the synthesis of chlorophyll. Sabagh et al. [32] reported similar results. Under deep tillage, SPAD value was significantly higher than that under conventional tillage and the improvement was 1.6, 1.5, 1.6 and 0.9 at 49, 64, 103 and 132

DAS, respectively. Nitrogen application significantly increased leaf chlorophyll content with the maximum value obtained with 100 kg N  $ha^{-1}$  at all growth stages. One of the main components of chlorophyll synthesis is nitrogen. Chlorophyll content increases with an increase in nitrogen availability to the plant resulting in higher SPAD value in plots with higher nitrogen rates and more irrigations. Application of irrigation water and deep tillage maintained optimum soil moisture status in the root zone that accelerated the nitrogen uptake by the plant. SPAD value was highest (46.87) and lowest (40.1) with the application of N at the rate of 270 and 90 kg  $ha^{-1}$  as reported by Shengri et al. [33].

### 3.5 Relative Leaf Water Content (RLWC)

Relative leaf water content (RLWC) increased linearly with an increase in growth and the highest RLWC was recorded at 152 DAS (Table 3). Irrigation had a significant effect on RLWC at all growth stages as depicted in Table 3. At 70 DAS, one irrigation resulted in 3.7 per cent significantly higher RLWC than no irrigation. Two irrigations significantly increased the RLWC by 4.5 and 8.3 per cent over one irrigation and no irrigation, respectively at 104 DAS. The corresponding increment recorded at 152 DAS was 1.9 and 3.4 per cent, respectively. The results indicated that an increase in irrigation frequency led to increased relative leaf water content. The high RLWC is associated with higher dry matter production rates because cell turgidity is important with the opening and closing of stomata, expansion of leaves and the movement of water and nutrients to the various parts of the plant [34]. Relative water content as well as osmotic pressure decreased significantly by prolonging irrigation intervals up to 15 days [35]. The relative water content of canola significantly reduced with increasing intervals of irrigation to 45 days as observed by Sabagh et al. [32]. Nitrogen application resulted in a significant increase in RLWC throughout the growing period of the crop (Table 3).  $N_{100}$  recorded 79.4, 83 and 87.4 per cent RLWC over  $N_0$  (77.9, 80.7 and 85.4%) at 70, 104 and 152 DAS. The highest value of RLWC was recorded in 100 kg N  $ha^{-1}$  (10.87% increase over the control) as reported by Namvar and Khandan [36]. It has been shown that increasing nitrogen application will increase protein synthesis, increase cell wall thickness and cause absorption of extra water by protoplasm and improve the relative water content [37]. Tillage had no significant effect on the RLWC of canola.

**Table 2. Effect of irrigation, tillage and nitrogen rates on plant height and SPAD value of canola at various growth stages**

Treatment	Plant height (cm)				At maturity	SPAD			
	64 DAS	91 DAS	132 DAS	154 DAS		49 DAS	64 DAS	103 DAS	132 DAS
<b>Irrigation</b>									
I <sub>0</sub>	18.4	80.1	110.6	144.1	154.1	42.7	43.9	46.7	44.6
I <sub>1</sub>	28.5	95.2	134.0	149.4	169.8	44.6	46.4	49.3	46.7
I <sub>2</sub>	29.1	102.9	154.9	166.7	188.3	44.6	46.4	51.4	49.1
CD (0.05)	2.7	4.5	11.2	16.7	7.5	1.3	1.5	1.2	1.4
<b>Tillage</b>									
CT	24.8	92.5	132.5	152.6	170.4	43.2	44.8	48.3	46.3
DT	25.8	93.0	133.9	154.2	171.0	44.7	46.3	50.0	47.3
CD (0.05)	NS	NS	NS	NS	NS	1.0	1.2	1.0	NS
<b>Nitrogen rate</b>									
N <sub>0</sub>	18.9	83.2	117.2	143.6	161.7	40.7	42.3	45.8	43.7
N <sub>50</sub>	25.4	91.6	134.6	151.4	169.8	43.5	44.5	48.2	46.0
N <sub>75</sub>	27.6	96.9	138.5	157.2	173.7	44.9	46.7	50.6	48.0
N <sub>100</sub>	29.3	99.1	142.3	161.3	177.8	46.7	48.7	51.9	49.5
CD (0.05)	1.2	2.4	4.6	3.9	2.8	1.1	1.3	0.9	0.7

**Table 3. Effect of irrigation, tillage and nitrogen rates on relative leaf water content (RLWC, %) of canola**

Treatment	RLWC (%)		
	70 DAS	104 DAS	152 DAS
<b>Irrigation</b>			
I <sub>0</sub>	76.6	77.9	84.6
I <sub>1</sub>	79.4	80.7	85.9
I <sub>2</sub>	79.3	84.4	87.5
CD (0.05)	1.7	0.8	2.2
<b>Tillage</b>			
CT	78.2	80.9	85.6
DT	78.6	81.1	86.4
CD (0.05)	NS	NS	NS
<b>Nitrogen rate</b>			
N <sub>0</sub>	77.7	79.9	84.9
N <sub>100</sub>	79.2	82.1	87.1
CD (0.05)	0.6	0.7	0.7

### 3.6 Yield Attributes and Yield

#### 3.6.1 Number of primary and secondary branches

Irrigation regimes, tillage and nitrogen rates significantly affected the production of primary and secondary branches per plant (Table 4). Two irrigations produced the maximum number of primary (5.2) and secondary branches (8.2) followed by one irrigation (4.0 and 7.0) and no post-sowing irrigation (3.3 and 6.3), respectively. Shivran et al. (2018) reported that three irrigations (30-35 DAS, flowering and silique development) produced the highest number of

primary branches in mustard. Two irrigations (pre-bloom stage + pod filling stage) produced a significantly higher number of primary and secondary branches per plant than one irrigation (pre-bloom stage) in mustard [38]. A significant effect of irrigation regimes on secondary branches was also reported by Rathore et al. [39] in Indian mustard. This might be due to the reason that better water content in plants enhances growth and better growth produces a higher number of branches. Deep tillage increased the production of primary branches by 18 per cent and secondary branches by 10.3 per cent over conventional tillage. The number of branches plant<sup>-1</sup> declined significantly with

decreasing the intensity of tillage in Indian mustard [40]. Better soil moisture status promoted better root growth resulting in improved shoot growth and a higher number of branches. This is in confirmation with the finding of Mondal et al. [41] and Belal [42]. The plots applied with  $N_{100}$  resulted in the highest number of primary branches with a mean value of 5.9 which was 1.2, 2.2 and 3.6 higher than  $N_{75}$ ,  $N_{50}$  and  $N_0$ . A similar trend was observed for secondary branches. This agreed with the findings of Akbar et al. [43] who reported that the addition of 100 kg N ha<sup>-1</sup> produced a maximum (11) number of branches plant<sup>-1</sup>, while a minimum (5) number of branches plant<sup>-1</sup> in control plots of canola crop. These results are in line with those reported by Kumar et al. [22]. The reason could be more vegetative growth with increasing nitrogen level which results in more number of branches plant<sup>-1</sup> in plots with 100 kg N ha<sup>-1</sup> as compared to control plots.

### 3.6.2 Number of siliquae per plant and grain per siliqua

Two irrigations produced significantly more siliquae per plant (371) and grains per siliqua (23.3) as compared to one irrigation (351, 21.2) and no irrigation (328, 19.5), respectively (Table 4). The favourable effect of two irrigations on the sink component (number of siliquae and number

of grains) could be attributed to better development of the plants in terms of plant height, number of branches and dry biomass production leading to increased bearing capacity due to optimum growth on account of favourable moisture during entire crop growing period [44,45]. Tyagi and Upadhyay [46] observed that the application of two irrigations at 40 and 75 DAS produced significantly the highest number of siliquae plant<sup>-1</sup> (101.9) and several grains siliqua<sup>-1</sup> (12) followed by one irrigation at 40 DAS and the lowest under no post sowing irrigation in Indian mustard. Similar findings were also reported by Jat et al. [47] and Bharat et al. [48]. Plants in deep-tilled plots resulted in 22.3 grains per siliqua which was significantly higher than in conventional tilled plots (20.3 grains per siliqua) whereas for several siliquae per plant, the value was numerically higher but could not be reached at a level of significance. Poddar and Kundu [49] reported that with an increase in tillage intensity increased number of seeds per siliqua. The nitrogen rates of  $N_{100}$ ,  $N_{75}$  and  $N_{50}$  resulted in a significant increase in the number of siliquae per plant by 26, 16 and 8 per cent over  $N_0$  and the number of seeds per siliqua by 25, 19 and 7 per cent over  $N_0$ . Increasing nitrogen application rate resulted in the production of a higher number of siliquae per plant [50]. Similar results were reported by Tufail et al. [51] and Islam et al. (2018) for number of seeds per siliqua.

**Table 4. Effect of irrigation, tillage and nitrogen rates on yield attributes, yield and oil content of canola**

Treatment	Primary Branches	Secondary Branches	Siliqua per plant	Grain per siliqua	1000 grain weight (g)	Grain yield (t ha <sup>-1</sup> )	Stover yield (t ha <sup>-1</sup> )	Oil content (%)
<b>Irrigation</b>								
$I_0$	3.26	6.26	328	19.5	2.71	1.41	5.18	42.31
$I_1$	3.98	6.98	351	21.2	3.35	1.60	5.87	42.44
$I_2$	5.24	8.24	371	23.3	3.88	1.76	6.43	42.56
<b>CD (0.05)</b>	0.23	0.25	28	1.1	0.17	0.12	0.43	NS
<b>Tillage</b>								
<b>CT</b>	3.82	6.82	344	20.3	3.24	1.52	5.56	42.41
<b>DT</b>	4.50	7.50	356	22.3	3.39	1.66	6.09	42.47
<b>CD (0.05)</b>	0.19	0.21	NS	0.9	0.14	0.10	0.35	NS
<b>Nitrogen rate</b>								
$N_0$	2.28	5.28	310	18.9	2.82	1.08	4.05	42.63
$N_{50}$	3.69	6.70	334	20.3	3.24	1.56	5.67	42.52
$N_{75}$	4.74	7.74	361	22.4	3.50	1.74	6.34	42.41
$N_{100}$	5.93	8.93	394	23.7	3.69	1.97	7.23	42.19
<b>CD (0.05)</b>	0.30	0.31	8	0.7	0.08	0.10	0.29	0.09

### 3.6.3 Thousand seed weight

Significantly higher mean thousand-grain weight was recorded with two irrigations (3.88 g) in comparison to one irrigation (3.35 g) and no post-sowing irrigation (2.71 g) as shown in Table 4. Bharat et al. [48] reported that with an increase in irrigation amount, 1000 grain weight increased in Indian mustard. Deep tillage (3.39 g) resulted in a significantly higher thousand-grain weight than CT (3.24 g). Sub-soiling resulted in 8 per cent higher 1000 grain weight as compared to conventional tillage in soybean [52]. The effect of nitrogen rates on 1000 grain weight was significant with the maximum mean value obtained with N<sub>100</sub> (3.69 g) followed by N<sub>75</sub> (3.50 g), N<sub>50</sub> (3.24 g) and least with N<sub>0</sub> (2.82 g). The maximum 1000 grain weight was observed from the plot fertilized with 70 kg ha<sup>-1</sup>, while the minimum at 40 kg ha<sup>-1</sup> as reported by Ali et al. 2019. The results are in line with Hamid and Shaheen [53] who reported that proper nitrogen level increases the weight of grain due to high uptake of nitrogen which results in heavier grain.

### 3.6.4 Grain and stover yield

Irrigation, tillage and nitrogen rates had substantial effects on the grain yield of canola (Table 4). Two irrigations recorded 10 and 9.5 per cent significantly higher grain and stover yield than one irrigation (1.6 t ha<sup>-1</sup> and 1.4 t ha<sup>-1</sup>) and 25 and 24 per cent over no post-sowing irrigation (1.4 t ha<sup>-1</sup> and 5.2 t ha<sup>-1</sup>), respectively. This might be due to higher photosynthesis and translocation of assimilates toward reproductive structure owing to sufficient soil moisture. The results conformed with Ray et al. [54] and Shivran et al. [55]. The DT plots registered an increase of 9.2 and 9.5 per cent higher seed and stover yield over CT, respectively. Deep tillage provided better root growth and better moisture extraction that helped the crop initially develop an adequate source (as reflected by high biomass accumulation) as compared to CT. Pal and Phogat [56] concluded from their experiment that deep ploughing significantly increased the

seed yield of mustard over conventional tillage. An increase in nitrogen dose from N<sub>0</sub> to N<sub>100</sub> also increased grain yield significantly. N<sub>100</sub> recorded a significantly higher grain yield of 1.97 t ha<sup>-1</sup> than N<sub>75</sub> (1.74 t ha<sup>-1</sup>), N<sub>50</sub> (1.56 t ha<sup>-1</sup>) and N<sub>0</sub> (1.08 t ha<sup>-1</sup>). Increasing N rates improved grain yield as reported by Ali et al. [57] in canola.

### 3.6.5 Oil content

Significant results were obtained for different nitrogen rates while irrigation and tillage had no significant effect (Table 4). Two irrigations resulted in 42.56 per cent of oil content which was comparable with one irrigation (42.44 per cent) and no irrigation (42.31 per cent). The oil content of rain-fed canola and mustard was 7% and 5% lower than the irrigated ones, respectively as observed by Zeleke et al. [20]. A significant effect of several irrigations on the oil content of Indian mustard was also observed by Shivran et al. [55]. Oil content decreased with an increase in nitrogen dose. The highest nitrogen dose of 100 kg ha<sup>-1</sup> resulted in a significantly lower oil content of 42.19 per cent against 75 kg N ha<sup>-1</sup> (42.41%), 50 kg N ha<sup>-1</sup> (42.52%) and 0 kg N ha<sup>-1</sup> (42.63%). The possible reason for the decrease in oil content with N increase may be because N is the major constituent of protein so it might increase the percentage of grain protein, as a result, there might be a decrease in the percentage of oil content since it has an inverse relationship with protein [58]. The results agree with those documented by Olama et al. [59] and Khan et al. [60].

### 3.6.6 Yield attributing parameters and seed yield relationship

The mean value of different yield attributing parameters like primary branches, secondary branches, siliquae per plant, seeds per siliqua and 1000- grain weight were recorded and correlated with the grain yield of canola. The regression model was developed with these parameters and the grain yield of canola (Table 5). Among these parameters, the maximum

**Table 5. Yield attributing parameters and grain yield relationship**

Dependent variable (Y)	Independent Variable (X)	Regression equation	R <sup>2</sup>
Seed yield	Primary branches	4.18X-5.83	0.780*
do	Secondary branches	2.12X+0.55	0.887*
do	Siliquae per plant	0.10X-18.31	0.897*
do	Grain per siliqua	1.21X-10.19	0.773*
do	1000- grain weight	5.21X-1.69	0.683*

\* Significance at 5% level

variation in seed yield was explained by siliquae per plant (89.7%) followed by number of secondary branches (88.7%), number of primary branches (78.0%), seeds per siliqua (77.3%) and 1000- grain weight (68.3%).

#### 4. CONCLUSION

This study demonstrated that frequent irrigation (two irrigations) and higher nitrogen rate (100 kg N ha<sup>-1</sup>) along with deep tillage significantly increased the growth, yield attributes, yield and oil content of canola oilseed rape. Two irrigations produced 10 per cent higher mean grain yield over one irrigation and 25 per cent over no irrigation. Deep tillage incremented seed yield by 9.2 per cent over conventional tillage and 100 kg N ha<sup>-1</sup> produced the highest grain yield. The CT plots retained more soil moisture than DT plots and among N levels, N<sub>0</sub> recorded higher soil moisture content throughout the crop growing season. Root density in upper 60 cm soil depth was higher in I<sub>2</sub> followed by I<sub>1</sub> and I<sub>0</sub> irrigation regimes whereas below 60 cm, it was higher under I<sub>0</sub>. Higher root density was recorded under DT and N<sub>100</sub> plots. It may be concluded that for higher productivity canola can be grown under deep tillage with sufficient irrigation (I<sub>2</sub>) and N fertilization (N<sub>100</sub>). The study demonstrated that frequent irrigation (two irrigations), higher nitrogen rate (100 kg N ha<sup>-1</sup>) along with deep tillage had no negative influence on plant establishment and have likelihood of crop yield improvement for irrigated canola production in Punjab.

#### ACKNOWLEDGEMENT

The authors are grateful to Punjab Agricultural University, Ludhiana, Punjab, India for providing funds for completion of this research project of M. Sc. thesis.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Jat RK, Sapkota TB, Singh RG, Jat ML, Kumar M, Gupta RK. Seven years of conservation agriculture in a rice-wheat rotation of Eastern Gangetic Plains of South Asia: yield trends and economic profitability. *Field Crops Res.* 2014; 164:199–210.

2. Singh VV, Bhagirath R, Singh M, Meena ML, Chauhan JS. Generation mean analysis for water stress tolerance parameters in Indian mustard (*Brassica juncea* (L.) Czern & Coss) Crosses. *J Breeding Genetics.* 2014;46(1):76-80.
3. Khodabin G, Lightburn K, Hashemi SM, Moghada MS, Jalilian A. Evaluation of nitrate leaching, fatty acids, physiological traits and yield of rapeseed (*Brassica napus*) in response to tillage, irrigation and fertilizer management. *Plant Soil.* 2022; 473(1-2):423-40.
4. Anonymous, Package of practices for Rabi Crops of Punjab. pp 44. Punjab Agricultural University, Ludhiana; 2018.
5. Anonymous, 2019. Production and productivity of rapeseed and mustard in India. Available:<http://www.indiastat.com>
6. Bhattarai M. Seed Yield, Oil and Protein Content, and Pollinator Potential of Winter Canola (Doctoral dissertation, Tennessee State University); 2023.
7. Sohrabi T, Movahhedi DM, Salehi A, Balouchi HR. Effect of Nitroxin Bio-Fertilizer Application Accompanied with Nitrogen on Canola (*Brassica napus*) Yield Quantity and Quality. *Isfahan University Technol. J. Crop Prod. Process.* 2023;10: 12(4):65-81.
8. Hirzel J, Undurraga P, León L, Matus I. Effect of three crop rotations and four residue levels on canola and bean grain yield and residue production. *J. Agric. Sci.* 2023;161(2):272-278.
9. Conyers MK, Moroni JS, Poile GJ, Oates AA, Lowrie R, Swan AD, et al. Response of canola and cereals to amendment of subsurface soil acidity and a hardpan. *Crop Pasture Sci.* 2023.;2: 45-67.
10. Pecan U, Pintar M, Kastelec D. Variability of in situ soil water retention curves under different tillage systems and growing seasons. *Soil Tillage Res.* 2023;233: 105779.
11. Ahmad N, Rashid M, Vaes AG. Fertilizer and their use in Pakistan, 2nd ed. NFDC. Pub.No. 4/96, Islamabad. 1996;274.
12. Schneider F, Don A, Hennings I, Schmittmann O, Seidel SJ. The effect of deep tillage on crop yield—What do we really know? *Soil Till Res.* 2017;174:193-204.
13. Yadav RP, Tripathi ML, Trivedi SK. Yield and quality of Indian mustard (*Brassica juncea*) as influenced by irrigation and

- nutrient levels. Indian J Agron. 2010; 55(1):56-59.
14. Malagoli P, Laine P, Rossato L, Ourry A. Dynamics of nitrogen uptake and mobilization in field grown winter oilseed rape (*Brassica napus*) from stem extension to harvest. Annals Bot. 2005;95: 853-861.
  15. Malhi SS, Lemke R. Tillage, crop residue and N fertilizer effects on crop yield, nutrient uptake, soil quality and nitrous oxide gas emissions in a second 4-yr rotation cycle. Soil Till Res. 2007;96:269-283.
  16. Gajri PR, Arora VK, Kumar K. A procedure for determining average root length density in row crops by single-site augering. Plant Soil. 1994;160:41-47.
  17. Alexander DE, Silvela L, Collins SFI, Rodgers RC. Analysis of oil content in maize by wide line NMR. J Am Oil Chem Soc. 1967;44:555-558.
  18. Sarkar S, Rana SK. Role of tillage on productivity and water use pattern of rice-wheat cropping system. J Indian Soc Soil Sci. 1999;47(3):532-534.
  19. Mandal KG, Hati KM, Misra AK, Bandyopadhyay KK. Root biomass, crop response and water-yield relationship of mustard (*Brassica juncea* L.) grown under combinations of irrigation and nutrient application. Irrig Sci. 2010;28:271-280.
  20. Zeleke KT, Lucketts DJ, Cowley RB. Response of canola (*Brassica napus* L.) and mustard (*B. juncea* L.) to different watering regimes. Exp Agric. 2014; 50(4):573-590.
  21. Arora VK, Gajri PR, Chaudhary MR. Effects of conventional and deep tillage on mustard for efficient water and nitrogen use in coarse textured soils. Soil Till Res. 1993;26:327-340.
  22. Kumar K, Kumar Y, Katiyar NK. Effect of plant geometry, nitrogen level and antitranspirants on physiological growth, yield attributes, WUE and economics of mustard (*Brassica juncea*) under semi-arid conditions of western Uttar Pradesh. J Pharmacogn Phytochem. 2018;7(2):226-229.
  23. Taylor AJ, Smith CJ, Wilson IB. Effect of irrigation and nitrogen fertilizer on yield, oil content, nitrogen accumulation and water use of canola (*Brassica napus* L.). Fertilizer Res. 1991;29:249-260.
  24. Beard T, Maaz T, Borrelli K, Harsh J, Pan W. Nitrogen affects wheat and canola silica accumulation, soil silica forms, and crusting. J Environ Qual. 2018;47:1380-1388.
  25. Kaur R, Arora VK. Deep tillage and residue mulch effects on productivity and water and nitrogen economy of spring maize in north-west India. Agric Water Manage. 2019;213:724-731.
  26. Dreccer MF, Schapendonk AHCM, Slafer GA, Rabbinge R. Comparative response of wheat and oilseed rape to nitrogen supply: absorption and utilization efficiency of radiation and nitrogen during the reproductive stages determining yield. Plant Soil. 2000;220:189-205.
  27. Liu W, Ma G, Wang C, Wang J, Lu H, Li S, Feng W, Xie Y, Ma D, Kang G. Irrigation and nitrogen regimes promote the use of soil water and nitrate nitrogen from deep soil layers by regulating root growth in wheat. Front Plant Sci. 2018; 9:32.
  28. Gour R, Kushwaha HS, Mangal S. Optimization basal and foliar application of nitrogen for enhancing productivity of mustard [*Brassica juncea* (L.) Czern and Coss]. Int J Chem Studies. 2019;7(1): 1553-1556.
  29. Saud RK, Singh BP, Pannu RK. Effect of limited irrigation and nitrogen levels on growth, yield attributes and yield of Indian mustard (*Brassica juncea* L.). Agric Sci Digest. 2016;36(2):142-145.
  30. Pattam K, Pannu RK, Dhaka AK, Sharma KD. Effect of dates of sowing and nitrogen levels on growth and yield of Indian mustard. Int J Curr Microbiol App Sci. 2017;6(9):1029-1036.
  31. Khan S, Anwar S, Kuai J, Ullah S, Fahad S, Zhou G. Optimization of nitrogen rate and planting density for improving yield, nitrogen use efficiency and lodging resistance in oilseed rape. Front Plant Sci. 2017;8:532.
  32. Sabagh AEL, Abdelaal KAA, Barutcular C. Impact of antioxidants supplementation on growth, yield and quality traits of canola (*Brassica napus* L.) under irrigation intervals in north Nile delta of Egypt. J Exp Biol Agric Sci. 2017;5(2):163-172.
  33. Shengri AH, Gharineh MH, Bakhshandeh A, Fathi G, Karmollachaab A. Effect of terminal heat stress and nitrogen on morphophysiological characteristics, ANUE and yield of canola (*Brassica napus* L.). Int J Agric Crop Sci. 2012.4:284-288.
  34. Begum FA, Paul NK. Influence of soil moisture on growth, water use and yield of

- mustard. J Agron Crop Sci. 1993;170:136-141.
35. Leithy SM, Leila BA, Abdallah EF, Gaballah MS. Response of canola plants to antitranspirant levels and limited irrigation. Am Eurasian J Sustain Agric. 2015;9(4):83-87.
  36. Namvar A, Khandan T. Inoculation of rapeseed under different rates of inorganic nitrogen and sulfur fertilizer: impact on water relations, cell membrane stability, chlorophyll content and yield. Archives Agron Soil Sci. 2015;61(8):1137-1149.
  37. Saneoka H, Moghaieb REA, Premachandra GS, Fujita K. Nitrogen nutrition and water stress effects on cell membrane stability and leaf water relations in *Agrostis palustris* Huds. Env Exp Bot. 2004;52:131-138.
  38. Singh PK, Singh AK, Singh RK. Effect of different dates of sowing and irrigation scheduling on growth and yield of mustard (*Brassica juncea* L.). Res Environ Life Sci. 2016;9(2):200-202.
  39. Rathore SS, Shekhawat K, Dass A, Premi OP, Rathore BS, Singh VK. Deficit irrigation scheduling and superabsorbent polymer-hydrogel enhance seed yield, water productivity and economics of Indian mustard under semi-arid ecologies. Irrig Drain. 2019;68:531-541.
  40. Mishra J, Singh RK, Yadaw D, Das A, Sahoo S. Effect of tillage and irrigation frequency on growth of Indian mustard [*Brassica juncea* (L.) Czernj and Cosson]. Int J Chemical Studies. 2019;7(1):2127-2130.
  41. Mondal N, Hossain S, Bhuiya S. Tillage and mulching effects on conservation of residual soil moisture, yield attributes and yield of mustard (cv. Daulat.). Bangladesh J Agric Res. 2008;33(4):597-606.
  42. Belal M. Effect of irrigation and sowing method on yield and yield attributes of mustard. Rajshahi Univ J Life Earth Agric Sci. 2013;41:65-70.
  43. Akbar H, Tufail M, Ali S, Jan A. Nitrogen use efficiency and morpho-phenological traits of canola (*Brassica napus* L.) as influenced by shoots cutting and nitrogen levels. EC Agric. 2016;2:530-535.
  44. Chauhan DR, Ram M, Singh I. Response of Indian mustard to irrigation and fertilization with varicose sources and levels of sulphur. Indian J Agron. 2002; 47:422-426.
  45. Pirri I, Sharma SN. Effect of sulphur on yield attributes and yield of Indian mustard (*B. juncea*) as influenced by irrigation. Indian J Agric Sci. 2007;77:188-190.
  46. Tyagi PK, Upadhyay AK. Growth, yield and water use efficiency of Indian mustard (*Brassica juncea* L.) as influenced by irrigation frequency and row spacing. J Oilseed Brassica. 2017;8(1):27-36.
  47. Jat AL, Rathore BS, Desai AG, Shah SK. Production potential, water productivity and economic feasibility of Indian mustard (*Brassica juncea*) under deficit and adequate irrigation scheduling with hydrogel. Indian J Agric Sci. 2018; 88(2):212-215.
  48. Bharat R, Kumar J, Rai SK, Gupta R. Effect of hydrogel and irrigation scheduling on water use efficiency and productivity of Indian mustard (*Brassica juncea* L.) in Jammu region. J Oilseed Brassica. 2019;10(2):63-66.
  49. Poddar P, Kundu AL. Influence of tillage systems and different dates of sowing on growth, productivity and economics of Rabi rapeseed after Kharif paddy. Environ Ecol. 2007;25(3):644-647.
  50. Ren T, Liu B, Lu J, Deng Z, Li X, Cong R. Optimal plant density and N fertilization to achieve higher seed yield and lower N surplus for winter oilseed rape (*Brassica napus* L.). Field Crops Res. 2017;204:199-207.
  51. Tufail M, Akbar H, Ali S, Jan A, Khan A. Nitrogen levels and shoots cutting influenced on oil contents, yield and yield attributes of Canola. Pure App Bio. 2015;4(1):31-37.
  52. Ghosh PK, Mohanty M, Bandyopadhyay KK, Painuli DK, Misra AK. Growth, competition, yield advantage and economics in soybean/pigeon pea intercropping system in semi-arid tropics of India: I. Effect of subsoiling. Field Crops Res. 2006;96(1):80-89.
  53. Hamid and Shaheen MA. Response of seed yield, yield components and oil content of canola and nitrogen fertilizer rate diversity. Environ Arid Land Agric Sci. 2007.;20:21-31.
  54. Ray K, Sengupta K, Pal AK, Banerjee H. Effects of sulphur fertilization on yield, S uptake and quality of Indian mustard under varied irrigation regimes. Plant Soil Environ. 2015;61(1):6-10.
  55. Shivran H, Kumar S, Tomar R, Chauhan GV. Effect of irrigation schedules on



- productivity and water use efficiency in Indian mustard (*Brassica juncea* L.). Int J Chem Studies. 2018;6(4):15-17.
56. Pal D, Phogat VK. Effect of deep tillage and gypsum on yield and N, P, K and S uptake by mustard. J Indian Soc Soil Sci. 2005;53(1):134-136.
57. Ali W, Anwar S, Khan MH, Iqbal J, Kamal A, Ahmad Z. Canola yield response to levels and application methods of nitrogen. Agrotechnol. 2019;8:189.
58. Cheema MA, Malik MA, Hussain A, Shah SH, Basra MA. Effect of time and rate of nitrogen and phosphorus application on the growth and the seed and oil yield of canola (*Brassica napus* L.). J Agron Crop Sci. 2001;186:103-110.
59. Olama V, Ronaghi A, Karimian N, Yasrebi J, Hamidi R, Tavajjoh M, Kazemi MR. Seed quality and micronutrient contents and translocations in rapeseed (*Brassica napus* L.) as affected by nitrogen and zinc fertilizers. Archives Agron Soil Sci. 2014; 60(3):423-435.
60. Khan S, Anwar S, Kuai J, Noman A, Shahid M, Din M, Ali A, Zhou G. Alteration in yield and oil quality traits of winter rapeseed by lodging at different planting density and nitrogen rates. Scientific Reports. 2018;8:634.

© 2023 Kaur et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/102670>