



Effects of Nano Chitosan and Bio Capsule on Growth, Yield and Quality of Chilli (*Capsicum annuum* L.) in Poly-House Condition

Y. Pabitri Devi ^a[⊙]* and Samir E. Topno ^a[#]

^a *Department of Horticulture, Naini Agricultural Institute, Faculty of Agriculture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, U. P., India.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i232473

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92851>

Original Research Article

Received 06 August 2022
Accepted 12 October 2022
Published 15 October 2022

ABSTRACT

A field experiment was designed in a Randomized Block Design with 9 treatments replicated thrice on Chilli (*Capsicum annuum* L.) under poly-house condition. The Experiment examined the effect of nano chitosan, bio capsule and NPK @RDF application on growth, yield and quality of chilli and to work out the economics of various treatments. Various treatment combinations were prepared using different doses of Nano chitosan and Bio capsules in combination. It was observed that the treatment T₈ (NPK + Nano chitosan 120 ppm + Bio capsule 120 ppm) was superior over other treatments in terms of growth, yield and quality of Chilli i.e., plant height (63.73 cm), fruit yield per 200m² (3.27q), T.S.S. (9.67 ° Brix) and ascorbic acid (143.96 mg/100 g). Therefore, use of Nano chitosan @ 120 ppm along with 500 ppm Bio capsule could be used for getting better yield in chilli. Moreover, application of nano chitosan and bio capsule on nutritional quality of chilli could be studied in details further.

Keywords: *Nano chitosan; bio capsule; chilli; benefit cost ratio.*

[⊙] Post-Graduate Student;

[#] Assistant Professor;

*Corresponding author: E-mail: yumnampabitri1998@gmail.com;

1. INTRODUCTION

Chilli (*Capsicum annuum* L.) belongs to the family Solanaceae having diploid species with mostly $2n=24$ chromosomes but wild species with $2n=2x=26$ chromosomes have been reported [1,2]. Chilli was introduced into Europe in 1493 by Christopher Columbus, who discovered it in tropical America. Believed to be a native of Mexico and Peru, it was widely used by the people of Central and South America prior to Columbus's discovery. It spread so quickly that by 1542, three types of chilli were already introduced into India. India, Mexico, Japan, Ethiopia, Uganda, Nigeria, Thailand, Turkey, Indonesia, China and Pakistan are the major chilli growing countries. To some extent, it is also grown in Italy, Spain and the United States. India is the largest producer and consumers of chilli in the world. In India chilli is grown on an area of 7.75 lakh ha with production of 14.92 lakh tones (Anonymous 2014). Chilli are varieties of the berry-fruit of the plants from the genus capsicum. They are cultivated for their pungency (spiciness). Chilli peppers are widely used in many cuisines as a spice to add "heat" to dishes. Capsaicin and related compounds known as capsaicinoids are the substances giving chilli peppers their intensity when ingested or applied topically. Other varieties of capsicum include bell peppers (UK: sweet peppers), but while chilli peppers are (to varying degrees) pungent or "spicy", bell peppers are generally not and provide additional sweetness and flavor to a meal rather than "heat." Chilli is one of the most valuable crops in India. The crop is grown largely for its fruits all over India as a principal ingredient of various curries, and chutneys. It is used for vegetables, spices, condiments, sauces, and pickles. Dry chillies are used for curry powder. The red colour in chilli is due to "capsanthin". Pungency in chillies is due to the active constituent "capsaicin", an alkaloid is extracted from chillies and used to the medicine. Chilli's vitamins include vitamin C (109 mg), vitamin B6 (0.1mg), vitamin A (530 IU), and minerals like iron (0.5mg), copper (0.1mg), potassium (153 mg). It also contains amino acids making it a high nutritional value food. Green chillies are rich in dietary fibres and also contain zero cholesterol. As per studies, 100gms of chilli contains; total fat (0.1g), sodium (3.2mg), magnesium (11.2mg), carbohydrates (1.8g), dietary fibre (0.7g), glucose (0.8g), protein (0.9g), carbohydrate (4.3g), calcium (8.1mg), water content (39.5g) (USDA, 2017).

Polybags are lower cost alternatives to plastic pots. They are economical and easy to use by the residential greenhouse grower. These heavy-duty re-usable grow bags are made from a durable plastic that provides for a longer life. The bags have pre-punched drain holes and they are self-standing when filled with media. Widely used in greenhouse drip irrigation applications, they work very well with most all mediums and are excellent for bedding plants, tree seedlings, tomatoes, bell peppers, cucumbers, etc.

Chitosan is the N-acetyl derivative of chitin obtained by N-deacetylation. Chitosan is widely used in food and bioengineering industries for encapsulation of active food ingredients, enzyme immobilization, as a carrier for controlled drug delivery, in agriculture as a plant growth promoter. Chitosan is also a defense elicitor and an antimicrobial agent. Chitosan has interesting properties such as biodegradability, biocompatibility, bioactivity, nontoxicity and polycationic nature (Divya and Jisha, 2017). Nano chitosan has broad antimicrobial activity against fungal pathogens however, the bulk size limits its solubility which affects the antimicrobial property. Chitosan nanoparticles have great potential over the bulk counter parts as size can alter several properties compare to bulk material. The exclusive properties of these materials, such as a large surface area and greater reactivity, have also raised concerns about adverse effects on environmental health. Recently, IISR-ICAR (Indian Council of Agricultural Research) scientists have developed the technology to pack bio-fertilizers in tiny capsules. This eliminates the need for farmers to carry the sacks of biofertilizers. It consists of a carrier medium rich in live microorganisms. When applied to seed, soil or living plants, it increases soil nutrients or makes them biologically available. Easy and reliable technology of storing and delivering GPR bioagents in hard gelatin capsule termed as Bio capsule. It uses a select combination of beneficial microorganisms such as Trichoderma, Pseudomonas and Bacillus. They form a mutually beneficial or symbiotic relationship with host plants as they grow in the soil.

2. MATERIALS AND METHODS

The experiment was conducted in Randomized Block Design comprising 9 treatments replicated thrice during August 2021- February 2022 at Jacob Institute of Bio-technology and Bio-engineering (JIBB) poly-house, Faculty of Engineering and Technology, Sam Higginbottom

University of Agriculture, Technology and Sciences, Prayagraj. The plants were planted with a spacing of 60 cm between the rows and 50 cm between the plants. The observations were recorded on randomly selected five plants on characters. The treatment details has been given in Table 1. In general, soil properties of experimental site showed a typical alluvial soil of eastern region of Uttar Pradesh. Soil was sandy loam in texture, slightly acidic in reaction and having low electrical conductivity, very high in organic carbon, low in available nitrogen low in available phosphorus and moderately high in available potassium given in Table 2. Weeding and plant protection measure were followed as and when needed. Observations were recorded at different stages of growth periods. The observations were recorded for characters viz. Plant height (cm), No. of branches/plant, Days to first flowering, Days to 50% flowering, No. of flower per plant, No. of fruit per plant, Fruit weight (g), Fruit weight per plant (g/plant), Fruit

length (cm), Fruit diameter (cm), Fruit yield per plant, fruit yield per hectare (q), Total soluble solid (TSS), Acidity (%), Ascorbic acid (mg/100 g of fruit). The data were statistically analysed by the method suggested by Fisher and Yates, 1936.

3. RESULTS AND DISCUSSION

In the present investigation an attempt has been made to study the effect of different treatment combinations of nano chitosan, bio capsules and NPK@ RDF on growth, yield and quality of chilli in Poly-house condition are discussed. The results obtained are presented in relevant Tables as follows.

3.1 Growth Parameters

Growth parameters comprised of Plant height (30, 60, 90 DAT), number of branches per plant. The data for these characters observed has been listed out in Tables 3 and 4.

Table 1. Details of treatments used

Treatment notation	Treatment combination
T ₀	NPK (RDF)
T ₁	Bio capsule (250ppm)
T ₂	Bio capsule (500ppm)
T ₃	Nano chitosan (60ppm)
T ₄	Nano chitosan (120ppm)
T ₅	NPK + Bio capsule (500ppm)
T ₆	NPK + Nano chitosan (120ppm)
T ₇	NPK + Bio capsule (250ppm) + Nano chitosan (60ppm)
T ₈	NPK + Bio capsule (500ppm) + Nano chitosan (120ppm)

Table 2. Physical and chemical properties of soil at horticulture research field (SHUATS)

Sl. No.	Particulars	Value (0-30 depth)	Method followed
Physical properties (Initial reading)			
1	Sand	58.70	Boyounce Hydrometer (Piper, 1966)
2	Silt	19.46	
3	Clay	21.84	
4	Textural class	Sandy loam	
Chemical properties (Initial reading)			
1	Soil pH	7.3	Potentiometry (Jackson, 1973)
3	Organic carbon (%)	0.56	Walkley and Black's method (Jackson, 1973)
4	Available nitrogen (Kg ha ⁻¹)	49 kg/ha	Alkaline permanganate method (Subbaiah and Asija, 1956)
5	Available phosphorus (Kg ha ⁻¹)	70 kg/ha	Bray's method (Jackson, 1973).
6	Available potash (kg ha ⁻¹)	67 kg/ha	Ammonium acetate method (Jackson, 1973)

Source: Soil analysis was done by KVK (Krishi Vigyan Kendra, Prayagraj, U.P.)

3.1.1 Plant height (cm) [30, 60 and 90 DAT]

The data pertaining to plant height are presented in Table 3. At 30 DAT the maximum height was observed in T₈ (31.20cm) followed by T₇ (28.53cm). The minimum plant height was observed in T₃ (21.40cm). At 60 DAT the maximum height was observed in T₈ (46.87cm) followed by T₇ (44.60cm). The minimum plant height was observed in T₃ (32.80cm). At 90 DAT the maximum height was observed in T₈ (63.73cm) followed by T₇ (60.80cm). The minimum plant height was observed in T₃ (45.40cm). Treatment T₈ (NPK100% + Bio capsule 500ppm + Nano chitosan 120ppm) was significantly superior and recorded maximum plant height (63.73cm). Bio capsules played an important role in the fixation of nutrients (N&P) in plants. The probable reason for increased plant height due to more uptake of applied nutrients by the plants, needed for protein and protoplasm synthesis for higher rate of meiosis, resulting in better photosynthesis and plant growth and ultimately increased the plant height. And Nano chitosan prevent the pathogenic attack. Similar result was given by Khati et al. [3], Chodhary et al. [4] and Hidangmayum et al. [5].

3.1.2 Number of branches/plant

The data pertaining to Number of branches per plant are presented in Table 4. Analysis of variance revealed a significant variation in term of number of branches per plant due to different treatments. At 30 DAT the maximum branches were observed in T₈ (5.47) followed by T₇ (5.20). The minimum branches were observed in T₃ (3.20). At 60 DAT the maximum branches were observed in T₈ (10.53) followed by T₇ (10.07). The minimum branches were observed in T₃ (7.13). At 90 DAT the maximum branches were observed in T₈ (17.60) followed by T₇ (16.80). The minimum branches were observed in T₃ (12.93). Treatment T₈ (N.P.K. 100 % + Bio capsule 500 ppm + Nano chitosan 120 ppm) was significantly superior and recorded maximum no of branches (17.60). Similar result was given by Jayvanth et al., (2018), Almaliotis et al. [6], Bosland et al. [7] and Carrizo et al. [8].

3.2 Flowering Parameter

3.2.1 Days to first flowering, days to 50% flowering and number of flowers per plant

The data on day to First Flowering, days to 50% flowering and number of flowers per plant

were recorded and presented in Table 5. Days to First Flowering varied significantly due to the effect of different treatments. Early Flowering was observed in T₈ (42.17) followed by T₇. The maximum day to First Flowering was observed in T₃ (50.83). Similar result was given by Turan and Sahin [9]. Early flowering (42.17) was recorded for Treatment T₈ (N. P.K. 100 % +Nano chitosan 120 ppm + Bio capsule 500 ppm). Days to 50% flowering varied significantly due to the effect of different treatments. Early Flowering was observed in T₈ (52.92) followed by T₇ (53.67). The maximum day to 50% Flowering was observed in T₃ (62.08). Similar result was given by Turan and Sahin [9,10]. Early flowering (52.92) was recorded for Treatment T₈ (N. P.K. 100 % +Nano chitosan 120 ppm + Bio capsule 500 ppm). Maximum No. of Flowers/Plant was observed in T₈ (271.08) followed by T₇ (268.92). The minimum No. of Flowers/Plant was observed in T₃ (262.58). It is due to the presence of N.P.K, Bio capsule and nano chitosan. Bio capsule have ability to fix the nutrient given by fertilizer in soil and tends towards plant mechanism. N.P.K. had played important role in enhancing the reproductive growth in plant and in other hand chitosan had the ability of cation exchange capacity in soil so that it made the availability of micronutrient like Zn, B, Fe and Cl. Similar result was given by Khan et al. [11], Islam et al. [12] and Kraft et al. [13]. The result was close conformity with Gerjes et al. [14], Orzali et al. [15] and Zayed et al. [16] for number of flowers per plant.

3.3 Yield Parameters

3.3.1 Number of fruits per plant

Results of the present study indicated that with the progression of growth stages there was statistically significant increase in No. of Fruit/Plant (Table 6). Higher No. of Fruit/Plant was observed in T₈ (148.67) followed by T₇ (146.83). The minimum No. of Fruit/Plant was observed in T₃ (138.42). It is evident from the data that treatment T₈ (N. P.K. 100 % +Nano chitosan 120 ppm + Bio capsule 500 ppm) showed best result amongst the other treatment which is at par with each other whereas treatment T₃ (Nano chitosan 60 ppm). This result was close conformity with Gerjes et al. [14], Malerba et al. [17] and Maluin et al. [18].

3.3.2 Fruit weight (g), fruit yield per plant (kg/plant), fruit yield per hectare (q)

Fruit weight of chilli showed differences among the different. Results of the present study indicated statistically significant increase in Fruit weight (g), fruit yield per plant (kg/plant) and fruit yield per hectare (q) (Table 6). Maximum Fruit weight (g) was observed T8 (7.28 g) followed by T7 (6.59g). The minimum Fruit weight (g) was observed in T3 (3.49 g). The maximum fruit weight (7.28g), was recorded for Treatment T8 (N.P.K. 100 % +Nano chitosan 120 ppm + Bio capsule 500 ppm). Maximum Fruit yield (g/plant) was observed in T8 (1082.31g) followed by T7 (967.24g). The minimum Fruit yield (kg/plant) was observed in T3 (483.57g). Higher Fruit yield per hectare (q) was observed in T8 (7.24 q)

followed by T7 (6.48q). The minimum Fruit yield per hectare (q) was observed in T3 (3.27q).The increase in fruit weight may be due to the combination of N.P.K. fertilizer and Bio capsule and Nano chitosan which promoted vegetative growth and development and have associated with acceleration of higher rate of photosynthesis and their accumulation in economic part of plant. Nano chitosan accumulate micronutrient availability to plants and Bio capsule fixed nutrient given through the N.P.K for plants reproductive phase resulting increased fruit weight. This result was close conformity with Sharif et al. [19], Ayoola et al. [20] and Kocięcka et al. [21], and for fruit yield by Mondal et al. [22], Bakshi et al. [23], Kumar et al. [24], Makinde et al. [25].

Table 3. Effect of nano chitosan and bio capsule on different treatment combinations of chilli plant for plant height [30, 60, 90 DAT]

Treatments	30DAT	60DAT	90DAT
T ₀ - N P K (RDF)	25.00	37.33	54.13
T ₁ - Bio capsule (250ppm)	23.05	34.80	47.87
T ₂ - Bio capsule (500ppm)	23.67	35.47	51.20
T ₃ - Nano chitosan (60ppm)	21.40	32.80	45.40
T ₄ - Nano chitosan (120ppm)	22.33	34.07	47.60
T ₅ - NPK + Bio capsule (500ppm)	27.27	41.93	58.67
T ₆ - NPK + Nano chitosan (120ppm)	25.93	39.00	56.00
T ₇ - NPK + Bio capsule (250ppm) + Nano chitosan (60ppm)	28.53	44.60	60.80
T ₈ - NPK + Bio capsule (500ppm) + Nano chitosan (120ppm)	31.20	46.87	63.73
F-TEST	S	S	S
SE(d)	2.03	1.75	1.74
C.V.	9.79	5.57	3.95
C.D. AT 5%	4.30	3.72	3.69

Table 4. Effect of nano chitosan and bio capsule on different treatment combinations of chilli plant for number of branches per plant [30, 60, 90 DAT]

Treatments	30DAT	60DAT	90DAT
T ₀ - N P K (RDF)	4.27	9.13	14.80
T ₁ - Bio capsule (250ppm)	3.60	7.67	13.73
T ₂ - Bio capsule (500ppm)	4.07	8.20	14.47
T ₃ - Nano chitosan (60ppm)	3.20	7.13	12.93
T ₄ - Nano chitosan (120ppm)	3.33	7.40	13.20
T ₅ - NPK + Bio capsule (500ppm)	4.93	9.80	16.47
T ₆ - NPK + Nano chitosan (120ppm)	4.53	9.40	16.00
T ₇ - NPK + Bio capsule (250ppm) + Nano chitosan (60ppm)	5.20	10.07	16.80
T ₈ - NPK + Bio capsule (500ppm) + Nano chitosan (120ppm)	5.47	10.53	17.60
F-TEST	S	S	S
SE(d)	0.24	0.41	0.35
C.V.	6.99	5.64	2.87
C.D. AT 5%	0.52	0.86	0.75

Table 5. Effect of nano chitosan and bio capsule on different treatment combinations of chilli plant for days to first flowering, days to 50% flowering and number of flowers per plant

Treatments	Days to first flower	Days to 50% flower	No. Of flowers/plant
T ₀ - N P K (RDF)	46.00	56.50	265.58
T ₁ - Bio capsule (250ppm)	48.17	59.50	264.00
T ₂ - Bio capsule (500 ppm)	47.33	58.00	264.50
T ₃ - Nano chitosan (60 ppm)	50.83	62.08	262.58
T ₄ - Nano chitosan (120 ppm)	49.08	61.00	263.42
T ₅ - NPK + Bio capsule (500 ppm)	43.58	54.17	267.83
T ₆ - NPK + Nano chitosan (120 ppm)	44.67	55.50	267.42
T ₇ - NPK + Bio capsule (250 ppm) + Nano chitosan (60 ppm)	43.00	53.67	268.92
T ₈ - NPK + Bio capsule (500 ppm) + Nano chitosan (120 ppm)	42.17	52.92	271.08
F-TEST	S	S	S
SE(d)	1.07	1.09	2.00
C.V.	2.85	2.35	0.92
C.D. AT 5%	2.28	2.32	4.24

Table 6. Effect of nano chitosan and bio capsule on different treatment combinations of chilli plant for fruit weight, fruit yield per plant, fruit yield per hectare

Treatments	No. of fruits/plant	Fruit weight (g)	Fruit weight/plant (kg/plant)	Fruit yield quintal/200m ²
T ₀ - N P K (RDF)	142.58	4.70	670.11	4.50
T ₁ - Bio capsule (250ppm)	139.58	3.95	550.65	3.70
T ₂ - Bio capsule (500ppm)	141.67	4.28	606.34	4.07
T ₃ - Nano chitosan (60ppm)	138.42	3.49	483.57	3.27
T ₄ - Nano chitosan (120ppm)	139.08	3.67	510.99	3.44
T ₅ - NPK + Bio capsule (500ppm)	146.00	5.91	862.18	5.78
T ₆ - NPK + Nano chitosan (120ppm)	144.08	5.29	762.73	5.11
T ₇ - NPK + Bio capsule (250ppm) + Nano chitosan (60ppm)	146.83	6.59	967.24	6.48
T ₈ - NPK + Bio capsule (500ppm) + Nano chitosan (120ppm)	148.67	7.28	1082.31	7.24
F-TEST	S	S	S	S
SE(d)	1.73	0.09	14.30	0.10
C.V.	1.48	2.14	2.43	2.49
C.D. AT 5%	3.67	0.18	30.31	0.21

Table 7. Effect of nano chitosan and bio capsule on different treatment combinations of chilli plant for fruit length and diameter (cm)

Treatments	Fruit length (cm)	Fruit diameter (cm)
T ₀ - N P K (RDF)	7.57	0.93
T ₁ - Bio capsule (250ppm)	6.87	0.84
T ₂ - Bio capsule (500ppm)	7.20	0.92
T ₃ - Nano chitosan (60ppm)	6.27	0.81
T ₄ - Nano chitosan (120ppm)	6.43	0.84
T ₅ - NPK + Bio capsule (500ppm)	8.80	1.06
T ₆ - NPK + Nano chitosan (120ppm)	8.03	0.99
T ₇ - NPK + Bio capsule (250ppm) + Nano chitosan (60ppm)	9.27	1.13
T ₈ - NPK + Bio capsule (500ppm) + Nano chitosan (120ppm)	10.13	1.21
F-TEST	S	S
SE(d)	0.32	0.04
C.V.	5.01	5.55
C.D. AT 5%	0.68	0.09

Table 8. Effect of Nano chitosan and bio capsule on different treatment combinations of chilli plant for TSS, acidity and Vitamin C content

Treatments	T.S.S. (°Brix)	Acidity %	Vitamin C mg/100g
T ₀ - N P K (RDF)	8.57	0.219	131.41
T ₁ - Bio capsule (250ppm)	7.92	0.234	125.74
T ₂ - Bio capsule (500ppm)	8.39	0.227	127.05
T ₃ - Nano chitosan (60ppm)	6.97	0.252	117.10
T ₄ - Nano chitosan (120ppm)	7.26	0.247	124.82
T ₅ - NPK + Bio capsule (500ppm)	9.21	0.196	136.87
T ₆ - NPK + Nano chitosan (120ppm)	8.85	0.216	134.01
T ₇ - NPK + Bio capsule (250ppm) + Nano chitosan (60ppm)	9.46	0.182	140.02
T ₈ - NPK + Bio capsule (500ppm) + Nano chitosan (120ppm)	9.67	0.172	143.96
F-TEST	S	S	S
SE(d)	0.22	0.01	3.98
C.V.	3.18	7.98	3.71
C.D. AT 5%	0.47	0.03	8.43

3.3.3 Fruit length (cm) and fruit diameter (cm)

Results of the present study indicated that with the progression of growth stages there was statistically significant increase in fruit length (cm) and fruit diameter (cm) (Table 7). Maximum fruit length (cm) was observed in T₈ (10.13cm) followed by T₇ (9.27cm). The minimum fruit length (cm) was observed in T₃ (6.27cm). Maximum fruit Diameter (cm) was observed in T₈ (1.21cm) followed by T₇ (1.13cm), T₅ (1.06cm), T₆ (0.99cm), T₀ (0.93cm), T₂ (0.92cm), T₁ (0.84cm), T₄ (0.84cm) and T₃. The minimum fruit Diameter (cm) was observed in T₃ (0.81cm). It is confirmed from the data that treatment T₈ (N. P.K. 100 % +Nano chitosan 120 ppm + Bio capsule 500 ppm) showed best result amongst the other treatment which is at par with each other whereas treatment T₄ (Nano chitosan 60 ppm). Similar findings were observed by Rafique et al. [26] and Singh et al. [27] for fruit length and Duhana et al. [28], Khalifa et al. [29] and Malerba et al. [17] for fruit diameter.

3.4 Quality Parameter

3.4.1 TSS (°Brix), acidity, vitamin C content (mg/100 g)

The total Soluble Solid, acidity, Vitamin C content of Chilli showed significant variation among different treatment which is presented in Table 8. Maximum value of total Soluble Solid was observed in T₈ (9.67°B) followed by T₇ (9.46°B). The minimum total Soluble Solid was observed in T₃ (6.97°B). The highest TSS (9.67°B) value was observed for T₈ (N.P.K. 100 % +Nano

chitosan 120 ppm + Bio capsule 500 ppm). Based on the data it is found that in nine treatment combination, T₈ (NPK % + Bio capsule 500ppm + Nano chitosan 120ppm) recorded minimum (0.172%) for acidity followed by T₇ (NPK % + Bio capsule 250ppm + Nano chitosan 60ppm) with (0.182%). The maximum vitamin C content was observed in T₈ (NPK % + Bio capsule 500ppm + Nano chitosan 120ppm) (143.96 mg) followed by T₇ (NPK % + Bio capsule 250ppm + Nano chitosan 60ppm) with (140.02 mg). The minimum Vitamin C was observed in T₃ (Nano chitosan 60ppm) with (117.10 mg). When the nutrient supply became insufficient, the limited synthesized carbohydrates meet the requirements of only vegetative parts. Contrary to this, combination of N.P.K. with Bio capsule supplied adequate level of nutrients because of Bio capsule prevent the leaching of nutrient and fixed the nutrient for plant growth. Thus, synthesized carbohydrates translocated to the fruits, which ultimately increased the total soluble solids of fruit. Similar finding was given by Duhana et al. [28], Chiou et al. [30] and Gadalla et al. [31] for TSS, acidity and Vitamin C content.

4. CONCLUSION

On the basis of the present investigation, it is concluded that the various treatments applied to enhance the vegetative and growth of chilli treatment T₈ (NPK (RDF) + Nano chitosan 120 ppm + Bio capsule 500ppm) was found to be superior among others, followed by T₇ (NPK (RDF) + Nano chitosan 60ppm + Bio capsule 250ppm) and the lowest was T₃ (Nano chitosan

60 ppm) in every aspect of growth, yield, quality. Therefore, use of Nano chitosan @ 120 ppm along with 500 ppm Bio capsule could be used for getting better yield in chilli. Moreover, application of Nano chitosan and bio capsule on nutritional quality of chilli could be studied in details further.

ACKNOWLEDGEMENT

The author would like express heartfelt gratitude to Dr. Samir E. Topno (Assistant Professor, Department of Horticulture, NAI, SHUATS, Prayagraj) for his continuous guidance and supervision throughout our research period.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Pickersgill BARBARA. Cytogenetics and evolution of capsicum L. Chromosome Engineering in Plants: Genetics, Breeding, Evolution, Part B. Elsevier, Amsterdam. 1991;139-160.
- Pickersgill B. Relationships between weedy and cultivated forms in some species of chilli peppers (*Genus Capsicum*). Evolution. 1971;25(4):683-691.
- Khati P, Chaudhary P, Gangola S, Bhatt P, Sharma A. Nano chitosan supports growth of *Zea mays* and also maintains soil health following growth. 3 Biotech. 2017;7(1):1-9.
- Choudhary RC, Kumari S, Kumaraswamy RV, Sharma G, Kumar A, Budhwar S, Saharan V. Chitosan nanomaterials for smart delivery of bioactive compounds in agriculture. In Nanoscale Engineering in Agricultural Management. CRC Press. 2019;124-139.
- Hidangmayum A, Dwivedi P, Katiyar D, Hemantaranjan A. Application of chitosan on plant responses with special reference to abiotic stress. Physiology and Molecular Biology of Plants. 2019;25(2):313-326.
- Almaliotis D, Bladenopoulou S, Chatzissavvidis CH, Karagiannidis N. Broccoli (*Brassica oleracea* L. Var. *Italica*) yield as affected by soil fertility parameters in northern Greece. Acta Hort. 2007;729: 409-413.
- Bosland PW. Capsicums: Innovative uses of an ancient crop. Progress in new crops. ASHS Press, Arlington, VA. 1996;479-487.
- Carrizo García C, Barfuss MH, Sehr EM, Barboza GE, Samuel R, Moscone EA, Ehrendorfer F. Phylogenetic relationships, diversification and expansion of chilli peppers (*Capsicum*, *Solanaceae*). Annals of Botany. 2016;118(1):35-51.
- Turan M, Gulluce M, Cakmakci R, Oztas T, Sahin F, Gilkes RJ, Prakongkep N. The effect of PGPR strain on wheat yield and quality parameters. In proceedings of the 19th world congress of soil science: Soil solutions for a changing world. Brisbane, Australia. 2010;209:212.
- Baris O, Sahin F, Turan M, Orhan F, Gulluce M. Use of plant-growth-promoting rhizobacteria (PGPR) seed inoculation as alternative fertilizer inputs in wheat and barley production. Communications in Soil Science and Plant Analysis. 2014;45(18): 2457-2467.
- Khan, Insaf, Singh, Devendra, Jat Bhanwar Lal. Effects of biofertilizers on plant growth and yield characters of *Pisum sativum* L. Adv. Res. J. Crop Improv. 2017; 8(1):99-108.
- Islam S, Akanda AM, Prova A, Sultana F, Hossain MM. Isolation and identification of plant growth promoting rhizobacteria from cucumber rhizosphere and their effect on plant growth promotion and disease suppression. Front. Microbiol. 2015;6: 1360. DOI:10.3389/fmicb.2015.0136
- Kraft KH, Brown CH, Nabhan GP, Luedeling E, Luna Ruiz JDJ, Coppens d'Eeckenbrugge G, Gepts P. Multiple lines of evidence for the origin of domesticated chilli pepper, *Capsicum annuum*, in Mexico. Proceedings of the National Academy of Sciences. 2014;111(17): 6165-6170.
- Geries L, Omnia HS, Marey RA. Soaking and foliar application with chitosan and nano chitosan to enhancing growth, productivity and quality of onion crop. Plant Arch. 2020;20(2):3584-3591.
- Orzali L, Corsi B, Forni C, Riccioni L. Chitosan in agriculture: A new challenge for managing plant disease. Biological Activities and Application of Marine Polysaccharides. 2017;87-96.
- Zayed MM, Elkafafi SH, Zedan AM, Dawoud SF. Effect of nano chitosan on growth, physiological and biochemical parameters of *Phaseolus vulgaris* under salt stress. Journal of Plant Production. 2017;8(5):577-585.

17. Malerba M, Cerana R. Chitosan effects on plant systems. *International Journal of Molecular Sciences*. 2016;17(7):996.
18. Maluin FN, Hussein MZ. Chitosan-based agronanochemicals as a sustainable alternative in crop protection. *Molecules*. 2020;25(7):1611.
19. Sharif R, Mujtaba M, Ur Rahman M, Shalmani A, Ahmad H, Anwar T, Wang X. The multifunctional role of chitosan in horticultural crops; A review. *Molecules*. 2018;23(4):872.
20. Ayoola OT, Adeniyen ON. Influence of poultry manure and NPK fertilizer on yield and yield components of crops under different cropping systems in south west Nigeria. *African Journal of Biotechnology*. 2006;5(15).
21. Kocięcka J, Liberacki D. The potential of using chitosan on cereal crops in the face of climate change. *Plants*. 2021;10(6):1160.
22. Mondal MM, Malek MA, Puteh AB, Ismail MR, Ashrafuzzaman M, Naher L. Effect of foliar application of chitosan on growth and yield in okra. *Australian Journal of Crop Science*. 2012;6(5):918-921.
23. Bakshi M, Wali VK, Bakshi P, Sharma A, Sharma D, Shah RA. Integrated nutrient management induced changes in nutrient uptake, fruit yield and quality of Kinnow mandarin. *Indian Journal of Agricultural Sciences*. 2017;87(3):414-8.
24. Kumar UJ, Bahadur V, Prasad VM, Mishra S, Shukla PK. Effect of different concentrations of iron oxide and zinc oxide nanoparticles on growth and yield of strawberry (*Fragaria x ananassa* Duch) cv. Chandler. *International Journal of Current Microbiology and Applied Sciences*. 2017;6(8):2440-2445.
25. Makinde EA, Ayoola OT, Akande MO. Effects of organo-mineral fertilizer application on the growth and yield of egusi melon. *Australian Journal of Basic and applied sciences*. 2007;1(1):15-19.
26. Munazza Rafique, Aneela Riaz, Ashfaq Anjum, Amjad Qureshi M, Fakhar Mujeeb. Role of bioinoculants for improving growth and yield of okra (*Abelmoschus esculentus*). *Universal Journal of Agricultural Research*. 2018;6(3):105-112.
27. Singh A, Singh JN. Effect of biofertilizers and bioregulators on growth, yield and nutrient status of strawberry cv. Sweet Charlie. *Indian Journal of Horticulture*. 2009;66(2):220-224.
28. Duhan JS, Kumar R, Kumar N, Kaur P, Nehra K, Duhan S. Nanotechnology: The new perspective in precision agriculture. *Biotechnol Rep*. 2017;15:11–23.
29. Khalifa NS, Hasaneen MN. The effect of chitosan–PMAA–NPK nanofertilizer on *Pisum sativum* plants. *3 Biotech*. 2018;8(4):1-12.
30. Chiou KL, Hastorf CA. A systematic approach to species–level identification of chile pepper (*Capsicum spp.*) seeds: Establishing the groundwork for tracking the domestication and movement of chile peppers through the Americas and beyond. *Economic Botany*. 2014;68(3): 316-336.
31. Gadalla MB. Knowledge, attitude and practice about health benefits of chilli pepper among students of university of science and technology; 2020.

© 2022 Devi and Topno; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92851>