



Application of Multivariate Principal Component Analysis for Characterization of Leaf Litters in Northern Nigeria

Folasade Aanuoluwapo Akinsola ^{a*}, Ishaku Yo'ila Amapu ^a,
Eunice Yemisi Oyinlola ^a
and Tawakalitu Akanji Abdulsalam ^b

^a Department of Soil Science, Faculty Agriculture, Ahmadu Bello University, Zaria, Nigeria.
^b Vetplace Animal hospital, Yaba, Lagos, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author FAA conceptualize the study and wrote the first draft of the manuscript. Authors IYA and EYO wrote the protocol and manage the analyses of the study. Author TAA performed the statistical analysis. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ASRJ/2024/v8i1141

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106715>

Original Research Article

Received: 08/08/2023
Accepted: 12/10/2023
Published: 02/01/2024

ABSTRACT

This study was to objectively describe the interrelationship between mass loss and leaf chemical parameters of litter species: *Khaya senegalensis*, *Mangifera indica*, *Gmelina arborea*, and *Eucalyptus camaldulensis* using principal component analysis. Leaf litters were analyzed for chemical compositions. Mass loss and ten litter chemical parameters such as organic carbon, total nitrogen, total phosphorus, potassium, magnesium, calcium, sodium, total soluble polyphenol, total sulphur and carbon to nitrogen ratio were investigated. Principal Component Analysis (PCA) was used to identify the variation of litter chemical properties. The results showed significant

*Corresponding author: E-mail: olomolafolashade@gmail.com;

relationships between mass loss and litter chemical parameters in PC₁ and PC₂ axes. *Khaya senegalensis* has the highest loadings 84% followed by *Gmelina arborea* 77%, *Eucalyptus camaldulensis* 75.3% and *Mangifera indica* 64.5%. In conclusion, *Khaya senegalensis* showed a propensity to have a faster response in driving biogeochemical cycling during decomposition.

Keywords: Leaf litter; chemical composition; mass loss; decomposition.

1. INTRODUCTION

Litter mass loss or decay is the sum of carbon dioxide, CO₂, release and discharge of compounds, which contains both carbon compounds and nutrients [1]. According to [2], the breakdown of leaf litter is an important part of the global biogeochemical cycles that affect soil carbon storage, nutrient availability, and plant production. Designing agroforestry systems for soil conservation and improvement requires an evaluation of the quality of the litter and the rate of leaf biomass breakdown [3,4]. The management of chemical element and compound levels in agroforestry systems is once again emphasized [5,2]. Many studies concluded that the rate of leaf litter decomposition depended on tree species, the chemical composition of the leaves, and environmental factors such as temperature and soil moisture [6,7].

A multivariate technique called principal component analysis (PCA) can be used to reduce the dimensionality of such datasets, improving their interpretability while minimizing information loss [8,9]. It accomplishes this by producing fresh, uncorrelated variables that maximize variance one after the other. As a result, PCA is an adaptive data analysis technique [10]. Finding such new variables, the principal components, reduces to solving an eigenvalue/eigenvector issue, and the new variables are specified by the dataset at hand, not a priori. Many researchers have used the independent factor scores derived from multivariate technique of principal component factor analysis to soil and leaf litter properties [3,11,12]. The modelling component of leaf litter decomposition is widely reported in scientific literatures [13,14]. In Sub-Saharan Africa, there is dearth of information on the interrelationships among mass loss and litter chemistry traits of leaf litters using a multivariate approach. Understanding leaf litter decomposition dynamics is imperative owing to nutrient release rates and synchronization. The overarching aim is to use principal components analysis to account for interrelationship between mass loss and

chemical parameters of leaf litters in Northern Nigeria.

2. MATERIALS AND METHODS

2.1 Experimental Site and Climatic Conditions

The study was carried out at Institute for Agricultural Research, Ahmadu Bello University, Zaria, Kaduna State, Northern Guinea Savannah zone of Nigeria field (IAR plot R14). The experimental area in Samaru has a geo reference: 10°10' 0" N and 70°37' 60"E and 688 m a.s.l. [15]. This region is characterized by two distinct seasons: the dry season comprising the cold dry period also known as the harmattan, November-December and the hot dry period between April-June as well as a warm rainy season between July-September with rainfall of about 1060mm annually. October and March constitute traditional months between rainy and cold dry season and between hot dry period and rainy season. Warm conditions and high relative humidity prevail during rainy season. The region is characterized by a lot of leaf falls during the dry season and very high temperatures which affect plant growth and developmental process [16].

2.2 Collection and Processing of Leaf Samples

Fall leaves of African mahogany, *Khaya senegalensis*; mango, *Mangifera indica*; Beech wood, *Gmelina arborea*; and river red gum, *Eucalyptus camaldulensis*, were picked from the selected tree species. The collected leaf litter samples were cleaned and all sediments and dirt particles were removed by using a soft brush with running tap water followed by final rinsing in distilled water. Each sample was air-dried under shade at the Department of Soil Science Laboratory.

2.3 Chemical Analysis of Leaf Litter

Air dried leaf litter samples were ground in mortar and sieved through a 1mm mesh size sieve. The

fine powder was used for the estimation of C, N, P, K, Ca, Mg, lignin and total soluble polyphenols. The standard procedures that were adopted for the chemical analysis are presented below.

2.4 Total Carbon

The total carbon content was determined by igniting the samples at 550°C using the Walkley Black method [17].

2.5 Total Nitrogen

Nitrogen content in fresh leaf litter was determined by digesting 0.1 g of samples in 5 ml of concentrated sulphuric acid using digestion mixture (sodium sulphate: copper sulphate in 10:4 ratio and nitrogen in the digest was determined by Kjeldhal's method as reviewed by Saez-Plaza et al. [18].

2.6 Total Phosphorus

Approximately 0.2g of the powdered leaf sample was digested in tri-acid mixture (nitric acid: perchloric acid: sulphuric acid in 1:1:3 ratio) and the digest was made up to 100 ml. A known quantity of aliquot was taken to determine the phosphorus content by following chlorostannous reduced molybdophosphoric blue colour method in sulphuric acid system [19] and the colour intensity was read at 660 nm in UV spectrophotometer.

2.7 Total Basic Cations

Calcium (Ca), magnesium (Mg), potassium (K), and sodium (Na) were determined using flame photometry and Atomic Absorption Spectrophotometer (AAS) as appropriate after wet digestion [20].

2.8 Sulphur

This was determined turbidimetrically using spectrophotometer to read for absorbance at wavelength of 430nm. [21]

2.9 Total Soluble Polyphenols

Total soluble polyphenol was determined in the Pharmacognosy and Drug Development Laboratory of the Department of

Pharmacognosy, Ahmadu Bello University, Zaria using the Follin-Ciocalteu's method [22].

2.10 Data Analysis

Principal component analysis of morphometric measures and their contribution to mass loss was analyzed using the factor procedure of SAS (2014). The correlation coefficient of mass loss and litter chemical parameters was determined. From the correlation matrix, data for the principal component factor analysis was generated. Principal component analysis [23] is a method of transforming the variables in a multivariate data set X_1, X_2, \dots, X_P into variables Y_1, Y_2, \dots, Y_P which are uncorrelated with each other and account for the decreasing proportion of the total variance of the original variable.

3. RESULTS AND DISCUSSION

Fig. 1 shows the relationship between mass loss and chemical composition in litter of *Khaya senegalensis*. Mass loss had high negative and significant $p < 0.05$, association with potassium, organic carbon and phosphorus. Positive and significant association was recorded among sulphur, nitrogen, lignin and total phenol content. PC1 had the highest loadings with eigen value of 9.79 and percent variance of 75.3%. As seen in Fig. 2 *Gmelina Arborea*, Organic carbon, sodium, total phenol content, calcium and nitrogen had positive loadings with 84% total variance on the PC1 while magnesium, sulphur, nitrogen: phosphorus ratio, phosphorus, mass loss and potassium had negative loadings. Principal component analysis of mass loss and chemical composition of *Khaya senegalensis* leaf litter is shown in Fig. 3. Mass loss had positive relationship with calcium, sodium, magnesium, nitrogen: phosphorus ratio, organic carbon, carbon to nitrogen ratio. The relationship between lignin and total phosphorus and potassium and total phenol count was negative. PC1 had a percent variance of 64.5% while PC2 recorded 35.5% of the total variations.

Principal component analysis of mass loss and chemical composition of *Mangifera indica* are shown in Fig. 4. Sodium had positive relationship with calcium, sodium, nitrogen: phosphorus ratio, organic carbon, carbon to nitrogen ratio, lignin and total nitrogen. The relationship between mass loss, magnesium and total phosphorus was negative.




Fig. 1. Principal component matrix of mass loss and chemical parameters in *Eucalyptus Camaldulensis*

TN = total nitrogen, TK = total potassium, OC = organic carbon, TP = total phosphorus, TPC = total phenol content, C: N = carbon to nitrogen ratio, N: P = Nitrogen to phosphorus ratio, Mg = magnesium, Ca = Calcium, Na = sodium, S = Sulphur




Fig. 2. Principal component matrix of mass loss and chemical parameters in *Gmelina arborea*

TN = total nitrogen, TK = total potassium, OC = organic carbon, TP = total phosphorus, TPC = total phenol content, C: N = carbon to nitrogen ratio, N: P = Nitrogen to phosphorus ratio, Mg = magnesium, Ca = Calcium, Na = sodium, S = Sulphur




Fig. 3. Principal component matrix of mass loss and chemical parameters in *Khaya senegalensis*

TN = total nitrogen, TK = total potassium, OC = organic carbon, TP = total phosphorus, TPC = total phenol content, C: N = carbon to nitrogen ratio, N: P = Nitrogen to phosphorus ratio, Mg = magnesium, Ca = Calcium, Na = sodium, S = Sulphur




Fig. 4. Principal component matrix of mass loss and chemical parameters in *Mangifera indica*
 TN = total nitrogen, TK = total potassium, OC = organic carbon, TP = total phosphorus, TPC = total phenol content, C: N = carbon to nitrogen ratio, N: P = Nitrogen to phosphorus ratio, Mg = magnesium, Ca = Calcium, Na = sodium, S = Sulphur

In *Mangifera indica*, mass loss was strongly and negatively correlated with magnesium and total phosphorus which agrees with the observations of several studies [24,25]. The positive relationship between carbon: nitrogen ratio and nitrogen: potassium ratio was based on the premise that nitrogen concentration limits the activity of decomposers [26]. In *Gmelina arborea*, increase in mass loss will significantly cause a reduction in organic carbon, calcium, sodium, lignin, total phenol content and C: N this agrees with trend reported by Liu et al. [27]. who observed that mass-loss rates and late-stage litter negatively correlated to carbon, calcium and sodium. In *Eucalyptus camaldulensis*, mass loss was strongly, negatively and highly correlated with potassium, sulphur, total phenol content and C: N ratio. Many studies showed that litter mass loss was strongly correlated with nitrogen, potassium, sulphur and C: N ratio [13,25]. The concept of the regulatory effect of C: N ratio was based on the fact that nitrogen concentration limits the activity of decomposers [26,28]. Judging from the positive correlations between mass loss and potassium, magnesium, sulphur, nitrogen and C: N ratio, this implies good predictability of mass loss of litters of *Eucalyptus camaldulensis*.

4. CONCLUSIONS

Multivariate PCA showed noticeable variations among leaf litters parameters. Since these methods are of high accuracy and have different abilities, they could be used for indicator of soil health through litter decomposition. The main component factor analysis led to an objective

simultaneous analysis of organic carbon, total nitrogen, total phosphorus, potassium, magnesium, calcium, sodium, total soluble polyphenol, total sulphur and carbon to nitrogen ratio in different leaf litters rather than on individual basis. This resulted in the reduction of the leaf litters chemical parameters to two PCs in each leaf litter species.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Brady N, Weil R. Nutrient cycles and soil fertility. In: Anthony, V.R., Ed., elements of the nature and properties of soils, 3rd Edition, Pearson Education Inc, Upper Saddle River, NJ. 2010;396-420.
2. Tenkiano N, Sia D, Chauvet É. Leaf litter decomposition in Guinean savannah streams. *Inland Waters*. 2018;8(4):413-421.
3. Qu M, Wang Y, Huang B, Zhao Y. Source apportionment of soil heavy metals using robust absolute principal component scores-robust geographically weighted regression (RAPCS-RGWR) receptor model. *Science of the Total Environment*. 2018;626:203-210.
4. Simpson LT, Chapman SK, Simpson LM, Cherry JA. Do global change variables alter mangrove decomposition? A systematic review. *Global Ecology and Biogeography*; 2023.

5. Xiao W, Han YH, Chen Kumar Chen C, Guan Q. Multiple interactions between tree composition and diversity and microbial diversity under litter decomposition. *Geoderma*. 2019;341:161-171.
6. Asigbaase M, Dawoe E, Lomax BH, Sjogersten S. Temporal changes in litterfall and potential nutrient return in cocoa agroforestry systems under organic and conventional management, Ghana. *Heliyon*. 2021;7(10).
7. Tu L, Hu H, Hu T, Jian Z, Xianwei L, Li L, et al.. Litterfall, litter decomposition, and nutrient dynamics in two subtropical bamboo plantations of China. *Pedosphere*. 2014;24(01):84–97.
8. Silva LC, Lopes B, Pontes MJ, Blanquet I, Segatto ME, Marques C. Fast decision-making tool for monitoring recirculation aquaculture systems based on a multivariate statistical analysis. *Aquaculture*. 2021;530:735931.
9. Jolliffe IT. *Principal component analysis*, 2nd edn. New York, NY: Springer-Verlag; 2002.
10. Xie P, Ma Z, Du R, Lv M, Shen Y, Lu X, Cen H. Generating high-quality 3DMPCs by adaptive data acquisition and NeREF-based reflectance correction to facilitate efficient plant phenotyping; 2023. Available:arXiv preprint arXiv:2305.06777.
11. Liu Z, Jia M, Li Q, Lu S, Zhou D, Feng L, Yu J. Comparative analysis of the properties of biochars produced from different pecan feedstocks and pyrolysis temperatures. *Industrial Crops and Products*. 2023;197:116638.
12. Kooch Y, Ghorbanzadeh N, Wirth S, Novara A, Shah Piri A. Soil functional indicators in a mountain forest-rangeland mosaic of northern Iran. *Ecological Indicators*. 2021;26:107672.
13. Berg B, McClaugherty C. *Plant Litter. Decomposition, Humus Formation, Carbon Sequestration*, third ed. Springer Verlag, Heidelberg, Berlin. 2014;317.
14. Sánchez-López N, Hudak AT, Boschetti L, Silva CA, Robertson K, Loudermilk EL, Taylor MK. A spatially explicit model of tree leaf litter accumulation in fire maintained longleaf pine forests of the southeastern US. *Ecological Modelling*. 2023;481:110369.
15. Goggle Earth. *Goggle Earth Imagery*; 2023.
16. Institute for Agricultural Research Metrological station, IARMS. Metrological data from IAR metrological station, Ahmadu Bello University, Samaru, Zaria Nigeria; 2021.
17. FAO. Standard operating procedure for soil organic carbon. Walkey-Black method Titration and colorimetric method; 2019. Available:Fao.org
18. Saez-Plaza, Michalowski T, Navas MJ, Asuero AG. An overview of the Kjeldahl method of nitrogen determination part 1. Early history, chemistry of the procedure, and titrimetric finish. *Critical Review in Analytical Chemistry*. 2013;4:43-45.
19. Bray RH, Kurtz,L. Determination of total, organic, and available forms of phosphorus in soils. *Soil Science*. 1945; 59(1):39-46.
20. Anderson C, Peterson M, Curtin D. Base cations, KC and Ca2C, have contrasting effects on soil carbon, nitrogen and denitrification dynamics as pH rises. *Soil Biology and Biochemistry*. 2017;3: 99-107.
21. Tabatabai MA, Bremmer JM. A simple turbidimetric method of determining total sulphur in plant materials. *Agronomy Journal*. 1970;62(6):805-806.
22. Moyer RA, Hummer KE, Finn CE, Frei B, Wrolstad RE. Anthocyanins, phenolics, and antioxidant capacity in diverse small fruits: Vaccinium, Rubus, and Ribes. *Journal of Agriculture and Food Chemistry*. 2002;50(3):519–525.
23. Everitt BS, Landau S, Leese M. *Cluster Analysis*. 4th edition. Arnold Publisher, London; 2001.
24. Jiang L, Wang H, Li S, Fu X, Dai X, Yan H, Kou L. Mycorrhizal and environmental controls over root trait–decomposition linkage of woody trees. *New Phytologist*. 2021;229(1):284-295.
25. Liu P, Huang J, Sun OJ, Han X. Litter decomposition and nutrient release as affected by soil nitrogen availability and litter quality in a semiarid grassland ecosystem. *Oecologia*. 2010; 162:771-780.
26. Ma Y, Cai R, Zhong H, Wu L, Ge G. The home-field advantage of litter decomposition in lake wetlands and the community characteristics of bacterial and eukaryotic decomposers. *Plant and Soil*. 2023;483(1-2):109-130.
27. Liu S, Yang R, Hou C. Effect of enzyme activity changes on decomposition characteristics of leaf litter mixed decomposition of configured tree species

- in Ecological Tea Garden. Agriculture. 2023;13(2):394.
28. Parthey ST, Thavathasan NV. Agronomic potentials of rarely used agroforestry species for smallholder agriculture in sub-Saharan Africa: An exploratory study. Communication in soil science and plant analysis. 2013;44(11):1733-1746.

© 2024 Akinsola et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/106715>