



Carbon Sequestration Potential of Roadside Trees in Southern Punjab, Pakistan

Muhammad Zubair ^{a*}, Sidra Khan ^a and Syed Bilal Hussain ^b

^a Department of Forestry and Range Management, Bahauddin Zakariya University, Multan, Pakistan.

^b Institute of Molecular Biology and Biotechnology, Bahauddin Zakariya University, Multan, Pakistan.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRAF/2022/v8i4174

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92869>

Original Research Article

Received 15 August 2022

Accepted 21 October 2022

Published 25 October 2022

ABSTRACT

Trees along roadside play an important role in climate change mitigation through carbon sequestration. Vehicular pollution is not only responsible for environmental degradation but also cause various health issues to inhabitants resides in the vicinity of such roads. Present study was conducted to observe the role of trees grown along roadside and their underneath soil are efficient in carbon storage at four different sites of Multan city, Punjab, Pakistan. Four sites covering whole Multan city were chosen namely Khaniwal road (S1); Nangshah road (S2); Shujaabad road (S3). and Boson road (S4). The most abundant of the tree species viz-a-viz, *Dalbergia sissoo*, *Eucalyptus camadulensis* and *Vachellia nilotica* were sampled having 10 trees across 04 sites and measured for their diameter and height, whereas, underneath each tree soil samples were extracted at two depths viz., (0-20) (20-40) were recorded during the field visit. The study calculated biomass using allometric equations while soil organic carbon and organic carbon was assessed using Walkely Black method. Although species have different height and diameter, hence, their sequestration rate was also different. Data revealed that in all four sites the carbon sequestration rate remained higher in *E. camaldulensis* for biomass (Above ground, Belowground and total biomass), carbon and soil organic carbon estimation. as compared to *D.sissoo* and *V. nilotica* at all four sites. This research concluded that *E. camaldulensis* species may be be planted along the roadside due to its higher carbon sequestration rate in maintaining health roadside ecosystem.

Keywords: Climate change; mitigation; transport; biomass; vegetation.

1. INTRODUCTION

Vegetation growing along roadside perform an ecological function not only on reducing pollution load but also sequester carbon and help mitigate climate change [1]. Depending upon the types of species and their composition different land uses differ in their carbon storage potential as well as their ecological functions [2]. Roadside plantation can be potential contributor in decreasing the amount of CO₂ in atmosphere by collecting in the form of biomass. The importance of plantation in cities along the roadside should not be neglected as it improves the pollution caused by automobile. The roadside plantation additionally has friendly and attractive means to inhabitants and great ecological value to cities as portion of green set-up. However, in urban areas plantation are affected by anthropogenic activities due to which degradation of urban forest and decrease the life span of trees as compared to trees in rural areas occurred. The main stressor of urban trees is air pollution, it has negative impact on tree health and reduce the life of tree. Removal of pollution through plants can be undying, although plants naturally work as a provisional preservation place for dust particles [3]. For appropriate repair of roadsides, there is need to know the ecological relationship of roadside plantation and their part in carbon storage plantation of an area [4]. Depend on the types of species and their structure different land uses change in carbon storage potential as well as their ecological functions [2]. Therefore, maintains the roads and their construction are very important perception of the economy, government, and country's development [5].

Now a days climate change is the main issue all around the world having great importance, caused by excessive energy use and anthropogenic activities [6,7]. Carbon sequestration is extremely important as climate change and global warming up of the earth surface. Biomass of plant indicate the sequestration rate of carbon in plants. In plant the density of wood plays a major role of carbon sequestration. By adding the ornamental beauty of urban life plants play an important role in cooling effect as transpiration rate and through photosynthesis carbon dioxide sequestration and release of oxygen in the atmosphere [8]. Carbon sequestration by plantations is a low cost and profitable technology to remove carbon dioxide from the atmosphere since plant supply and

sequester carbon at greater amount as compare to other technologies. The range of carbon supply by one plant is approximately 1 to 8 MgC ha⁻¹ yr⁻¹. Different part of trees sequesters carbon as biomass. Terrestrial carbon sequestration is a low cost mitigation option by trees. The accurate amount of carbon removed from the atmosphere could be measured through tree biomass that provide accurate estimation [9]. Trees in urban environment provide a lot of ecosystem services to the residents also remove carbon dioxide from the atmosphere during photosynthesis and tree also sequester carbon in different parts as biomass.

In view of the above and based on potential of tree species to sequester carbon for emission mitigation, the focus of this study is to understand role of trees in sequestering carbon in woody biomass growing along roadsides.

2. MATERIALS AND METHODS

2.1 Location

The study was conducted in district Multan (30.1575°N, 71.5249°E), province of Punjab, Pakistan. Multan is the fifth biggest population city of Pakistan and it is also known as the city of Saints. It is located in a corner generated by five rivers of central Pakistan. The city has developed to become a powerful political and economic epicenter for the country, with a dry port and admirable transport link. According to the Köppen–Geiger classification, Multan district fall within the desert climate (BWh). The average annual precipitation is 175mm and average annual temperature is 25.6°C.

2.2 Sampling

Four different road sites; *Khanewal* road (30.1356° N, 71.3329° E) [Site1], *Nangshah* road (30.1100° N, 71.4214° E) [Site2], *Shujabad* road (30.853° N, 71.2629° E) [Site3] and *Bosan* road (30.1647° N, 71.2951°E) [Site 4] were selected for sampling and data collection. The data were collected from 100 square meter along the roadside. Furthermore, three different trees species being excessively available/grown along roadsides as *V. nilotica* locally called as *Kikar*, *D. sissoo* locally called as *Shisham* and *E. camaldulenses* locally called as *sufaida* comprised of ten replications totaling twelve trees species along the road side with a total of 120

trees were selected from four different road side to estimate the carbon in biomass and soil. Tree heights were measured by clinometer and girth measured by steel tape. Girth at 1.37m above the ground level and terminal height for each tree's species were measured and recorded with in the 100 square meter along the roadside. For above ground and below ground biomass soil sample were collected through the equipment name auger. Soil sample was collected at the depth of 20 cm and 40 cm.

2.3 Biomass Determination

By using the allometric equation above ground and below ground biomass was estimated from collected data, belowground biomass was calculated from the above ground biomass supposed to be 26% [10]. By adding the above and below ground biomass the total biomass was obtained. For all tree species trees allometric equations was used to analyzed above ground and below ground biomass. From the growth parameter of trees (DBH and height) above ground and below ground biomass was calculated. The tress biomass was calculated in kgs which was then converted to biomass in Mg per hectare and Mg per hectare in carbon estimation.

Allometric equation were used to calculate the above and belowground biomass of tree species.

2.4 Estimation of Total Biomass

Total biomass was calculated by adding the above and below ground biomass:

$$\text{Total biomass} = \text{AGB} + \text{BGB}$$

2.5 Carbon Stock Estimation

Measuring the potential of carbon capture and storage in forests, accurate assessment of carbon in live tree tissue is required. Mostly regional and global scale assessments, carbon content has been assumed to 50% of tree biomass that is not accurate because of the great variation in carbon content among tree species. According to [14], Angiosperms in subtropical climate has a carbon content of approximately 48.1% in species biomass. As Pakistan has a subtropical type of climate and the study deals only with broad leaved species, so tree biomass was multiplied by 48.1% ($0.475 \times \text{biomass}$) for calculating carbon storage.

$$\text{Carbon} = 0.475 \times \text{biomass}$$

2.6 Soil Carbon Estimation

Soil carbon was assessed by the formula:

$$\text{SC (Mg per hectare)} = 10 \times \text{Bulk Density} \times 15 \times \text{Carbon} / 100 \times 10000 / 1000$$

Soil samples were collected from four different roadside at two depth (0-20cm) (20-40cm) by using soil auger. A total of 120 sample, 60 for each depth were collected. Sample were collected in polythene bags. Bulk density was measured by using [15] method, soil parameters used for the analysis were Soil pH, EC, Soil organic matter percentage, Soil carbon and Bulk density.

2.7 Statistical Analysis

All the data were analyzed by using Statistics 8.1 software. Furthermore, one way ANOVA including LSD (least significant difference between means) were also applied to analyze the difference between average values of the data.

3. RESULTS

3.1 Tree Height and Diameter

Results pertaining to diameter and height are presented in Table 2. Results revealed that in all the four site *E. camaldulensis* species had maximum height having 16.36 m, 12.96 m, 16.61m and 16.25 m respectively, whereas for diameter, *E. camaldulensis* had maximum diameter in S1-3, except S4 wherein *D. sissoo* showed higher diameter (37.04cm) to all species.

3.2 Species Biomass Estimation

The average biomass was recorded in S1 of species *E. camaldulensis* 13.7 Mg ha⁻¹, *V. nilotica* contain 0.98 Mg ha⁻¹, *D. sissoo* 1.01 Mg ha⁻¹. In S2, average biomass of *E. camaldulensis* 7.19 Mg ha⁻¹, *V. nilotica* contain 1.52 Mg ha⁻¹ and *D. sissoo* 1.02 Mg ha⁻¹. Whereas, in S3, *E. camaldulensis* showed average biomass of 13.05Mg ha⁻¹, *V. nilotica* contain 1.05 Mg ha⁻¹ and *D. sissoo* 0.98 Mg ha⁻¹. However, in S4, *E. camaldulensis* showed biomass of 6.55 Mg ha⁻¹, *V. nilotica* contain 2.41 Mg ha⁻¹ and *D. sissoo* 2.06 Mg ha⁻¹ (Table 3).

Table 1. Allometric equation for the calculation of above and below ground biomass

Species	Component	Allometric equation	Source
<i>D. sissoo</i>	AGB	$e^{-3.1141 * D^2} H^{0.9719}$	[11]
	BGB	$AGB^{0.26}$	
<i>V. nilotica</i>	AGB	$\text{LogY} = -1.0646 + 0.9098 \times \text{logD}^2 H$	[12]
	BGB	$\text{LogY} = -1.3952 + 0.8253 \times \text{logD}^2 H$	
<i>E. camaldulensis</i>	AGB	$\text{LnY} = -2.2660 + 2.4663 \times \text{ln D}^2 H$	[13]
	BGB	$BGB = AGB \times 0.26$	[10]

Table 2. Height and diameter of tree species

Site	Area	Species name	Height (m)	Diameter (cm)
1	S1	<i>D. sissoo</i>	11.24 ^b ± 1.64	25.64 ^b ± 4.21
		<i>V. nilotica</i>	9.96 ^c ± 0.70	21.35 ^c ± 2.26
		<i>E. camaldulensis</i>	16.36 ^a ± 0.94	40.93 ^a ± 3.56
2	S2	<i>D. sissoo</i>	11.64 ^{ab} ± 1.38	25.12 ^b ± 4.20
		<i>V. nilotica</i>	10.14 ^b ± 0.88	26.31 ^b ± 4.00
		<i>E. camaldulensis</i>	12.95 ^a ± 2.23	38.11 ^a ± 7.42
3	S3	<i>D. sissoo</i>	11.15 ^b ± 1.50	25.72 ^b ± 3.95
		<i>V. nilotica</i>	9.63 ^b ± 1.47	22.40 ^b ± 3.47
		<i>E. camaldulensis</i>	16.61 ^a ± 2.20	37.61 ^a ± 3.70
4	S4	<i>D. sissoo</i>	11.12 ^b ± 3.07	37.04 ^a ± 8.04
		<i>V. nilotica</i>	11.24 ^b ± 2.50	29.36 ^b ± 7.98
		<i>E. camaldulensis</i>	18.01 ^a ± 2.08	16.25 ^c ± 3.47

S1, Khaniwal road; S2, Nangshah road; S3, Shujaabad road; S4, Boson road

Table 3. Biomass of different tree species along the roadside

Site	Area	Scientific name	AGB Mg ha ⁻¹	BGB Mg ha ⁻¹	TB Mg ha ⁻¹
1	S1	<i>D. sissoo</i>	0.80 ^b ± 0.32	0.20 ^b ± 0.08	1.01 ^b ± 0.40
		<i>V. nilotica</i>	0.78 ^b ± 0.00	0.20 ^b ± 0.00	0.98 ^b ± 0.00
		<i>E. camaldulensis</i>	10.45 ^a ± 2.08	2.71 ^a ± 0.54	13.17 ^a ± 2.62
2	S2	<i>D. sissoo</i>	0.81 ^b ± 0.27	0.21 ^b ± 0.07	1.02 ^b ± 0.34
		<i>V. nilotica</i>	1.21 ^b ± 0.00	0.31 ^b ± 0.00	1.52 ^b ± 0.00
		<i>E. camaldulensis</i>	5.70 ^a ± 2.67	1.48 ^a ± 0.69	7.19 ^a ± 3.36
3	S3	<i>D. sissoo</i>	7.85 ^a ± 2.90	0.20 ^b ± 0.07	0.98 ^b ± 0.00
		<i>V. nilotica</i>	0.83 ^b ± 0.00	0.21 ^b ± 0.00	1.05 ^b ± 0.00
		<i>E. camaldulensis</i>	10.35 ^a ± 4.26	2.69 ^a ± 0.00	13.05 ^a ± 0.00
4	S4	<i>D. sissoo</i>	1.63 ^b ± 0.74	0.42 ^b ± 0.19	2.06 ^b ± 0.94
		<i>V. nilotica</i>	1.91 ^b ± 1.37	0.43 ^b ± 0.00	2.41 ^b ± 1.72
		<i>E. camaldulensis</i>	5.25 ^a ± 0.00	1.35 ^a ± 0.00	6.55 ^a ± 0.00

S1, Khaniwal road; S2, Nangshah road; S3, Shujaabad road; S4, Boson road

Table 4. Above ground, below ground, total carbon and carbon sequestration of different tree species along the roadside

Site	Area	Species	AGC kgha ⁻¹	BGC kg hac ⁻¹	TC kg hac ⁻¹	CO ₂
1	S1	<i>D. sissoo</i>	0.38 ^b ± 0.15	0.10 ^b ± 0.04	0.48 ^b ± 0.19	1.77 ^b ± 0.71
		<i>V. nilotica</i>	0.37 ^b ± 0.00	0.09 ^b ± 9.96	0.47 ^b ± 0.00	1.76 ^b ± 0.44
		<i>E. camaldulensis</i>	5.01 ^a ± 1.00	1.30 ^a ± 0.26	6.32 ^a ± 1.26	23.5 ^a ± 4.62
2	S2	<i>D. sissoo</i>	0.39 ^b ± 0.13	0.10 ^b ± 0.03	0.49 ^b ± 0.16	1.80 ^b ± 0.60
		<i>V. nilotica</i>	0.60 ^b ± 0.21	0.15 ^b ± 0.00	0.75 ^b ± 0.21	2.78 ^b ± 0.97
		<i>E. camaldulensis</i>	2.73 ^a ± 1.28	0.71 ^a ± 0.33	3.45 ^a ± 1.61	12.64 ^a ± 5.91
3	S3	<i>D. sissoo</i>	3.76 ^a ± 1.39	0.09 ^b ± 0.00	3.86 ^b ± 1.39	14.16 ^b ± 5.24
		<i>V. nilotica</i>	0.41 ^b ± 0.15	0.10 ^b ± 0.04	0.52 ^c ± 0.19	1.90 ^c ± 0.72
		<i>E. camaldulensis</i>	4.97 ^a ± 0.00	1.29 ^a ± 0.53	6.26 ^a ± 2.57	22.94 ^a ± 9.44
4	S4	<i>D. sissoo</i>	0.78 ^b ± 0.35	0.20 ^b ± 0.09	0.99 ^b ± 0.42	3.63 ^b ± 1.65
		<i>V. nilotica</i>	0.92 ^b ± 0.65	0.23 ^b ± 0.17	1.15 ^b ± 0.82	4.24 ^b ± 3.03
		<i>E. camaldulensis</i>	2.52 ^a ± 0.84	0.65 ^a ± 0.21	3.18 ^a ± 1.06	11.65 ^a ± 3.88

S1, Khaniwal road; S2, Nangshah road; S3, Shujaabad road; S4, Boson road

Table 5. Organic carbon and soil organic carbon of different tree species

Site	Area	Species name	OC 0-20	OC 20-40	0-20 SOC Mgh ¹	20-40 SOC Mgh ¹
1	S1	<i>D. sissoo</i>	0.56 ^a	0.39 ^b	13.8 ^a	7.86 ^a
		<i>V. nilotica</i>	0.61 ^a	0.42 ^{ab}	15.0 ^a	11.0 ^a
		<i>E. camaldulensis</i>	0.113 ^b	0.55 ^a	2.66 ^b	17.1 ^a
2	S2	<i>D. sissoo</i>	0.51 ^a	0.48 ^a	12.3 ^a	12.5 ^a
		<i>Vachellia nilotica</i>	0.39 ^{ab}	0.28 ^b	9.78 ^{ab}	7.56 ^b
		<i>E. camaldulensis</i>	0.24 ^b	0.35 ^{ab}	5.80 ^b	8.53 ^{ab}
3	S3	<i>D. sissoo</i>	0.44 ^a	0.37 ^a	10.82 ^a	9.62 ^a
		<i>V. nilotica</i>	0.59 ^a	0.29 ^a	14.61 ^a	7.65 ^a
		<i>E.camaldulensis</i>	0.55 ^a	0.37 ^a	13.51 ^a	9.05 ^a
4	S4	<i>D. sissoo</i>	0.42 ^b	0.53 ^a	10.47 ^b	13.86 ^a
		<i>V. nilotica</i>	0.58 ^{ab}	0.53 ^a	14.56 ^{ab}	13.99 ^a
		<i>E. camaldulensis</i>	0.66 ^a	0.48 ^a	15.89 ^a	11.67 ^a

S1, Khaniwal road; S2, Nangshah road; S3, Shujaabad road; S4, Boson road

3.3 Species Carbon Estimation

Data regarding above ground, below ground carbon estimation is presented in Table 4. *E. camaldulesnses* showed higher above ground carbon (AGC) contents (5.01^a±1±1.00), Below Ground (BGC) contents (1.30^a±0.26), Total carbon (TC) contents (6.32^a±1.26) and Carbon sequestration (23.5^a±4.62) than *V. nilotica* and *D. sissoo* in all four sites, whereas *V. nilotica* showed lowest values AGC, BGC, TC and Total Carbon to all species in all sites.

3.4 Organic Carbon and Soil Organic Carbon Estimation

Li et al., (2010) revealed that soil organic carbon density depend on layers of soil, changing in weather pattern and soil texture. Soil carbon density is different in first layer of soil as compare of the deeper layer of the soil in the roadside due to maximum occurrence of sand component in the soil.

Organic carbon and soil organic carbon were determined from four different roadside. Two depth soil sample were determined first depth (0-20) and second depth (20-40) their result are as shown in Table 5. The method used by measuring the soil organic carbon by Walkely Black Method [15].

4. DISCUSSION

Present study aimed to assess the potential of forest tree species along roadsides in Pakistan. The rate of carbon sequestration depends on growth factors of the plants. The major anxiety between policy maker and scientist monitoring the current level of atmospheric carbon dioxide through constraining harm of biodiversity, increase afforestation, decrease deforestation and reforestation [16,17]. Individual trees of

similar species may vary in their biophysical and biomass structure in various sites. Different researcher stated that number of plants and species diverse with the nature of species, grade of biotic disruption, edaphic condition, stand condition and structure of species [18]. Urban areas are associated with increased emissions due to heavy traffic load and industrial development. Last few years in urban cities increasing population but also the transportation emission and industrialization the carbon dioxide level increase in the atmosphere [19]. To build up the proper structure in Pakistan, it is important to aware the climate change mitigation and understand the advantages of mitigation among government sector and local people, through campaign among different areas, provincial level, as district-wise climate change systems bring source to express policy systems for advanced tendencies by distributing data near the effectiveness, climate change influences and acknowledgement of perception response choices [20]. Pakistan has a well-developed transport system for public and cargo and thus emit lion share of greenhouse gases. Trees being perennial vegetation can store large quantities of carbon in their above and below ground biomass and the same found true when tree species sequester large quantities of carbon [21] and help in mitigating vehicular emissions.

Soil act as important ingredient of the terrestrial ecosystem and mainly dependent upon soil type, terrain, climatic conditions and agriculture and forestry management practices [16,22]. Soil underneath trees carries reservoir of carbon pool being increased amount of litter with corresponding increased organic matter contents [23]. Present study findings were well supported by studies [22,24] that soil organic carbon decreases with decreasing soil depth.

5. CONCLUSIONS

The current study acts as a good point to know how species perform along roadsides in areas of vehicular pollution. Additional and more work is needed to test more species to lay a foundation of growing them across roadsides for the growth of vegetation for mitigating emissions. Hence urban trees especially alongside roads remained helpful in sequestering carbon to mitigate the harmful effects of vehicular pollution and hence should be included in annual plantation drive. This research shall also work twofold, not only to mitigate emissions but also to withstand hard weather wherein high temperature and scanty precipitation is a norm. Moreover, species identified such could also be used around other problem sites like that of industries to resolve environmental problems in Pakistan.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Da Silva AM, Braga Alves C, Alves SH. Roadside vegetation: Estimation and potential for carbon sequestration. *iForest-Biogeosciences & Forestry*. 2010;3(5):124.
- Hicks C, Woroniecki S, Fancourt M, Bieri M, Garcia Robles H, Trumper K, Mant R. The relationship between biodiversity, carbon storage and the provision of other ecosystem services. Cambridge: Critical Review for the Forestry Component of the International Climate Fund. 2014;102.
- Nowak DJ. Institutionalizing urban forestry as a “biotechnology” to improve environmental quality. *Urban Forestry & Urban Greening*. 2006;5(2):93-100.
- Bozena Š. Road-side herbaceous vegetation: Life history groups and habitat preferences. *Polish Journal of Ecology*. 2010;58(1):69-79.
- Dierkes C, Geiger WF. Pollution retention capabilities of roadside soils. *Water Science and Technology*. 1999;39(2):201-208.
- Akbostancı E, Tunç Gİ, Türüt-Aşık S. CO₂ emissions of Turkish manufacturing industry: A decomposition analysis. *Applied Energy*. 2011;88(6):2273-2278.
- Hussain M, Liu G, Yousaf B, Ahmed R, Uzma F, Ali MU, Butt AR. Regional and sectoral assessment on climate-change in Pakistan: Social norms and indigenous perceptions on climate-change adaptation and mitigation in relation to global context. *Journal of Cleaner Production*. 2018;200:791-808.
- Akbari H. Shade trees reduce building energy use and CO₂ emissions from power plants. *Environmental Pollution*. 2002;116:S119-S126.
- Chavan SP, Late AM, Bhosale BJ, Nalawade PM, Mule MB. Effect of vehicular pollution on plants: A case study from Aurangabad city (MS), India. In 2010 2nd International Conference on Chemical, Biological and Environmental Engineering. IEEE. 2010; 216-219.
- Cairns MA, Brown S, Helmer EH, Baumgardner GA. Root biomass allocation in the world's upland forests. *Oecologia*. 1997;111:1-11.
- Brown S, Gillespie AJ, Lugo AE. Biomass estimation methods for tropical forests with applications to forest inventory data. *For. Sci*. 1989;35:881-902.
- Rawat L, Luna RK, Kholiya D, Kamboj SK. Biomass, productivity and nutrient retention in *Acacia Catechu* Willd. Plantations in Shivalik Hills. *Indian Forester*. 2008;134:212-225.
- Hawkins T. Biomass and Volume Tables for *Eucalyptus camaldulensis*, *Dalbergia sissoo*, *Acacia auriculiformis* and *Cassia siamea* in the Central Bhabar-Terai of Nepal; University of Oxford: Oxford, UK; 1987.
- Thomas SC, Martin AR. Carbon content of tree tissues: A synthesis. *Forests*. 2012; 3:332-352.
- Walkley A. A critical examination of a rapid method for determining organic carbon in soils: Effect of variations in digestion conditions and of organic soil constituents. *Soil Sci*. 1947;63: 251-263.
- Kanowski P, McDermott C, Cashore B., Post-Copenhagen strategies for the implementation of REDD+. In: Richardson K, Steffen W, Liverman D et al. (Eds.), *Climate Change: Global Risks, Challenges and Decisions*. Cambridge University Press, New York USA, 2011;429-430.
- Pandey SS, Maraseni TN, Cockfield G. Carbon stock dynamics in different vegetation dominated community forests under REDD+: a case from Nepal. *Forest Ecology and Management*. 2014;327: 40-47.

18. Arora P, Chaudhry S. Vegetation and soil carbon pools of mixed plantation of *Acacia nilotica* and *Dalbergia sissoo* under social forestry scheme in Kurukshetra, India. J. Mater. Environ. Sci. 2017;8:4565–4572.
19. Wallace A. Reducing Carbon Emissions by Households: The Effects of Foot printing and Personal Allowances, De Montfort University, Leicester, UK; 2009. Available:<https://www.dora.dmu.ac.uk/dspace/handle/2086/2402>]
20. Reckien D et al. Equity, environmental justice, and urban climate change. Climate Change and Cities: Second Assessment Report of the Urban Climate Change Research Network ed C Rosenzweig et al. (New York: Cambridge University Press); 2018.
21. Zubair M, Yasin G, Qazlbash SK, UI Haq A, Jamil A, Yaseen M, Rahman SU, Guo W. Carbon Sequestration by Native Tree Species around the Industrial Areas of Southern Punjab, Pakistan. Land. 2022; 11(9):1577.
22. Yasin G, Nawaz MF, Yousaf MTB, Gul S, Qadir I, Niazi NK, Sabir MA. Carbon stock and CO₂ sequestration rate in linearly planted *Vachellia nilotica* farm trees. Pak. J. Agric. Sci. 2020;57: 807–814.
23. Takimoto A, Nair PR; Nair VD. Carbon stock and sequestration potential of traditional and improved agroforestry systems in the West African Sahel. Agric. Ecosyst. Environ. 2008;125:159–166.
24. Nawaz M, Yousaf M, Yasin G, Gul S, Ahmed I, Abdullah M, Rafay M, Tanvir M, Asif M., Afzal S. Agroforestry status and its role to sequester atmospheric CO₂ under semi-arid climatic conditions in Pakistan. Appl. Ecol. Environ. Res. 2018;16:645–661.

© 2022 Zubair et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92869>