



An Experimental Study on the Effect of Sulphur and Boron on Growth and Yield of Sesame (*Sesamum indicum*)

G. Bhanu Teja^{a#} and Shikha Singh^{a†}

^a Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i232582

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/94306>

Original Research Article

Received 25 September 2022
Accepted 01 December 2022
Published 05 December 2022

ABSTRACT

A field experiment was carried out during *Zaid* 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The soil of the experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.36 %), available N (171.48 kg/ha), available P (15.2 kg/ha) and available K (232.5 kg/ha). The experiment was laid out on Randomized Block Design with nine treatments each replicated thrice based on one year experimentation. The treatments are T₁: 10 kg/ha Sulphur + 1 kg/ha Boron, T₂: 10 kg/ha Sulphur + 1.5 kg/ha Boron, T₃: 10 kg/ha Sulphur + 2 kg/ha Boron, T₄: 20 kg/ha Sulphur + 1 kg/ha Boron, T₅: 20 kg/ha Sulphur + 1.5 kg/ha Boron, T₆: 20 kg/ha Sulphur + 2 kg/ha Boron, T₇: 30 kg/ha Sulphur + 1 kg/ha Boron, T₈: 30 kg/ha Sulphur + 1.5 kg/ha Boron, T₉: 30 kg/ha Sulphur + 2 kg/ha Boron are used. The results showed that application of 30 kg/ha Sulphur + 2 kg/ha Boron was recorded significantly higher plant height (96.22 cm), Plant dry weight (17.24 g/plant), whereas significantly highest Crop growth rate (8.74 g/m²/day) was recorded with the treatment 20 kg/ha Sulphur + 2 kg/ha Boron. However, Significantly maximum No. of capsules/plant (52.88), No. of Seeds/capsule (62.16), Test weight (3.29 g), Seed yield (1.35 t/ha), Stover yield (6.24 t/ha), Harvest index (17.77 %) as compared to other treatments.

Keywords: Sulphur; boron; sesame; growth and yield.

[#] M.Sc. Scholar;

[†] Assistant Professor;

*Corresponding author: E-mail: tejb048@gmail.com;

1. INTRODUCTION

Oil seed crops play the second most important role in the Indian agricultural economy next to food grains in terms of area and production. As Indian climate is suitable for the cultivation of oilseeds crop; large varieties of oilseeds are cultivated here. The government of India has been pursuing several development programs to meet the requirement of increasing demand of oilseeds in the country owing to increase in population, improvement in the standard of living and increasing industrial requirements. The concerted efforts of these development programs/ schemes register significant improvement in annual growth of yield and area under oilseed crops. Although country is having an increasing production trend in domestic oil seeds but only 50 per cent of the total requirement is met from domestic production and nearly half of the requirement is still being made through imports" [1].

"Among the oilseed crops, sesame (*Sesamum indicum* L.) is one of the world most important and oldest known oil seed crops (Abou Gharbia et al. 1997). Sesame is a member of the Pedaliaceae plant family. Its cultivation has started in 1500 BC in the Middle East, Asia & Africa. India is the major producer of sesame (*Sesamum indicum* L.) and ranks first in both area (1.78 M ha) and production (0.81 Mt) with average productivity of 455 kg ha⁻¹. Odisha produces 0.09 million tonnes sesame seeds annually with average productivity of 403 kg ha⁻¹" [2]. "Higher nutritional, medicine and cooking quality has recognized it as 'the queen of oilseeds'. There is a decline in productivity in sesame due to its cultivation in marginal and sub-marginal lands and moreover poor crop management practice.

Sulphur has long been recognized as one of the essential elements for plant growth particularly for oilseed crops. Sulphur is a constituent of three amino acids commonly found in plants viz., cystine, cysteine and methionine, which are essential components of proteins. Sulphur increases the oil content and gives pungency to oil as it forms certain disulphide linkages. Oilseed crops require more sulphur than cereals as their oil storage organs are mostly proteins, rich in S. Deficiency of sulphur is known to hamper N metabolism in plants as well as synthesis of S-containing amino acids and thus exerts adverse effects on both seed and oil yield. Sulphur (S) is essential for growth and development, plays a key role in plant metabolism, indispensable for the

synthesis of essential oils and plays a vital role in chlorophyll formation" [3].

"Among the micronutrient deficiency, boron deficiency is the second most dominant problem globally. Among the micronutrients, boron deficiency is one of the most widespread micronutrient deficiency in India. Plants require boron for several growth processes like development of meristematic tissue, proper pollination and seed set, translocation of sugars, starches. It has been reported that boron is required for pollen germination and pollen tube growth" (Dugger, 1983).

2. MATERIALS AND METHODS

The present examination was carried out during *Zaid* 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, UP, which is located at 25.28°N latitude, 81.54°E longitude and 98 m altitude above the mean sea level. The experiment laid out in Randomized Block Design which consists of nine treatments with T₁: 10 kg/ha Sulphur + 1 kg/ha Boron, T₂: 10 kg/ha Sulphur + 1.5 kg/ha Boron, T₃: 10 kg/ha Sulphur + 2 kg/ha Boron, T₄: 20 kg/ha Sulphur + 1 kg/ha Boron, T₅: 20 kg/ha Sulphur + 1.5 kg/ha Boron, T₆: 20 kg/ha Sulphur + 2 kg/ha Boron, T₇: 30 kg/ha Sulphur + 1 kg/ha Boron, T₈: 30 kg/ha Sulphur + 1.5 kg/ha Boron, T₉: 30 kg/ha Sulphur + 2 kg/ha Boron are used. "The experimental site was uniform in topography and sandy loam in texture, nearly neutral in soil reaction (P^H 7.1), low in Organic carbon (0.38%), medium available N (225 kg ha⁻¹), higher available P (19.50 kg ha⁻¹) and medium available K (213.7 kg ha⁻¹). In the period from germination to harvest several plant growth parameters were recorded at frequent intervals along with it after harvest several yield parameters were recorded those parameters are growth parameters, plant height, and plant dry weight are recorded. The yield parameters like Capsules/plant, Seeds/capsule, Test weight, Seed yield and stover yield were recorded and statistically analyzed using analysis of variance (ANOVA) as applicable to Randomized Block Design" (Gomez K.A. and Gomez A.A. 1984) [4].

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

3.1.1 Plant height

Significantly highest plant height (96.22 cm) was observed in the treatment with 30 kg/ha Sulphur

+ 2 kg/ha Boron over all the other treatments. However, the treatments with an application of 20 kg/ha Sulphur + 2 kg/ha Boron (95.46 cm) and 30 kg/ha Sulphur + 1.5 kg/ha Boron (95.73 cm) which were found to be at par with treatment 30 kg/ha Sulphur + 2 kg/ha Boron as compared to all the treatments.

“The presence of sulphur plays a vital role photosynthetic process of plant which has a direct bearing on plant growth and development” [5].

“The increase in plant height might be due to the involvement of Boron in different physiological processes like enzymes activation, electron transport, chlorophyll formation, stomatal regulation, etc. which gradually increased plant height” [6].

3.1.2 Plant dry weight (g/plant)

Treatment with 30 kg/ha Sulphur + 2 kg/ha Boron was recorded with significantly maximum dry weight (17.24 g/plant) over all the treatments. However, the treatments with 20 kg/ha Sulphur + 2 kg/ha Boron (17.06 g/plant) and 30 kg/ha Sulphur + 1.5 kg/ha Boron (16.95 g/plant) which were found to be statistically at par 30 kg/ha Sulphur + 2 kg/ha Boron.

Sulfur has a crucial part in the photosynthetic process of plants, which directly affects plant growth and development. Yadav et al. 2008 noted comparable outcomes. Boron administration often affects cell division, and nitrogen uptake from the soil may improve plant development as reflected in terms of plant dry weight. The findings were consistent with those of Mamatha et al. [7].

3.2 Yield Attributes and Yield

3.2.1 Capsules/plant

Significantly Maximum No. of capsules/plant (52.88) was recorded with the treatment of application of 30 kg/ha Sulphur + 2 kg/ha Boron over all the treatments. However, the treatments 20 kg/ha Sulphur + 2 kg/ha Boron (51.93) and 30 kg/ha Sulphur + 1.5 kg/ha Boron (52.43) which was found to be statistically at par with 30 kg/ha Sulphur + 2 kg/ha Boron.

3.2.2 Seeds/capsule

Significantly Highest No. of Seeds/capsule (61.26) was recorded with the treatment of

application of 30 kg/ha Sulphur + 2 kg/ha Boron over all the treatments. However, the treatment 20 kg/ha Sulphur + 2 kg/ha Boron (59.77) and 30 kg/ha Sulphur + 1.5 kg/ha Boron (60.54) which were found to be statistically at par with 30 kg/ha Sulphur + 2 kg/ha Boron.

The presented result revealed that sulphur application @ 30 kg/ha Sulphur recorded significantly at harvest, a number of capsules/plant and number of seeds/capsule. The present findings are in close agreement with those reported by Sriramchandracharan [8] Bhosale et al. [9].

“The application of boron to sesame generally improves capsule growth by synthesizing tryptophan and auxin. The enhancement effect on capsules/plant, seeds per capsule attributed to the favorable influence of boron application to crops on nutrient metabolism, biological activity and growth parameters which in turn influenced higher enzyme activity which in turn encouraged more capsules/plant and seeds/capsule” [10].

3.2.3 Test weight

Significantly highest Test weight (3.29 g) was recorded with the treatment application of 30 kg/ha Sulphur + 2 kg/ha Boron over all the treatments. However, the treatment with (3.05 g) in 20 kg/ha Sulphur + 2 kg/ha Boron and 30 kg/ha Sulphur + 1.5 kg/ha Boron (3.15 g) which were found to be statistically at par with 30 kg/ha Sulphur + 2 kg/ha Boron.

The number of capsules/plant, and test weight in sesame increased dramatically as Sulphur fertilization levels increased up to 30 kg/ha, while the number of seeds/capsules also increased significantly. Shinde et al. [11] found outcomes that were comparable.

3.2.4 Seed yield (t/ha)

Significantly highest Seed yield (1.35 t/ha) was recorded with the treatment application of 30 kg/ha Sulphur + 2 kg/ha Boron over all the treatments. However, the treatments with (1.14 t/ha) in 20 kg/ha Sulphur + 2 kg/ha Boron and 30 kg/ha Sulphur + 1.5 kg/ha Boron (1.25 t/ha) which were found to be statistically at par with 30 kg/ha Sulphur + 2 kg/ha Boron.

Therefore, 40 kg/ha sulphur was found a suitable for getting higher yield of sesame. Yadav et al. (1996); Jadav et al. [12] observed that the seed

and stover yield of sesame increased with application of sulphur. The higher and positive response to added sulphur may be assigned to low to medium status of available S of the soil under study and due to the stimulating effect of applied S in the synthesis of chloroplast and greater photosynthesis efficiency, increased dry matter yield of the plant.

“Boron plays a vital role in increasing because it takes place in many physiological processes of plants such as chlorophyll formation, stomatal regulation, starch utilization which enhanced seed yield” [13].

3.2.5 Stover yield (t/ha)

Significantly highest Stover yield (6.24 t/ha) was recorded with the treatment application of 30 kg/ha Sulphur + 2 kg/ha Boron over all the

treatments. However, the treatments with (5.88 t/ha) in 20 kg/ha Sulphur + 2 kg/ha Boron and 30 kg/ha Sulphur + 1.5 kg/ha Boron (6.11 t/ha) which were found to be statistically at par with 30 kg/ha Sulphur + 2 kg/ha Boron.

The increase in stover yield may be attributable to sulphur, which is essential for plants' photosynthetic processes and has a direct impact on plant growth and development. The outcomes support the conclusions made by Yadav et al. in 2008. Increased amounts of boron were associated with the best outcomes because they may have a positive impact on plant metabolism, biological process activity, and photosynthetic pigment and enzyme activity. These effects in turn promoted vegetative development and higher biomass accumulation. Similar findings were made by Ravichandra et al. [14].

Table 1. Effect of Sulphur and Boron on growth attributes of Sesame

| Treatments | Plant height (cm) | Dry weight (g/plant) |
|---------------------------------------|-------------------|----------------------|
| 1. 10 kg/ha Sulphur + 1 kg/ha Boron | 91.88 | 15.43 |
| 2. 10 kg/ha Sulphur + 1.5 kg/ha Boron | 92.33 | 15.65 |
| 3. 10 kg/ha Sulphur + 2 kg/ha Boron | 93.83 | 16.10 |
| 4. 20 kg/ha Sulphur + 1 kg/ha Boron | 93.32 | 15.89 |
| 5. 20 kg/ha Sulphur + 1.5 kg/ha Boron | 94.10 | 16.33 |
| 6. 20 kg/ha Sulphur + 2 kg/ha Boron | 95.46 | 16.95 |
| 7. 30 kg/ha Sulphur + 1 kg/ha Boron | 94.78 | 16.75 |
| 8. 30 kg/ha Sulphur + 1.5 kg/ha Boron | 95.73 | 17.06 |
| 9. 30 kg/ha Sulphur + 2 kg/ha Boron | 96.22 | 17.24 |
| F- test | S | S |
| S. EM (\pm) | 0.29 | 0.11 |
| C. D. (P = 0.05) | 0.86 | 0.33 |

Table 2. Effect of sulphur and boron on yield attributes and yield of sesame

| Treatments | Capsules/ plant | Seeds/ capsule | Test Weight (g) | Seed yield (t/ha) | Stover yield (t/ha) | Harvest Index (%) |
|---------------------------------------|-----------------|----------------|-----------------|-------------------|---------------------|-------------------|
| 1. 10 kg/ha Sulphur + 1 kg/ha Boron | 48.67 | 55.87 | 2.33 | 0.75 | 4.82 | 13.47 |
| 2. 10 kg/ha Sulphur + 1.5 kg/ha Boron | 49.68 | 56.79 | 2.44 | 0.80 | 4.92 | 13.97 |
| 3. 10 kg/ha Sulphur + 2 kg/ha Boron | 50.54 | 57.82 | 2.75 | 0.93 | 5.15 | 15.28 |
| 4. 20 kg/ha Sulphur + 1 kg/ha Boron | 50.30 | 57.23 | 2.56 | 0.88 | 4.96 | 15.03 |
| 5. 20 kg/ha Sulphur + 1.5 kg/ha Boron | 51.09 | 58.71 | 2.83 | 1.03 | 5.42 | 15.92 |
| 6. 20 kg/ha Sulphur + 2 kg/ha Boron | 51.93 | 59.77 | 3.05 | 1.14 | 5.88 | 16.18 |
| 7. 30 kg/ha Sulphur + 1 kg/ha Boron | 51.49 | 59.36 | 2.96 | 1.06 | 5.74 | 15.57 |
| 8. 30 kg/ha Sulphur + 1.5 kg/ha Boron | 52.43 | 60.54 | 3.15 | 1.25 | 6.11 | 17.00 |
| 9. 30 kg/ha Sulphur + 2 kg/ha Boron | 52.88 | 61.26 | 3.29 | 1.35 | 6.24 | 17.77 |
| F test | S | S | S | S | S | S |
| S. EM (\pm) | 0.33 | 0.55 | 0.10 | 0.07 | 0.12 | 0.54 |
| CD (P = 0.05) | 0.98 | 1.65 | 0.30 | 0.22 | 0.37 | 1.61 |

3.2.6 Harvest index (%)

Significantly highest Harvest index (17.77 %) was recorded with the treatment application of 30 kg/ha Sulphur + 2 kg/ha Boron over all the treatments. However, the treatments with (16.18 %) in 20 kg/ha Sulphur + 2 kg/ha Boron and 30 kg/ha Sulphur + 1.5 kg/ha Boron (17.00 %) which were found to be statistically at par with 30 kg/ha Sulphur + 2 kg/ha Boron.

4. CONCLUSION

In conclusion, grain production was considerably greater (1.35 t/ha) when treatment 30 kg/ha Sulphur + 2 kg/ha Boron was applied compared to other treatments. Since the results are based on research conducted over a single season.

ACKNOWLEDGEMENT

I express thankfulness to my advisor Dr. Shikha Singh and all the faculty members of Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj -211007, Uttar Pradesh. For providing us essential facilities to undertake the studies.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Modhavadiya VL, Solanki MS, Hirpara DV, Jadeja AS, Ranpariya VS. Effect of sulphur and boron on yield of summer Sesamum (*Sesamum indicum* L.) under medium black calcareous soils of south Saurashtra region of Gujarat. International Journal of Chemical Studies. 2017;6(1):378-380.
2. Murmu S, Murmu K, Satapathy M. Effect of sulphur fertilization on growth, yield and quality of sesamum (*Sesamum indicum* L.) in Mid Central Zone of Odisha. International Journal Biological Research. Env. Agril. Sci. 2015;1(1):5-12.
3. Singh MK, Singh TK, Singh UN. Effect of nitrogen and sulphur levels on yield and quality of linseed under dryland conditions. Journal of Oilseeds Research. 2000; 17(1):162-164.
4. Akram SW, Singh R, Tripathi P, Roopa G. Effect of phosphorus and sulphur on yield and economics of wheat (*Triticum aestivum* L.). The Pharma Innovation Journal. 2022;11(5):599-602.
5. Yadav RA, Tripathi AK, Yadav AK. Effect of micronutrients in combinations with organic manures on production and net returns of Sesame (*Sesamum indicum* L.) in Bundelkhand tract of Uttar Pradesh. Annal Agricultural Research. 2008;30(1): 53-58.
6. Choudhary, Md. Ullah MU, Md Rahman H, Md. Shahidullslam A. Effect of boron and nitrogen fertilization on cowpea growth, nodulation and grain yield in Rangamati, Bangladesh. Legume Research. 2000; 23(1):9-14.
7. Mamatha K, Vidya S, Laxmi N. Influence of different levels of boron along with application of FYM on nutrient uptake and yield of Sesame (*Sesamum indicum* L.). International Journal of Farm Sciences. 2016;6(2):37-45.
8. Sriramchandrasekharan MV. Integrated Sulphur management on sesame. Journal of Interacademia. 2004;8(2):187-190.
9. Bhosale ND, Dabhi BM, Gaikwad VP, Agarwal MC. Influence of potassium and sulphur levels on yield, quality and economics of sesamum (*Sesamum indicum* L.) International Journal of Plant Sciences. 2011;6(2):335-337.
10. Yadav SN, Singh SK, Omkar K. Effect of boron on yield attributes, seed yield and oil content of mustard (*Brassica juncea* L.) on an inceptisol. Journal of the Indian Society of Soil Science. 2016;64(3):291-296.
11. Shinde SD, Raskar BS, Tamboli BD. Effect of spacing and sulphur levels on productivity of sesame (*Sesamum indicum* L.) under summer condition. Journal of Maharashtra Agricultural Universities. 2011;36:1:28-31.
12. Jadav DP, Padamani DR, Polara KB, Parmar KB, Babaria NB. Effect of different level of sulphur and potassium on growth, yield and yield attributes of sesame (*Sesamum indicum* L.). An Asian Journal of Soilscience. 2010;5(1):106-108.
13. Mallik K, Raj. Effect of different levels of phosphorus, sulphur and boron on growth and yield of sesame (*Sesamum indicum* L.). Journal of Oilseeds Research. 2015; 29(2):75-77.

14. Ravichandra K, Nagajyothi B, Jaipal S, Joy D, Krupakar A. Growth of groundnut (*Arachis hypogea* L.) and its yield as influenced by foliar spray of Boron along with rhizobium inoculation. Indian Journal of Dryland Agriculture and Development. 2015;30(1):60-63.

© 2022 Teja and Singh; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/94306>