



Comparative Analysis of Vision Screening Guidelines in the United States

Nathan J. Brown ^{a,b} and John A. Musser VI ^{b,c*}

^a Elson S. Floyd College of Medicine, Washington State University, Spokane, WA, USA.

^b Rural Ophthalmology Optometry Treatment & Screening 501(c)(3) Non-Profit, Little Rock, AR, USA.

^c Global Outreach Division, John A. Moran Eye Center, University of Utah, Salt Lake City, UT, USA.

Authors' contributions

This work was carried out in collaboration between both authors. Author NJB performed the statistical analysis, wrote the protocol, managed the analyses of the study and assisted with manuscript drafting. Author JAM designed the study, managed the analyses of the study and assisted with manuscript drafting. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/OR/2022/v17i330259

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91321>

Short Research Article

Received 17 June 2022
Accepted 25 August 2022
Published 30 August 2022

ABSTRACT

Aims: To assess current school vision screening guidelines of states spread across different regions of the United States to inform quality guideline parameters and help combat preventable pediatric vision loss.

Study Design: Cross-sectional comparative analysis.

Methodology: States were first ordered alphabetically and then selected using an online random number generator (Alphabet Inc., Mountain View, California). States were selected until 13 available screening guidelines were identified. Each guideline was assessed using a 10-point multi-factorial scoring criteria detailed in Table 1. Descriptive statistics (mean and standard deviation) were calculated for each scoring criteria using Microsoft® Excel (version 16.63.1, Redmond, WA).

Results: Most state guidelines included what ages to screen (84.62%, 11/13), how to screen (84.62%, 11/13), and how to follow-up with students to arrange (92.31%, 12/13) and confirm eye care delivery (76.92%, 10/13). Sadly, only the minority of state guidelines described at least two main causes of amblyopia (46.15%, 6/13), and a less than one-third of school nurse vision screening guidelines discussed the window of time to save vision in amblyopia (30.77%, 4/12). Worse yet, very few nurse vision screening guidelines explained that subjectively a child can't tell you if they are at risk of developing amblyopia (15.38%, 2/13), or included two treatments for amblyopia (7.69%, 1/13) in the vision screening educational program.

*Corresponding author: E-mail: john.a.musser@utah.edu;

Conclusion: This study found that most of the assessed vision screening guidelines outlined the process of screening and follow-up but failed to emphasize why screening is important, causes, and treatment options for amblyopia. Gaps in these training guidelines may contribute to delayed recognition and treatment for amblyopia—the leading cause of irreversible pediatric vision loss. Further improvements to vision screening guidelines are needed for school nurses and paraprofessional staff in the majority of the states evaluated.

Keywords: Amblyopia; lazy eye; vision screening; pediatric blindness.

1. INTRODUCTION

Amblyopia, also called lazy eye, is the leading cause of preventable pediatric vision loss [1]. It occurs early on in a child's life when the neural connections of the brain are developing with the eyes. Unfortunately, when the brain does not properly obtain stimulation from one eye, it gradually loses neural connections with that eye and ultimately the ability to see from that eye and control its position, hence lazy eye [2]. After age 7, structural and functional brain damage of the occipital lobe may be permanent [3]. However, amblyopia is often reversible if treated before the age of 5 [4,5]. A 2020 study published by Fu et al. in the British Journal of Ophthalmology recently determined that the 95% confidence interval of amblyopia prevalence is between 1.17% and 1.78% [6]. The most common cause of amblyopia is uncorrected refractive error [7]. Alternatively, more serious pathologies (cataracts, eyelid ptosis, vitreous hemorrhage, and cornea opacities) can cause stimulus loss resulting in what is termed deprivation amblyopia [8,9]. Additionally, Simmons found that children in medically underserved areas are at a disproportionately higher risk for underdiagnosis [10].

While not mandated by every state, school vision screenings serve a critical role in the early detection of amblyopia and other severe eye illnesses [11,12,13]. Many states' Department of Education provide screening guidelines to assist school nurses in these efforts. Once an eye care referral is made, children should be investigated for ocular pathology or the need for eyeglasses. If amblyopia is present, treatment for children aged 7 and younger is often an eye patch or atropine eye drops, both affordable and effective options [14]. By worsening vision in the stronger eye with the patch or atropine drops, the weaker eye can begin to catch up. The brain in young children begins to strengthen connections with the previously deprived eye with restoration in both vision and tracking abilities. Unfortunately, for children aged 8 and older this developmental

window has largely closed, and total restoration of vision and tracking is not as common. In fact, older children may require invasive ocular alignment surgery from a pediatric ophthalmologist with only partial improvement [15]. The purpose of this study was to conduct and present a comparative analysis of vision screening guidelines of 13 states spread across different regions of the United States.

2. METHODOLOGY

States were first ordered alphabetically and then selected using an online random number generator (Alphabet Inc., Mountain View, California). States were selected until 13 available screening guidelines were identified. Each guideline was assessed using a 10-point multi-factorial scoring criteria detailed in Table 1. Criteria were selected in three different domains of pediatric vision care: 1. Underlying cause 2. Screening approach 3. Follow-up and pediatric eye care importance. Descriptive statistics (mean and standard deviation) were calculated for each scoring criteria using Microsoft® Excel (version 16.63.1, Redmond, WA).

3. RESULTS AND DISCUSSION

Of the 27 states randomly selected, only 48.1% (13/27) provided public access to vision screening guidelines through each respective Department of Education website. The available guidelines were assessed on a 10-point grading scale. Scores ranged from 2-9 points with a mean of 5.25 points and a standard deviation of 2.33 points. Most state guidelines included what ages to screen (84.62%, 11/13), how to screen (84.62%, 11/13), and how to follow-up with students to arrange (92.31%, 12/13) and confirm eye care delivery (76.92%, 10/13).

Surprisingly, just over half of state guidelines referred to false negative and/or false positive screening risks (53.85%, 7/13) or described the main cause of preventable pediatric blindness—amblyopia (53.85%, 7/13). Sadly,

only the minority of state guidelines described at least two main causes of amblyopia (46.15%, 6/13), and a less than one-third of school nurse vision screening guidelines discussed the window of time to save vision in amblyopia (30.77%, 4/12). Worse yet, very few nurse vision screening guidelines explained that typically a child cannot tell you if they are at risk of developing amblyopia (15.38%, 2/13) because

their brain utilizes the good eye and on a day-to-day basis there is normal visual acuity in the good eye. Only one state (Colorado) provided education on the two treatments for amblyopia (7.69%, 1/13). Further data analysis and scores are summarized in Tables 2 and 3. Fig. 1 provides a visual representation of the 13 states analyzed and their vision screening education ratings.

Table 1. Multi-factorial scoring criteria

Scoring data	
1	Outlines appropriate follow-up process? Yes = 1 pt. No = 0 pt
2	Describes how to screen for amblyopia? Yes = 1 pt. No = 0 pt
3	Is pre-kindergarten (age 5) vision screening advised? Yes = 1 pt. No = 0 pt
4	Outlines process to confirm eye care delivery? Yes = 1 pt. No = 0 pt
5	Describes what amblyopia is? Yes = 1 pt. No = 0 pt
6	References false negative/positive risk due to young age or bias? Yes = 1 pt. No = 0 pt
7	Describes at least two causes amblyopia? Yes = 1 pt. No = 0 pt
8	Describes the window of time to save vision (before age 6 or 7)? Yes = 1 pt. No = 0 pt
9	Answers if a child can tell you if they have amblyopia? Yes = 1 pt. No = 0 pt
10	Describes how amblyopia is treated for at least two causes? Yes = 1 pt. No = 0 pt

Table 2. Scoring criteria and outcomes for 13 state vision guidelines

Scoring Criteria	Yes	Yes (%)	No	No (%)
Outline appropriate follow up process?	12	92.31%	1	7.69%
Describes how to screen for amblyopia?	11	84.62%	2	15.38%
Is pre-kindergarten (age 5) vision screening advised?	11	84.62%	2	15.38%
Outlines process to confirm eye care delivery?	10	76.92%	3	23.08%
Describes what amblyopia is?	7	53.85%	6	46.15%
References false negative/positive risk due to young age or bias?	7	53.85%	6	46.15%
Describes at least two causes of amblyopia?	6	46.15%	7	53.85%
Describes the window of time to save vision (before age 6 or 7)?	4	30.77%	9	69.23%
Answers if a child can tell you if they have amblyopia?	2	15.38%	11	84.62%
Describes how amblyopia is treated for at least two causes?	1	7.69%	12	92.31%




Fig. 1. Visual representation of the 13 states analyzed and their vision screening education ratings

Table 3. The 13 states included in analysis with corresponding scores

State	Score
Colorado	9
Minnesota	8
Arizona	7
California	7
Louisiana	7
Connecticut	7
Missouri	6
Rhode Island	5
West Virginia	5
Oregon	3
Tennessee	3
Massachusetts	2
North Carolina	2
Average Score	5.25
Standard Deviation	2.33

4. CONCLUSION

This study found that most of the assessed vision screening guidelines outlined the process of screening and follow-up but failed to emphasize the main cause and treatment of amblyopia or why screening is important before the critical period of visual development. Inconsistencies between these guidelines may contribute to delayed recognition and treatment for amblyopia—the leading cause of irreversible pediatric vision loss. Leukocoria, defined as “white pupil,” is an ominous sign that school nurses should be trained to recognize. Cataracts and retinoblastoma, two potential causes of leukocoria, can greatly affect a child’s life and educational opportunities. In the case of retinoblastoma, delayed follow-up can be fatal.

Further additions to screening guidelines are needed in order to provide school nurses and paraprofessional staff a deeper understanding of amblyopia and why early intervention is crucial. Improving the quality of school vision screening education is an essential task to help combat preventable pediatric blindness.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The Washington State University Institutional Review Board determined that this study was exempt and not considered human subjects research.

ACKNOWLEDGEMENTS

This investigation was not supported by a funding agency. All authors attest that they meet the current ICMJE criteria for authorship.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. McConaghy JR, McGuirk R. Amblyopia: Detection and Treatment. *Am Fam Physician*. 2019;100(12):745-750.
2. de Ribot FM, Feldman BH, Bacal DA, et al. Amblyopia. *EyeWiki*. Available: <https://eyewiki.aao.org/Amblyopia>. Published July 3, 2022. Accessed August 11, 2022
3. Kates MM, Beal CJ. Amblyopia. *JAMA*. 2021;325(4):408. DOI:10.1001/jama.2020.5741
4. Hunter D, Cotter S. Early diagnosis of amblyopia. *Vis Neurosci*. 2018;35:E013. DOI:10.1017/S0952523817000207
5. Webber AL, Wood J. Amblyopia: Prevalence, natural history, functional effects and treatment. *Clinical and Experimental Optometry*. 2005;88(6):365-375.
6. Fu Z, Hong H, Su Z, Lou B, Pan C-W, Liu H. Global prevalence of amblyopia and disease burden projections through 2040: A systematic review and meta-analysis. *British Journal of Ophthalmology*. 2020;104(8): 1164-1170. Available: <http://dx.doi.org/10.1136/bjophthalmol-2019-314759>
7. Shakarchi AF, Guo X, Friedman DS, Repka MX, Collins ME. Vision Needs of Children Who Failed School-based Vision Screening with and without Eyeglasses. *Ophthalmic Epidemiol*. 2021;28(2):131-137. DOI:10.1080/09286586.2020.1800754
8. Magdalene D, Bhattacharjee H, Choudhury M, Multani PK, Singh A, Deshmukh S, Gupta K. Community outreach: An indicator for assessment of prevalence of amblyopia. *Indian Journal of Ophthalmology*. 2018; 66(7):940.
9. Eslayeh AH, Omar R, Fadzil NM. Refractive amblyopia among children aged 4–12 years in a hospital-based setting in Gaza Strip, Palestine. *Medical Hypothesis Discovery and Innovation in Ophthalmology*. 2021;10(3): 107-113.
10. Simmons K. Amblyopia characterization, treatment, and prophylaxis. *Survey of Ophthalmology*. 2005;50(2):123-166. Available: <https://doi.org/10.1016/j.survophthal.2004.12.005>
11. Jullien S. Vision screening in newborns and early childhood. *BMC Pediatr*. 2021;21 (Suppl 1):306. Published 2021 Sep 8 DOI:10.1186/s12887-021-02606-2
12. Allen D, Contreras P, Daviet R, Elliott K. ADHS Vision Screening Guidelines Task Force. Recommended vision screening guidelines for children ages 3 and older. Available: <https://www.azed.gov/sites/default/files/2018/04/visioncreening2010.pdf?id=5acc2bf3217e114c088c729>. Published December 2010. Accessed August 11, 2022.
13. Azizoğlu S, Crewther SG, Şerefhan F, Barutçu A, Göker S, Junghans BM. Evidence for the need for vision screening of school children in Turkey. *BMC Ophthalmology*. 2017 Dec;17(1):1-4.
14. Jefferis JM, Connor A, Clarke MP. Amblyopia. *BMJ*. 2015;351. Available: <https://doi.org/10.1136/bmj.h5811>
15. Silverstein M, Scharf K, Mayro EL, et al. Referral outcomes from a vision screening program for school-aged children. *Can J Ophthalmol*. 2021;56(1):43-48. DOI:10.1016/j.jcjo.2020.07.009

© 2022 Brown and Musser; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91321>