



Efficacy of Different Levels of NPK against Anthracnose of King Chilli (*Capsicum chinense*) Caused by *Colletotrichum gloeosporioides* (Penz.) and Impact on Their Growth Parameters and Yield in Manipur

**Sorokhaibam Romio Singh^a, N. Tiameren Ao^a,
Hijam Meronbala Devi^{a*}, Sh. Herojit Singh^b,
Siman Debbarma^a, L. Daiho^a and Hijam Shila Devi^c**

^a Department of Plant Pathology, School of Agricultural Sciences and Rural Development, Nagaland University, Medziphema Campus-797106 Nagaland, India.

^b Department of Agricultural Economics, School of Agricultural Sciences and Rural Development, Nagaland University, Medziphema Campus-797106 Nagaland, India.

^c Department of Entomology, School of Agricultural Sciences and Rural Development, Nagaland University, Medziphema Campus-797106 Nagaland, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/V12i121501

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/94278>

Original Research Article

Received: 27/10/2022

Accepted: 12/12/2022

Published: 14/12/2022

*Corresponding author: E-mail: meronbaladevi@nagalanduniversity.ac.in;

ABSTRACT

A significant disease in Capsicum that reduces crop output is anthracnose. In the current study, the effectiveness of various NPK dosages (N₁-120 Kg, N₂ -150 Kg, N₃-90 Kg, P₁-60 Kg, P₂-75 Kg, P₃-45 Kg, K₁-60 Kg, K₂-75 Kg, K₃-45 Kg) against *Colletotrichum gloeosporioides* which caused Anthracnose diseases in King chilli and their effects on growth parameters and yield were assessed. The experiment described in Randomized Block Design (RBD) was conducted at KVK Thoubal, Manipur, from October 2021 to April 2022. Results showed that amongst the fertilizers, application of increased doses of P and K exhibited significant management of anthracnose of King chilli with 25.00% and 19.66% disease severity and 'Control' (60.00%) revealed high disease severity and reduced yield. The increased level of P and K also produced maximum yield with 11.17 Kg/plot and 9.30 Kg/plot in fresh weight respectively. Thus application with increased level of P and K has paramount importance in combating anthracnose in King chilli which significantly increase in the growth parameters and yield.

Keywords: King chilli; anthracnose; *Colletotrichum gloeosporioides*; NPK.

1. INTRODUCTION

Chilli (*Capsicum annum* Lin.) belonging to the family Solanaceae is a high value cash crop. The introduction of chilli peppers in Asia is believed to be around 450-500 years old [1]. Chilli is a necessary spice of every Indian household and used in many ways such as spices, condiments, vegetables, sauce, pickles and chutney for culinary purpose [2]. Chillies are good resource for vitamins like A, B and C, among this Chilli contains more amount of vitamin C compared to Orange. Chilli also contains minerals like Ca, P, Fe, Na, Cu and steam volatile oils, carotenoids, fatty oils etc. [3].

The King chilli (*Capsicum chinense* Jacq.) is known globally for its spice quality and aroma which is indigenous to the North Eastern states of India. It is also a good source of income for the local farmers. Different local names are given to this chilli such as *Naga chilli* (Nagaland), *Bhut Jolokia* (Assam), and *U-Morok* (Manipur) [4,5]. It is mainly cultivated in the state of Nagaland, Assam, Manipur and to some level in Mizoram, Arunachal Pradesh and Meghalaya. It is also grown in the North Eastern part of Bangladesh [6].

The *Naga Raja Mircha* or *Naga Joloakia* is considered as the hottest chilli variety in the world, measuring 855,000 SHU [7]. The hotness or hot flavor of chillies are due to the presence of a group of seven closely related compounds called capsaicinoids, but capsaicin and dihydro capsaicin are responsible for their pungency. This is measured in Scoville Heat Units (SHU) or determined by high Performance Liquid Chromatography HPLC.

Anthracnose has been derived from the Greek word which means 'coal'. This is a name for diseases in plants that is determined by dark or dull colour, lesions that are sunken and also containing spores of the pathogen [8]. Normally, the symptoms of anthracnose on fruits of chilli show sunken necrosis with concentric rings of acervuli. Blemished fruits have reduced market value [9].

Anthracnose or fruit rot is a highly destructive pathogen that attacks chilli and causing major crop losses [10]. *Colletotrichum* spp. causing anthracnose includes *C. acutatum*, *C. gloeosporioides* and *C. coccodes* [11]. Among all these, the most destructive and widely distributed are *C. gloeosporioides* and *C. acutatum* [12,13].

The King chilli which is grown in Manipur endures huge losses due to *C. gloeosporioides* causing fruit rot disease [14]. This chilli, like any other Capsicum, suffers from the attack of various pest and diseases, of which anthracnose of fruit rot is considered as the most damaging one. The pathogen *C. gloeosporioides* attack leaf, twig and fruit of chilli resulting in die back.

Under the climatic conditions of Manipur with heavy rainfall and humidity, anthracnose or fruit rot disease on King chilli caused by the fungus *Colletotrichum gloeosporioides* has been found to be a limiting factor in optimum crop production [14]. Although management of the pathogen focus on cultural practices, bio-agents, fungicides and genetic resistance, very little is currently known about the influence of plant nutrition and sowing dates on anthracnose of chilli. Hence, the present study was undertaken on management of anthracnose in King chilli by

applying different level of NPK and different dates of sowing.

2. MATERIALS AND METHODS

The present investigations were carried out at the experimental field of KVK Thoubal, Manipur during 2021-22 entitled "Effect of different levels of NPK and sowing dates against anthracnose of King chilli (*Capsicum Chinense*) caused by *Colletotrichum gloeosporioides* in Manipur."

2.1 Isolation and Identification of the Pathogen

The pathogen *Colletotrichum gloeosporioides* was isolated from naturally infected King chilli plants showing typical symptoms of the disease. The infected parts of the samples showing typical symptoms were cut into small pieces. The cut pieces were surface sterilized in 1.0% of sodium hypochlorite solution for 30 second and then rinsed three times with sterile distilled water to remove traces of the sodium hypochlorite, dried in sterilized tissue paper and then inoculated artificially on potato dextrose agar (PDA) medium in Petri plates. The mycelium growing out of the plant tissue were sub-cultured on PDA medium and incubated at $25 \pm 1^\circ\text{C}$ for 7 to 10 days. Pure culture of the pathogen was obtained by single hyphal tip method.

The mycelium of 7days old culture was taken on a glass slide with the help of needle and the colony characterization was done [15]. The pathogen was identified based on its mycelial, conidial characteristics following standard mycological keys [16].

2.2 Maintenance and Preservation of Culture

The stock cultures were maintained on PDA slants in refrigerator at 4°C and the pathogen was sub-cultured at regular intervals for maintaining the live culture for further studies.

2.3 Field Trial

2.3.1 Sowing in nursery

A susceptible local variety of King chilli seeds used in this investigation was obtained from the local farmers of S. B Nursery farm Nepra-company, Thoubal. The Nursery bed was prepared by clearing all the weeds and

measuring 3m×2m size. The soil of the bed was prepared by applying 2 kg Vermicompost, 2 kg Vermiculite and 1 kg Coco peat before 2 days of sowing. Clean seeds were then selected and sown in the nursery plot on 19th October 2021. Seeds were sown in line keeping 5cm gap between each line at a depth of 3-5cm and covered thinly by soil. Irrigation was provided twice a day till transplanting stage of the plant.

2.3.2 Land preparation

Primary tillage for land preparation was done by tractor followed by secondary tillage 2-3 times with harrow to crush the clods and uproot and shred the remaining weeds and stubbles with the help of power tiller. This was done in order to attain the soil to become good tilt condition. The mixture of 3 kg Vermicompost, 2 kg Vermiculite, 1 kg Coco peat, 1 kg of rice husk, 1 kg of Charcoal powder were added to each and every plot for enhancing the growth, germination and aeration of the plant after 1 week of deep tillage. Transplanting was done 15 days after the application according to the layout and design of the experiment.

2.3.3 Transplanting

Transplanting was done when the seedlings attained the height of 10-15cm (5-6 weeks after sowing). Before pulling out the seedlings, light irrigation was given to loosen the soil, in order to avoid the root damage during uprooting of the seedlings.

2.3.4 Duration of Experiment

The field experiment was conducted under agro-climatic condition of KVK Thoubal, during October, 2021 to May, 2022.

2.4 Experiment Design

2.4.1 Evaluation of different levels of NPK

Design of the experiment was RBD, Number of replication = 3, Number of King chilli cultivar = 1(local Raja Mircha), Treatment =10, Chemical nutrients applied were Urea (N 46%), DAP (18 % N and 46 % P_2O_5), MOP (KCl, 66.5% K), Total no. of plot = 30, Plot size = 2 x1.5 m², Row to row = 60cm, Plant to plant = 45cm, No. of row in a plot = 2, No. of plant in a row = 4, No. of plant in a plot = 8, Distance between two plot in a block = 0.5 m, Block to block = 1 m, Net area = 90 m². There were 10 Treatment including

untreated control plot $T_0 - N_0P_0K_0$, $T_1 - N_1P_1K_1$ (RD 120 : 60 : 60 Kg/ha), $T_2 - N_2P_2K_2$ (above RD 150: 75: 75 Kg/ha), $T_3 - N_3P_3K_3$ (below RD 90: 45: 45: Kg/ha), $T_4 - N_2P_1K_1$ (only N is high and rest were given at RD 150: 60: 60 Kg/ha), $T_5 - N_1P_2K_1$ (only P is high and rest were remain unchanged with 120:75: 60 Kg/ha), $T_6 - N_1P_1K_2$ (only K is above the RD and rest were kept at RD with 120:60:75 Kg/ha), $T_7 - N_3P_1K_1$ (Only N is below RD and rest were kept at RD with 90:60:60 Kg /ha) $T_8 - N_1P_3K_1$ (Only P was given below RD and rest were kept at RD) $T_9 - N_1P_1K_3$ (Only K is below RD and rest were given at RD).

2.5 Assessment of Disease Severity

The number of plants exhibiting symptoms at 60, 90, and 120 DAT were counted in order to determine the disease severity. Each plot's plants were randomly chosen from among them, and the five most common ones were evaluated and assessed for disease severity using the formula provided by [17,18].

$$PDI = \frac{\text{(Sum of each plant disease rating)}}{\text{(Total no. of plant assess} \times \text{maximum disease rating plant)}} \times 100$$

Where PDI = Percent Disease Index

2.6 Statistical Analysis

ANOVA (analysis of variance) was used to statistically examine the experimental results [19]. By applying the Fisher Schedecor "F" tests of probability at the 5% level of significance, error mean square was used to examine the importance of various causes of variation.

3. RESULTS AND DISCUSSION

3.1 Symptoms of the Disease

The characteristic symptom of the disease was observed as circular, sunken and elliptical or oblong spots with black margins on chilli fruits. Badly infected fruits lost their normal colour and turned straw colour. In advance stages of the disease, concentric markings with black fructifications representing the fungal acervuli were observed. The spotted fruits drop off prematurely resulting reduction in yield. In severe cases, it also attacked the fruit stalk and spread along the stem causing die-back symptom. The spots on the fruits measured 5-10 mm in diameter and 10-45 mm in length (Plate 1).

In the field studies, the disease symptoms were observed as small, black, circular lesions on the skin of the fruits which spread towards the long axis giving elliptical shape. As the infection advanced, the spots turn either diffused or black and spread along the stem causing die-back symptoms. Similar observations of the symptoms were also reported by many research workers like Butler and Bisby [21], Chowdhury [22] and Rangaswami [23].

Chowdhury, [22] described the symptoms incited by *Colletotrichum gloeosporioides* on ripe chilli fruit, which appeared as sunken circular or angular lesions. Often multiple lesions coalesce to form severe fruit rot. Generally, the lesions were characterized by the presence of black colored spots in concentric rings at maturity. The dark spots when observed under microscope were the acervuli structure containing setae hairs entrapping the conidia of the pathogen. Further, the pathogen forms micro sclerotia in plant debris or seed, soil, which is the mode of survival under unfavourable conditions.

3.2 Isolation and Identification of Pathogen

The pathogen causing the anthracnose, fruit rot and die-back of King chilli was isolated and identified following the procedure described in materials and methods. The fungal colony appeared as circular and dense, white at first and grey at the later stage conidial masses appeared pink as salmon in colour (Plate 2). Conidia were found to be cylindrical, oblong, -acicular, single celled and hyaline measuring 18.23-25.78 x 6.45-9.89 µm. Conidiophores were hyaline faintly brown, cylindrical, septate or aseptate. Setae were dark, brown, rigid, septate and tapering towards the apex, straight and curved measuring 51.56-128.90 x 4.29-6.45µm (Plate 3).

Based on the above characteristic features the pathogen was identified as the fungus *Colletotrichum gloeosporioides*. (syd.) Butler and Bisby. These observations and finding were similar to those of plant pathologist like Butler and Bisby [21] and Rangaswami [23].

3.3 Field Trial

3.3.1 Effect of different levels of NPK on severity of anthracnose of king chilli

Data on the disease severity as influenced by the treatment of NPK were depicted in Table 2.

Observations were recorded on 60 DAT, 90 DAT and 120 DAT. The results of this study revealed that there was minimum disease severity on $T_6 - N_1P_1K_2$ and disease severity has been observed at 30 days interval with 15.16% (60 DAT), 16.16% (90 DAT) and 19.66 (120 DAT) % which was followed by $T_5 - N_1P_2K_1$ with 19.64 %, 21.33 % and 25.00 % at 60, 90 and 120 DAT respectively. The maximum disease severity was observed in untreated control plot (T_0) with 51.27 % (60 DAT), 55.18 % (90 DAT) and 60.00 % (120 DAT) which was followed by $T_4 - N_2P_1K_1$ with 45.00%, 52.66 % and 58.00 % at 60, 90 and 120 DAT respectively. So, application of NPK with increased level of P and K has paramount significance in combating anthracnose in King chilli.

Nitrogen is an important fertilizer and is essential for cellular components [24]. However, increased dose of Nitrogen (N_2) increased in anthracnose severity [25]. Also increased dose of Nitrogen is known to promote succulent growth and also make the cell walls thinner [26].

Other studies have shown correlations between Nitrogen concentration and severity of anthracnose, in case of chilli. Choi et al. [27] showed that, as tissue Nitrogen content decreased, Ca and Mn contents, which have a direct role in phenolic metabolism, increased in the chilli.

All the treatments of 'Phosphorous' reduced anthracnose severity as compared to control but maximum reduction in disease incidence was in case of P_2 because it increased disease resistance by stimulating the defense system of the plant as it plays a vital role in different types of physiological and biochemical process [28-30].

Potassium may increase synthesis of host defense products and thereby reduce disease severity while phosphorous has been shown to inhibit anthracnose (caused by *C. gloeosporioides* or *C. acutatum*) in chilli [31] and decrease postharvest disease development on chilli.

Table 1. (0 – 9 scale) for accessing PDI of anthracnose of king chilli [20]

Rating scale	Description	Degree of Resistance
0	No symptoms on the leaf or branch or fruits	Immune
1	Small, irregular brown spots covering 1% or less area of the leaf or branch or fruits	Resistant
3	Brown, dirty, pin headed spots covering 1-10% area on the leaf or branch or fruits	Moderately
5	Dark brown, dirty black margin covering 1% of the area of the leaf, or branch or fruits	Moderately susceptible
7	Dark brown, circular or irregular spots with blackish covering 26-50% area of leaf or branch or fruits.	Susceptible
9	Dark brown, circular irregular spots with blackish covering 50 % and above area of leaf or branch or fruits.	Highly susceptible



(a)



(b)



(c)

Fig. 1(a-c). Symptoms of the disease

Table 2. Effect of different levels of NPK against severity of anthracnose on king chilli

Treatment	Anthracnose disease severity (%)		
	60 (DAT)	90 (DAT)	120 (DAT)
T ₀ – N ₀ P ₀ K ₀	51.28 (45.73)	55.19 (47.96)	60.00 (50.77)
T ₁ – N ₁ P ₁ K ₁	35.33 (36.47)	48.14 (43.93)	55.00 (47.87)
T ₂ – N ₂ P ₂ K ₂	31.00 (33.83)	36.00 (36.65)	38.67 (38.26)
T ₃ – N ₃ P ₃ K ₃	26.74 (31.14)	30.33 (33.42)	35.33 (36.77)
T ₄ – N ₂ P ₁ K ₁	45.00 (42.13)	52.67 (46.53)	58.00 (49.51)
T ₅ – N ₁ P ₂ K ₁	19.65 (26.31)	21.33 (27.54)	25.00 (30.00)
T ₆ – N ₁ P ₁ K ₂	15.17 (22.91)	16.17 (23.71)	19.67 (26.32)
T ₇ – N ₃ P ₁ K ₁	22.67 (28.43)	26.00 (30.67)	28.00 (32.00)
T ₈ – N ₁ P ₃ K ₁	38.67 (38.44)	41.00 (39.81)	44.50 (41.83)
T ₉ – N ₁ P ₁ K ₃	41.81 (40.28)	43.00 (45.50)	46.33 (46.00)
SEm±	0.52	0.32	0.19
CD (P=0.05)	1.56	0.95	0.57

Note: Figures in the table are mean values and those in parentheses are arc sine transformed values

The results obtained in the present investigation was also found in agreement with the findings of Gaddam and Sobita [32] who reported that minimum disease intensity was recorded in T₁ (NPK at above the recommended doses - 26.13%) followed by T₅ (P and K above the recommended doses- 26.40%). Maximum disease severity was recorded in T₄ (NPK at very low recommended doses 37.77%) as compared to control. Chowdhury and Rahim [33] also reported that combined application of NPK, Gypsum and Zinc sulphate (250 g + 250 g + 350 g + 100 g + 10 g) reduced the anthracnose severity in chilli (23.33%) and the highest disease severity (64%) was found in T₁₂ (control).

The reduction of disease severity in the present investigation with increased doses of P and K may be attributed to the increase disease resistance by stimulating the defense system of the plants. It is also known that appliances of P and K reduced plant diseases directly by different ways viz., by affecting pathogen metabolism, survival and development; by creating hindrance in food supply to pathogen by affecting internal metabolism of the plants and by disturbing the mode of survival and spread of the pathogen by changing the cell wall ultra-structure and functions of stomata [34,26].

3.3.2 Effect of different levels of NPK in different growth parameters of King chilli

3.3.2.1 Plant height

Data on plant height as influenced by the treatment was presented in Table 3. The observations were recorded at 30, 60, and 90

DAT. The maximum plant height was observed in T₂ – N₂P₂K₂ with 35.00 cm, 45.16 cm, and 85.83 cm respectively. This was followed by T₆ – N₁P₁K₂ with 16.66 cm, 26.83 cm, and 80.00 cm respectively. The minimum height was recorded at T₀ (untreated plot) with 10.17 cm, 20.50 cm, and 55.33 cm on 30, 60, and 90 DAT respectively.

From the above differences in plant height, it was discovered that treatment T₂ – N₂P₂K₂ had the tallest plant height (85.83 cm), followed by treatment T₆ – N₁P₁K₂ with (80.00 cm), and treatment T₉ – N₁P₁K₃ had the shortest plant height among the treatments with 59.90 cm. This might be as a result of increased NPK availability and uptake, which gradually increased plant height.

The present finding was found in agreement with the earlier report made by Gaddam and Sobita [32] who reported that the maximum plant height in T₁ (NPK at 150; 50; 50 Kg/ha – 54.66 cm) followed by T₂ (NPK at 250; 100; 100 Kg/ha – 51.10cm). While the untreated control found the shortest plant height (35.54cm). According to Adeola et al. [35] the effect of different levels of NPK considerably influenced the plant heights of the chilli, which dramatically ranged from 51.00 to 73.00 cm.

Dubey [36] measured the maximum plant height at 30, 60, 90, 120, and 150 days after planting in T₇: NPK@ 175:55:45 Kg/ha with (35.76 cm, 50.10 cm, 66.73 cm, 89.10 cm, and 116.70 cm respectively), followed by T₉: NPK @ 175:70:55 Kg/ha with (31.03 cm, 45.37 cm, 61.97 cm, 84.37 cm, and the minimum was recorded in T₁ (control) NPK@ 135:40:35 Kg/ha with (18.46cm,

32.87cm, 49.47cm, 71.77cm and 99.30cm respectively. He observed that plant height rose progressively as NPK levels rose.

3.3.2.2 Number of fruits per plant

Data pertaining to number of fruits as influenced by the treatment was presented in Table 4. The observations were recorded at 90 DAT.

The Maximum Number of fruits was recorded in $T_5 - N_1P_2K_1$ with 85 fruits per plant which was followed by $T_6 - N_1P_1K_2$ with 80 fruits per plant and $T_0 - N_0P_0K_0$ plot was observed to give the minimum no. of fruits per plant. Hassan and Kamal Uddin [37] who revealed that number of hybrid fruits per plant was significant due to the combined fertilizer application of NPK. The number of fruits plant⁻¹ was statistically maximum with the application of NPK 100% with 56.64 fruits plant⁻¹ as compared to control with 24.26 fruits plant⁻¹ (without NPK). The results presented here support those of Johnson and Decoteau [38], who discovered that fruit count and weight per plant increased linearly with increasing N rate as well as increased disease resistance in chilli crops.

3.3.2.3 Yield

Data pertaining to fruit yield per plant (fresh weight in gm), per plot (weight in kg) and per hectare (weight in ton) as influenced by the treatment was presented in Table 5. The observations were recorded at 90 DAT.

The maximum yield of fruits per plant, per plot and per hectare was recorded in $T_5 - N_1P_2K_1$ with

853.33 g plant⁻¹, 11.17 Kg plot⁻¹ and 37.00 ton hectare⁻¹ followed by $T_6 - N_1P_1K_2$ with 800.00 g plant⁻¹, 9.30 Kg⁻¹ plot and 31.00 ton hectare⁻¹. Conversely, the minimum yield was obtained at $T_0 - N_0P_0K_0$ with 205.00 gm plant⁻¹, 1.65 Kg plot⁻¹ and 5.167 ton hectare⁻¹. The aforementioned conclusion is consistent with that of Ayodele et al. [39]. Similar findings were also reported by Hasan and Kamal Uddin [37], who found that the effect of NPK fertilizer on chilli hybrid plants caused a substantial variation in seed output, ranging from 14.33 to 40.27 Kg ha⁻¹.

The combination of N and P fertilizers and different level of N fertilizers of the plant had found no significant difference. Significant increased of plant fresh weight was registered at P_{60} Kg/ha (72.20g). However control (P_0 Kg/ha) treatment had relatively lowest plant fresh weight accumulation than other treatments [38].

Table 3. Effect of different levels of NPK on plant height of King chilli

Treatment	Plant height in cm		
	30 DAT	60DAT	90DAT
$T_0 - N_0P_0K_0$	10.17	20.50	55.33
$T_1 - N_1P_1K_1$	20.17	30.00	73.00
$T_2 - N_2P_2K_2$	35.00	45.17	85.83
$T_3 - N_3P_3K_3$	15.17	17.17	66.80
$T_4 - N_2P_1K_1$	25.33	33.17	69.00
$T_5 - N_1P_2K_1$	29.50	40.33	75.47
$T_6 - N_1P_1K_2$	16.67	26.83	80.00
$T_6 - N_1P_1K_2$	12.83	23.00	70.67
$T_8 - N_1P_3K_1$	22.83	43.17	63.23
$T_9 - N_1P_1K_3$	32.17	39.17	59.90
SEm±	0.16	0.08	0.21
CD (P=0.05)	0.68	0.50	0.79

Table 4. Effect of different levels of NPK on no. of fruit and yield of king chilli

Treatment	No.of fruit /Plant	Yield Of fresh weight (g) /Plant	Yield per plot (Kg)	Yield (ton)/ hectare
$T_0 - N_0P_0K_0$	20.00	205.00	1.65	5.16
$T_1 - N_1P_1K_1$	52.00	524.33	5.03	16.41
$T_2 - N_2P_2K_2$	75.00	755.00	7.07	23.18
$T_3 - N_3P_3K_3$	46.33	458.33	3.72	12.16
$T_4 - N_2P_1K_1$	70.00	712.00	4.04	13.00
$T_5 - N_1P_2K_1$	85.00	853.33	11.17	37.00
$T_6 - N_1P_1K_2$	80.00	800.00	9.30	31.00
$T_7 - N_3P_1K_1$	40.16	410.00	6.16	20.16
$T_8 - N_1P_3K_1$	36.00	365.00	2.30	7.56
$T_9 - N_1P_1K_3$	30.00	301.66	2.43	8.40
SEm±	0.04	31.85	0.47	0.05
CD (P=0.05)	0.35	9.68	0.07	0.38




Plate 1. (a), (b), (c), (d), (e), (f), (g), (h), (i) and (j) are the symptoms of anthracnose caused by *Colletotrichum gloeosporioides* on leaf, branches, twigs and fruit of king chilli plant

Table 5. Results of regression showing average effect of PDI on different growth parameters of king chilli

Growth parameters		Regression coefficients	SE	t	p - value	R2
Plant height	Intercept	48.00	15.14	3.17	< 0.01	0.01
	PDI	0.08	0.41	0.2	0.84	
No.of fruit /Plant	Intercept	98.64	6.01	16.41	< 0.01	0.70
	PDI	-1.19	0.15	-8.09	< 0.01	
Yield Of fresh weight /Plant	Intercept	989.42	60.66	16.31	< 0.01	0.70
	PDI	-11.86	1.48	-8.00	< 0.01	
Yield per plot	Intercept	10.73	1.05	10.22	< 0.01	0.53
	PDI	-0.14	0.02	-5.57	< 0.01	
Yield/ hectare	Intercept	35.64	3.45	10.35	< 0.01	0.54
	PDI	-0.48	0.08	-5.70	< 0.01	



Plate 3. Conidia of *Colletotrichum gloeosporioides* under 40X compound microscope (cylindrical, hyaline, septate structure)



Plate 2. Growth of *Colletotrichum gloeosporioides* in PDA

The regression result as given in Table 5 shows there was no significant correlation between disease severity and plant height. As only 1

percent of the variation in plant height of King chilli can be predicted by disease severity. In case of regression analysis between disease severity and no. of fruits/plant and yield of fresh weight/ plant, there was significant negative correlation between no. of fruits/plant and yield of fresh weight/ plant and disease severity of Anthracnose and 70% of the variation in yield of King chilli can be predicted by PDI of the disease.

4. CONCLUSIONS

With the above given findings on the present investigation, it can be concluded that application of 'P' and 'K' at higher or recommended doses were effective in inhibiting anthracnose disease severity on King chilli. Application of NPK at above the normal recommended doses gave better plant height of King chilli. Furthermore, application of P & K above the normal recommended dosage produced maximum number of fruits per plant and gave better yields that were significantly higher than rest of the treatments. Application of increased level of P and K has therefore paramount importance in combating anthracnose disease severity of King chilli.

COMPETING INTERESTS

Authors have declared that no competing interests exist."

REFERENCES

1. Berke TG, Shieh SC. Capsicum, chillies, paprika, bird's eye chilli In: Peter KV (Ed.) handbook of herbs and spices. Wood

- Head Publishing Limited, Cambridge. 2002:111-122.
2. Patel VK, Gupta SP, Patel KL. Economics performance of chilli (*Capsicum Annuum* L.) cultivation in Raigarh district of Chhattisgarh state. International Journal of Agricultural Science and Research (IJASR). 2015;5:363-368.
 3. Abdul Kareem M, Tatagar MH, Krishna D, Kurubetta Mesta R, Masuthi DK, Waseem MA. Evaluation of bio-efficacy and phytotoxicity of pyraclostrobin 20% WG against anthracnose disease chilli. Journal of Pure and Applied Microbiology. 2016;10(3):2397-2399.
 4. Sanatombi KS, Sen-Mandi, Sharma GJ. DNA profiling of capsicum landraces of Manipur. Scientia Horticulturae. 2010;124:405-408.
 5. Verma PK, Rawat KK, Das N, Pradhan B. A botanical enigma of India's hottest chilli 'bhoot jolokia' (*Capsicum Chinense* Jacq.). New York Science Journal. 2013;6(11):49-51.
 6. Bhuyan MH, Rahman MB, Ara Rumman SML, Sarker JC. Evaluation of naga chilli (*Capsicum chinense* Jacq.) genotypes under North Eastern region of Bangladesh. Scientia Agriculturae. 2015;12(1):40-45.
 7. Mathur R, Dangi RS, Das SC, Malhotra RC. Hottest chilli variety in India. Current Science. 2000;79(3):287-289.
 8. Issac S. Fungal plant interaction. Chapman and Hall Press, London. 1992;115.
 9. Manandhar JB, Hartman GL, Wang TC. Anthracnose development on pepper fruits inoculated with *Colletotrichum gloeosporioides*. Plant Disease. 1995;79:380-383.
 10. Saxena AK, Hoshi IJ, Saxena JB. Additions to the host range of *Colletotrichum capsici*. India Phytopathology. 2016;35:166-212.
 11. Park KS, Kim C. Identification, distribution, and etiological characteristics of anthracnose fungi of red pepper in Korea. Korean Journal of Plant Pathology. 1992;8:61-69.
 12. Babu BS, Pandravada SR, Rao SK, Varaprasad KS. Global sources of pepper genetic resources against arthropods, nematodes and pathogens. Crop protection. 2011;30(5):389-400.
 13. Voorrips RE, Finkers R, Sanjaya L, Groenwold R. QTLmapping of anthracnose (*Colletotrichum* spp.) resistance in a cross between *Capsicum annuum* and *C. chinense*. Theoretical and Applied Genetics. 2004;109:1275-1282.
 14. Malangmeih, Rahaman SM. Economics of fresh naga king chilli in Manipur, India –a case study. International Journal of Environment, Ecology. 2016;6:151-162.
 15. Mew TW, Gonzales P. A handbook of rice seed-borne fungi. Los Banos, Philippines: International Rice Research Institute (IRRI) and Enfield, N.H., USA: Science Publishers, Inc. 2002:83.
 16. Barnett HL, Hunter BB. Illustrated genera of imperfect fungi. Burgess Publishing Company, Minnesota, USA; 1972.
 17. Chester KS. Plant disease losses: Their appraisal and interpretation. Plant Disease Reporter Supplement. 1959;193:189-362.
 18. Wheeler GC, Wheeler J. The larva of *Acanthognathus* (Hymenoptera: Formicidae). Psyche (Cambridge). 1969; 76:110-113.
 19. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. Indian Council of Agricultural Research Publication. 1978:87-89.
 20. Mayee CD, Datar VV. Phytopathometry. Marathwada Agricultural University Tech. Bul. 1(special Bul. 3); 1986.
 21. Butler EJ, Bisby GR. The fungi of India. Imperial Council of Agricultural Research Science Monograph-I. 1931;18: 237-241.
 22. Chowdhury S. Studies on the development and control of fruit rot of chillies. Indian Phytopathology. 1957;10:55-62.
 23. Rangaswami G. Disease of crop plants in India, Printice-Hall of India Private Ltd. New Delhi, India. 1994:570.
 24. Huber DM, Thompson IA. Nitrogen and plant disease. In: Datnoff LE, Elmer WH, Huber DM (ed) .Mineral nutrition and plant disease, APS Press, St Paul. 2007:31-44.
 25. Huber DM. The role of mineral nutrition in defense. In: Harsfall JG, Cowling EB. (ed). Plant disease, an advanced treatise how plants defend themselves, Academic Press, New York. 1980:381-406.
 26. Atiq Muhammad, Nazir Javed. Management of leaf rust of wheat through different levels of NPK and sowing times. Advances in Zoology and Botany. 2017;5(4):39-44.
 27. Choi JM, Jeong SK, cha KH, Chung HJ, Seo KS. Deficiency symptoms, growth characteristics and nutrient uptake of 'Nyoho' chilli as affected by controlled nitrogen concentrations in fertilizer

- sololution. Journal of the Korean society for Horticultural Science. 2000;41:339-441.
28. Khan R, Gurmani AR, Gurmani AH, Zia MS. Effect of phosphorus application on wheat and rice yield under wheat- rice system. Sarhad Journal of Agriculture. 2007;23(4):851-856.
 29. PPI. Effects of potassium on plant diseases. In: Potassium for Agriculture, Potash & Phosphate Institute, Norcross, GA. 1998:37-39.
 30. Snyder CS, Mascagni HJ. Phosphorus and potassium increase wheat yields and help reduce disease damage [On-line]. Int. Plant Nutrition Institute. News and Views from the Southeast Region; 1998. Available:<http://www.ipni.net/news>
 31. Biggs AR, Peterson CA. Effect of chemical applications to peach bark wounds on accumulation of lignin and suberin and susceptibility to *Leucostoma persoonii*. Phytopathology. 1999;80(86):1-5.
 32. Gaddam P, Sobita S. Integrated approaches for management of anthracnose of chilli (*Capsicum annum* L). Journal of Pharmacognosy and Photochemistry. 2019;8(1):422-427.
 33. Chowdhury, Rahim. Effect of fertilizers on the incidence of anthracnose disease, yield and quality of chilli. Journal of Agriculture & Rural Development. 2007;5(1&2):70-78.
 34. Perrenoud S. Potassium and Plant Health, 2nd ed. International Potash Institute, Bern, Switzerland. 1990:368. IPI Research Topics No. 3
 35. Adeola RG, Tijani EH, Makinde, EA. Ameliorate the effects of poultry manure and NPK fertilizer on the performance of pepper cropped with two cassava varieties. Global Journal of Science Frontier Research. 2011;11:661-669.
 36. Dubey AK. Effect of NPK on plant growth, yield and quality of capsicum (*Capsicum annum* L.) C.V. Swarna Under Shade Net Condition. International Journal of Current Microbiology and Applied Sciences. 2017;6(3):1085-1091.
 37. Hassan M, Kamal Uddin MD. Impact of nitrogen and phosphorus fertilizer on growth and yield of Bambara groundnut. Plant Archives. 2019;19:501-504.
 38. Johnson CD, Decoteau DR. Nitrogen and potassium fertility affects jalapeno pepper plant growth, pod yield and pungency. Journal of Horticultural Science. 1996;31(7):1119-1123.
 39. Ayodele OJ, Alabi EO, Aluko M. Nitrogen fertilizer effects on growth, yield and chemical composition of chilli hot pepper (Rodo). International Journal of Agriculture and Crop Sciences. 2015; 8(5):666-673.

© 2022 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/94278>